ELSEVIER

Inorganica Chimica Acta 283 (1998) 281–291

Inorganica Chimica Acta

Cumulative Author Index of Volumes 267–283

Abbott, E.H., **268**, 311
Abboud, K.A., **282**, 90
Abd-el-Khalek, H., **275–276**, 1
Abney, K.D., **274**, 232
Abram, S., **278**, 118
Abram, U., **271**, 199
Adams, E.K., **281**, 43
Adams, M.W.W., **273**, 409
Addison, A.W., **278**, 217
Adrowski, M.J., **278**, 113
Agterberg, F.P.W., **267**, 183
Aguiar, A.P., **273**, 196
Aguilera, F., **274**, 15
Aguirre, G., **274**, 108
Ahmed, L.S., **278**, 229
Aika, K., **280**, 183
Aikawa, M., **282**, 38
Aime, S., **275–276**, 521
Ainscough, E.W., **267**, 27; **277**, 37
Airola, K., **277**, 55
Akashi, H., **282**, 50
Akazawa, M., **271**, 13
Akhmedov, N.G., **280**, 328
Akhrem, A.A., **275–276**, 510
Akhrem, I., **280**, 355
Akizawa, H., **283**, 44
Akrivos, P.D., **271**, 57, 195; **281**, 121
Alarcón, C.J., **278**, 61
Albano, G., **277**, 225
Albinati, A., **274**, 219
Albright, T.A., **272**, 95
Alessio, E., **273**, 62
Alguindigue, S.S., **270**, 227
Alkam, H., **271**, 195
Allevi, M., **282**, 17
Alonso, G., **274**, 108
Alper, H., **280**, 172
Alzuet, G., **273**, 334
Amadelli, R., **272**, 197
Amador, U., **279**, 30
Amao, Y., **267**, 257
Amatore, C., **273**, 76
Amorim, M.T., **278**, 51
Amstutz, N., **271**, 231
Anacardio, R., **282**, 119
Anastassopoulou, J.D., **273**, 201
Anderegg, G., **274**, 219
Andersen, B., **273**, 41
Andersen, N.H., **272**, 109
Anderson, G.K., **271**, 236
Ando, I., **282**, 247
Ando, Y., **274**, 184
Anesti, V., **269**, 332
Angermund, K., **270**, 273
Angus, P.M., **268**, 85
Anis, S., **278**, 185
Antipin, M.Yu., **280**, 87, 288
Antonietti, M., **280**, 348
Antonov, D.Yu., **278**, 122
Antov, L., **272**, 291
Aoki, K., **277**, 8; **281**, 36
Aoki, M., **272**, 80
Aoki, S., **270**, 424
Aoyama, Y., **275–276**, 159
Appleton, T.G., **279**, 152
Aquino, M.A.S., **279**, 121
Arakawa, R., **279**, 165
Arancibia, A., **269**, 246
Arena, C.G., **277**, 119
Arena, G.E., **275–276**, 242
Aresta, M., **272**, 38
Argazzi, R., **275–276**, 380
Arion, V.B., **282**, 61
Armelao, L., **275–276**, 340
Arriortua, M.I., **278**, 150
Arulsamy, N., **274**, 155
Asano, H., **270**, 87
Ascenso, C., **273**, 279
Ascenzi, P., **267**, 109
Aschi, M., **275–276**, 192
Ashby, M.T., **270**, 227
Ashfar, A., **279**, 152
Aslanidis, P., **271**, 243; **281**, 121
Atencio, R., **275–276**, 528
Attinà, M., **275–276**, 192
Atwood, D.A., **277**, 157
Aukauloo, A., **278**, 159
Avnir, D., **280**, 21
Ayala, J.D., **274**, 122, 236
Ayres, D.J., **277**, 177
Azuma, N., **274**, 210; **281**, 207; **283**, 61
Azuma, S., **271**, 24

Babini, E., **275–276**, 230
Bachmann, C., **274**, 143
Baena, M.J., **273**, 244
Baggio, R., **278**, 108; **281**, 18
Baker, P.K., **279**, 65
Bakhmutov, V., **270**, 169
Bakhmutov, V.I., **278**, 122; **280**, 302
Bakhmutova, E.V., **280**, 302
Bakir, M., **270**, 511
Bal, W., **275–276**, 546; **283**, 1
Balahura, R.J., **274**, 215
Balch, A.L., **270**, 28
Balt, S., **269**, 203
Balzani, V., **277**, 225
Banci, L., **275–276**, 373
Bandoli, G., **269**, 235; **275–276**, 340, 385, 401; **277**, 163, 247; **278**, 96
Banerjee, R., **270**, 392
Banerjee, S., **275–276**, 551
Banße, W., **269**, 350
Baran, E.J., **268**, 239
Baran, P., **278**, 190
Barbier, J.-P., **268**, 13
Bardwell, D.A., **267**, 239, 323
Barnham, K.J., **273**, 8
Baron, A.J., **275–276**, 175
Barral, M.C., **277**, 76; **279**, 30
Barrera, J., **270**, 337
Barroso, N.P., **273**, 191
Bartczak, T.J., **277**, 242
Barthram, A.M., **267**, 1
Bartoli, J.-F., **272**, 4
Barton, S.J., **273**, 8
Barybin, M.V., **269**, 58
Bashall, A., **273**, 372
Basosi, R., **275–276**, 274, 557
Bastida, R., **267**, 55; **282**, 42
Bastos, C.M., **279**, 7
Baştürkmen, M., **271**, 19
Batista, A.A., **267**, 209
Battaglia, L.P., **278**, 209
Battioni, P., **272**, 4
Battistuzzi, G., **272**, 168; **273**, 397
Battistuzzi, R., **270**, 145
Bau, R., **269**, 162
Bavoso, A., **268**, 205
Bazhenova, M.A., **270**, 221
Bazhenova, T.A., **270**, 221; **280**, 288
Bazzicalupi, C., **268**, 63
Beauchamp, A.L., **279**, 159
Beck, C.M., **270**, 559
Beck, E.J., **279**, 121
Beck, S., **270**, 376
Beckett, M.A., **279**, 65
Beddoes, R.L., **273**, 225
Bedgood Jr., D.R., **282**, 200
Behrens, P., **269**, 73
Belicchi Ferrari, M., **269**, 297; **275–276**, 320
Belkova, N.V., **280**, 302
Bell, N.A., **277**, 14
Bell, R.A., **274**, 137
Belluco, U., **275–276**, 515
Belyakova, O.A., **280**, 275
Bencini, A., **268**, 63; **273**, 326; **274**, 90
Benetollo, F., **280**, 308
Bennett, J.E., **271**, 167
Bennett, J.L., **270**, 414
Bensimon, C., **280**, 172
Bercaw, J.E., **270**, 433, 467; **280**, 226
Berchet, L., **282**, 243
Bereman, R.D., **282**, 200
Berg, A., **275–276**, 493
Bergamo, A., **273**, 62
Berghoff, U., **269**, 135
Bergman, R.G., **270**, 20
Berkovich, E.G., **280**, 264
Bernadou, J., **272**, 228
Bernal, I., **282**, 96
Bernardinelli, G., **271**, 231; **275–276**, 279
Bernauer, K., **275–276**, 1, 9
Bertani, R., **275–276**, 515
Berthon, G., **283**, 211
Bertini, I., **269**, 125; **283**, 12
Bertolasi, V., **272**, 267
Besançon, J., **268**, 323
Beuter, M., **275–276**, 203
Beyer, L., **271**, 40
Bhalla, R., **273**, 225
Biagini-Cingi, M., **273**, 259
Bianchi, A., **273**, 326
Bianchi, M., **272**, 141
Bianchini, C., **272**, 1, 55; **280**, 302
Bickelhaupt, F., **269**, 203
Bignozzi, C.A., **275–276**, 380
Bigoli, F., **267**, 115; **273**, 175
Bikrani, M., **282**, 230
Bill, E., **282**, 61
Birnbaum, E.R., **270**, 433
Blaauw, R., **269**, 203
Blackman, A.G., **277**, 89
Blaton, N.M., **269**, 332
Blum, J., **280**, 21
Boča, M., **278**, 190
Boca, R., **278**, 127, 190
Boccarelli, A., **273**, 62
Bocelli, G., **270**, 123
Bochkarev, M.N., **280**, 138

Boersma, M.G., **275–276**, 98
Boffi, A., **267**, 109
Bögge, H., **271**, 9; **280**, 30
Bóka, B., **275–276**, 440
Bombieri, G., **274**, 122, 236; **280**, 308
Bommeli, H., **274**, 219
Bond, A.M., **267**, 281
Bonnamartini Corradi, A., **272**, 252
Bönnemann, H., **270**, 95
Bonomo, R.P., **282**, 180
Borbulevitch, O.Ya., **280**, 87
Borchert, T., **274**, 201
Bordunov, A.V., **278**, 6
Borgherini, F., **268**, 323
Borrás, J., **273**, 334
Borsari, M., **270**, 145; **272**, 168; **273**, 397; **275–276**, 230
Borzov, M.V., **280**, 257
Bosch, B., **270**, 446
Bösing, M., **274**, 251
Bosnich, B., **270**, 13
Bott, S.G., **267**, 223; **274**, 232
Böttcher, R., **281**, 160
Boubekeur, K., **279**, 144
Boucher, K.D., **267**, 159
Bouillez, A., **275–276**, 528
Bouwman, E., **279**, 217
Boyer, M.T., **270**, 8
Boyle, P.D., **282**, 200
Bradshaw, J.E., **275–276**, 106
Bradshaw, J.S., **278**, 6
Braga, D., **273**, 116
Brandt, K., **267**, 39
Brassat, I., **280**, 150
Brauch, T.W., **270**, 285
Brazzini, S., **273**, 326
Bregadze, V.I., **280**, 233
Breulmann, M., **280**, 348
Brewer, C.T., **278**, 197
Brewer, G., **278**, 197; **279**, 111
Brianese, N., **272**, 235
Briganti, F., **275–276**, 295
Brijoux, W., **270**, 95
Brinkmann, R., **270**, 95
Brintzinger, H.H., **270**, 376; **280**, 189
Brničević, N., **271**, 239
Brodie, A.M., **267**, 27; **277**, 37
Bronstein, L.M., **280**, 348
Brooker, S., **282**, 222
Broomhead, J.A., **271**, 151
Broughton, V., **275–276**, 279
Brown, D.A., **268**, 313; **280**, 30
Brown, K.L., **267**, 305; **279**, 178
Brown, T.M., **267**, 101
Browning, C.S., **271**, 112
Bruce, G.S., **275–276**, 250
Brüdgam, I., **269**, 23
Brüggeller, P., **274**, 143
Brügger, N., **275–276**, 1
Bruk, L.G., **280**, 202
Bruno, G., **277**, 119
Bruno, N., **271**, 217
Bryant, R.G., **275–276**, 106
Bubner, M., **273**, 234
Bucke, L.C., **277**, 89
Budker, V., **273**, 354
Buglyó, P., **268**, 297
Buijs, W., **267**, 183
Buisson, Y.L., **268**, 271
Bujdak, J., **272**, 89
Bullen, H.A., **278**, 241
Burdeniuc, J., **270**, 559
Burewicz, A., **271**, 167
Burger, K., **274**, 167; **275–276**, 130
Burrow, R.A., **271**, 112
Burrows, C.J., **275–276**, 314
Burth, R., **282**, 193
Busch, D.H., **270**, 151
Butenschön, H., **280**, 71
Buzás, N., **274**, 167

Cabeza, J.A., **268**, 131
Cacace, F., **275–276**, 192
Cafeo, G., **275–276**, 141
Calcaterra, A.D., **267**, 101
Calderazzo, F., **270**, 177
Calhorda, M.J., **275–276**, 263
Calogero, S., **282**, 1
Camm, G., **271**, 151
Campagna, T., **275–276**, 395
Cámpora, J., **269**, 191
Camus, A., **278**, 237
Cano, A.M., **280**, 1
Cano, J., **280**, 1
Canovese, L., **275–276**, 385; **277**, 247
Canters, G.W., **275–276**, 493
Cantrell, T.L., **268**, 211
Canty, A.J., **267**, 281
Cao, R., **267**, 313; **270**, 247; **271**, 93; **274**, 229; **275–276**, 557; **277**, 171
Capdevila, M., **278**, 10
Capitani, D., **282**, 17
Capozzi, F., **275–276**, 230
Capparelli, M., **275–276**, 528
Capps, K.B., **270**, 488
Carassiti, V., **272**, 197
Cargill Thompson, A.M.W., **267**, 239
Carlson-Day, K.M., **271**, 49
Carmona, E., **269**, 191; **273**, 244
Carotti, S., **281**, 90
Carrano, C.J., **273**, 14
Carrillo, D., **268**, 199
Carroll, R.I., **270**, 207
Carrondo, M.A., **273**, 213
Caruso, F., **267**, 151
Carvalho, M.F.N.N., **271**, 65
Casamento, M., **275–276**, 242
Casanova, J., **273**, 334
Casarin, M., **275–276**, 340
Casas, J.S., **267**, 169
Casella, L., **273**, 339; **282**, 180
Casellato, U., **272**, 235
Casey, C.P., **270**, 5; **280**, 316
Casey, M.T., **277**, 257; **279**, 24
Castan, P., **271**, 217
Castel, A., **277**, 98
Castiñeiras, A., **282**, 42
Castro, I., **278**, 159
Catellani, M., **270**, 123
Caulton, K.G., **270**, 261; **280**, 125
Cauzzi, D., **273**, 320
Cavaleiro, A.M.V., **281**, 225
Cecconi, F., **269**, 274
Celda, B., **275–276**, 73
Cenini, S., **272**, 18
Černák, J., **281**, 134
Cervera, B., **278**, 159
Cesario, M., **275–276**, 234
Cevik, S., **277**, 69
Chakravorty, A., **275–276**, 551
Challita-Bechara, A., **271**, 210
Chaloner, P.A., **281**, 165
Chan, A.S.C., **270**, 247
Chan, F., **272**, 95
Chan, K.S., **270**, 551
Chan, M.C.W., **272**, 176
Charmant, J.P.H., **270**, 382
Chau, D.E.K.-Y., **270**, 130
Chaudhuri, P., **274**, 111
Chavez, F.A., **269**, 269
Che, C.-M., **272**, 176
Chen, C., **267**, 81
Chen, C.-C., **270**, 247
Chen, F., **273**, 255
Chen, J., **267**, 313
Chen, J.-D., **269**, 302; **274**, 7
Chen, L., **267**, 271
Chen, Q., **268**, 163; **277**, 69
Chen, S., **283**, 17
Chen, Y., **267**, 11; **268**, 279, 287; **275–276**, 58; **277**, 46; **279**, 85
Chen, Y.-W., **275–276**, 215
Chen, Z., **273**, 255, 310
Cheng, J.Y.K., **272**, 176
Cherepanov, A., **275–276**, 493
Cherepanov, I.A., **280**, 71
Cherkashina, N.V., **280**, 339
Chernoglazova, N., **280**, 295
Chessa, G., **275–276**, 385
Cheung, K.-K., **272**, 176
Chia, L.-S., **282**, 10
Chibisov, A.K., **271**, 160
Chiesi-Villa, A., **270**, 298
Chikira, M., **279**, 51
Childs, B.J., **274**, 32
Chillemi, F., **273**, 339
Chiotellis, E., **268**, 169; **281**, 148
Chistyakov, A.L., **280**, 219
Chiumia, G.C., **271**, 210
Chiusoli, G.P., **270**, 123
Chizhevsky, I.T., **280**, 233
Chmielewski, P.J., **273**, 184
Chojnacki, J.A., **270**, 405
Chomič, J., **281**, 134
Choppin, G.R., **278**, 185
Choueiry, D., **280**, 8
Christoffers, J., **270**, 20
Chruscinska, E., **269**, 279
Chryssou, K., **268**, 169
Chua-anusorn, W., **267**, 7
Chung, Y.K., **279**, 116, 243; **281**, 229
Churakov, A.V., **280**, 257
Ciccarese, A., **275–276**, 410, 419
Ciofini, I., **274**, 90
Cipollini, R., **275–276**, 192
Clajus, D., **281**, 64
Clark, C.D., **279**, 200
Clark-Baldwin, K., **273**, 85; **275–276**, 215
Clearfiled, A., **270**, 207
Cleary, R.L., **267**, 1
Clegg, W., **271**, 57; **277**, 14
Clemente, D.A., **275–276**, 410, 419
Clouston, J.A., **274**, 215
Coffey, T.A., **274**, 243
Cohen, H., **270**, 440; **273**, 266
Colacio, E., **281**, 95
Cole-Hamilton, D.J., **280**, 99
Coleman, J.E., **275–276**, 481
Cölfen, H., **280**, 348
Collazo, C., **272**, 125
Collison, D., **273**, 225
Collomb-Dunand-Sauthier, M.-N., **282**, 243
Colton, R., **267**, 281
Coluccia, M., **273**, 62
Conan, F., **278**, 108
Conrad, O., **274**, 251
Consiglio, G., **282**, 163
Constable, E.C., **275–276**, 359; **277**, 225
Conte, V., **272**, 62
Cooper, A.C., **270**, 261
Corain, B., **275–276**, 401; **282**, 119
Cornia, A., **280**, 282
Corvaja, C., **275–276**, 401
Costa, C., **275–276**, 256
Costa, G., **270**, 145; **280**, 322
Costa, H.S., **273**, 196
Costa, I., **275–276**, 117
Costa, M.T.A.R.S., **280**, 308
Costamagna, J., **267**, 151; **273**, 191
Costanzo, F., **275–276**, 366
Costes, J.-P., **268**, 125; **274**, 73
Cotton, F.A., **267**, 173, 271; **271**, 187; **275–276**, 447, 541
Couchman, S.M., **267**, 1; **278**, 178
Coucouvanis, D., **283**, 111
Couto, F.A.R.S., **281**, 225
Coutsolelos, A.G., **275–276**, 182
Cowan, J.A., **275–276**, 24
Cox, B.G., **279**, 186
Cox, O., **271**, 137
Cox, P.J., **271**, 243
Crabtree, R.H., **270**, 559; **280**, 26
Craig, D.C., **268**, 135; **271**, 210; **274**, 32
Creighton, J.R., **271**, 57, 195; **277**, 14
Cresswell, R.J., **277**, 37
Crevier, T.J., **270**, 202
Crites, D.K., **273**, 346
Crociani, L., **282**, 119
Cronin, L., **269**, 241

Crowder, M.W., **268**, 211
Cucurull-Sanchez, L., **271**, 57; **277**, 14
Cuenca, T., **280**, 1
Cugini, F., **270**, 123
Cui, Y., **277**, 130; **282**, 261
Cummins, C.C., **278**, 103
Cunningham, C.T., **273**, 346
Curran, M., **277**, 257
Curtin, L.S., **282**, 155
Cyr, P.W., **270**, 130
Czernuscewicz, R.S., **273**, 14
Czernuszewicz, R.S., **281**, 25
Czerwinski, C.J., **280**, 316
Czuchajowski, L., **271**, 167

D'Agostino, A., **267**, 281
Dahan, F., **274**, 73; **279**, 69
Dahlke, P., **268**, 117
Dahrouch, M., **277**, 98
Dakternieks, D., **275–276**, 203
Dalian, E.B., **275–276**, 510
Dalley, N.K., **282**, 142
Dapporto, P., **275–276**, 168; **282**, 76
Daraghmeh, H., **279**, 127
Darensbourg, D.J., **270**, 405; **274**, 115
Daridor, A., **272**, 188; **280**, 118
Darkwa, J., **267**, 19
Dasgupta, T.P., **268**, 31
Dauter, Z., **273**, 213
Dave, B.C., **281**, 25
Davey, J.M., **281**, 10
Davies, H.O., **268**, 313
Davis, C.A., **273**, 372
Davis, J.J., **272**, 261
Davis, W.M., **267**, 165; **270**, 353
Davison, A., **267**, 165, 319; **271**, 191; **279**, 206
Dawson, J.H., **275–276**, 250
Day, M.W., **270**, 467; **280**, 226
de Blas, A., **267**, 55; **282**, 42
de Bolster, M.W.G., **269**, 203
de Castro, B., **271**, 83
Déchamps-Olivier, I., **268**, 13
Deere, P.T., **281**, 1
de Geest, D.J., **282**, 222
De Gioia, L., **273**, 339, 379
de Graaff, R.A.G., **267**, 193
Deibe, A.M.G., **274**, 73
Dekker, E.E., **275–276**, 215
De la Rosa, M.A., **272**, 109; **275–276**, 73
De Leonardis, P., **273**, 116
Deligeer, **275–276**, 289
del Mar Conejo, M., **272**, 125
DelMedico, A., **281**, 126
Del Pra, A., **274**, 122, 236
Delpuech, J.-J., **275–276**, 130
del Río, I., **268**, 131
Del Zotto, A., **277**, 111
Demadis, K.D., **270**, 511
Dementjev, A.P., **273**, 234
De Montauzon, D., **275–276**, 182
De Munno, G., **282**, 82, 252
Dendrinou-Samara, C., **272**, 24
Denti, G., **275–276**, 320
Depaoli, G., **275–276**, 340
De Pasquale, C., **278**, 76
Deplano, P., **267**, 115; **273**, 175
Deroche, A., **275–276**, 234
Deronzier, A., **282**, 243
De Santis, G., **267**, 177
de Sousa Moreira, I., **267**, 93
Deters, R., **269**, 117
Devereux, M., **277**, 257; **279**, 24
Devillanova, F.A., **267**, 115
De Vries, S., **275–276**, 493
de Witt, G.C., **275–276**, 37
Diaz, A., **275–276**, 557
Díaz, C., **269**, 246
Díaz, L.E., **273**, 367
Di Bilio, A.J., **275–276**, 470
Dick, S., **269**, 47
Di Furia, F., **272**, 62
Dilworth, J.R., **278**, 229
Dinjus, E., **269**, 283
Di Vaira, M., **273**, 151
Dixneuf, P.H., **272**, 188; **280**, 118
Dlháň, Ľ., **278**, 190
Do, Y., **283**, 145
Dobbs, A.J., **267**, 27
Dobos, S., **277**, 61
Dobson, G.R., **267**, 49; **271**, 105; **278**, 202
Doedens, R.J., **277**, 69
Doh, M.-K., **267**, 127
Dolgushin, F.M., **280**, 233, 243
D'Olieslager, J., **271**, 29
Domasevitch, K.V., **268**, 93
Doménech, A., **268**, 145
Domingo, L.R., **268**, 145
Domingos, Â., **271**, 65
Dominguez-Vera, J.M., **281**, 95
Donaire, A., **275–276**, 73
Dondana, A., **282**, 163
Doneva, S.N., **273**, 403
Dong, J., **273**, 409
Dong, S., **270**, 392
Donnadieu, B., **279**, 76
Donovan, R.J., **270**, 279
Dooley, D.M., **275–276**, 175
Døssing, A., **268**, 159
Douglas, B.E., **278**, 66
Downard, A.J., **267**, 281
Dräger, M., **275–276**, 203
Drago, V., **281**, 111
Drake, J.E., **271**, 75
Draper, J.D., **270**, 405; **274**, 115
Dreos, R., **272**, 74
Drew, M.G.B., **275–276**, 263; **279**, 65, 221
Driess, A., **269**, 143
Driessen, W., **271**, 19
Driessen, W.L., **267**, 183, 193; **282**, 257
Drogunova, G.I., **280**, 238
Drutkowski, U., **278**, 118
Drysdale, K.D., **279**, 121
Du, J.-L., **273**, 354
Duan, C.-Y., **271**, 99
Dubler, E., **279**, 136
Dubourdeaux, P., **282**, 243
Duckworth, P.A., **273**, 372
Dunbar, G.S., **282**, 222
du Preez, J.G.H., **269**, 235; **278**, 96
Dupuis, A., **268**, 125
Durán, M.L., **274**, 131
Duran, N., **278**, 10
Dyba, M., **283**, 1

Eaton, G.R., **273**, 354
Eaton, S.S., **273**, 354
Eckerle, A., **270**, 273
Efimenko, I.A., **278**, 209
Eglin, J.L., **268**, 151; **271**, 49
Einaga, H., **283**, 105, 160
Einerwold, J.G., **268**, 311
Eisenberg, R., **270**, 363
Eisenstein, O., **280**, 26
Elding, L.I., **277**, 236; **282**, 55
Elduque, A., **274**, 15
Elias, H., **278**, 127
Eliezer, I., **271**, 167
Ellis, J.E., **269**, 58
Ellis Jr., P.E., **270**, 162
Emelyanova, N.S., **280**, 288
Emge, T.J., **273**, 310
Eng, S.J., **278**, 170
Engberg, P., **268**, 159
Engel, M., **269**, 157
Englert, U., **269**, 43; **270**, 177; **280**, 150
English, A.M., **275–276**, 250
Enomoto, M., **283**, 80
Enright, G.D., **278**, 143
Ephritikhine, M., **279**, 210
Epstein, L.M., **280**, 302
Ercolani, C., **267**, 109
Eriksson, H., **277**, 232
Erker, G., **270**, 446
Erxleben, A., **283**, 223
Escalante-Tovar, S., **273**, 167
Escrivà, E., **279**, 58
Escuer, A., **269**, 313; **278**, 43
Espenson, J.H., **270**, 55
Estevan, F., **278**, 61
Ettorre, A., **282**, 17
Ezernitskaya, M.G., **280**, 71

Fabbrizzi, L., **267**, 177; **275–276**, 117
Fabretti, A.C., **280**, 282
Fabrizi de Biani, F., **273**, 24; **281**, 53
Faggi, C., **272**, 141
Faller, P., **272**, 150
Fallon, G.D., **279**, 192
Fanizzi, F.P., **275–276**, 410, 419, 500
Fantoni, A., **274**, 122
Fantucci, P., **270**, 479; **273**, 379
Fanwick, P.E., **267**, 123; **278**, 245
Faraone, F., **277**, 119
Fariati, **268**, 135
Farrar, D.H., **271**, 112; **274**, 82
Fattorusso, R., **278**, 76
Faulmann, C., **282**, 90
Faus, J., **268**, 263; **282**, 82
Favretto, M., **273**, 339
Fawcett, J., **274**, 177
Fawzi, R., **269**, 13
Fedin, V.P., **269**, 292; **271**, 228
Fedorov, V., **271**, 222
Fedorov, V.E., **269**, 292
Fedushkin, I.L., **280**, 138
Feher, A., **281**, 134
Fehlhammer, W.P., **269**, 23
Félix, V., **275–276**, 263
Fenton, D.E., **267**, 55; **282**, 42; **283**, 72
Feringa, B.L., **275–276**, 462
Fernández, C.O., **273**, 367
Fernandez, M.B.F., **274**, 73
Fernandez Garcia, M.I., **274**, 73
Fernando, Q., **269**, 224; **271**, 207
Ferrari, A.M., **272**, 252
Ferraro, J., **269**, 211
Ferrence, G.M., **270**, 111
Ferretti, V., **272**, 267
Filler, F.L., **282**, 155
Findsen, E.W., **271**, 119
Fischer, B., **279**, 136
Fischer, J., **283**, 98
Fischer, R., **281**, 181
Fisher, J.J., **270**, 5
Fitzpatrick, M.G., **281**, 101
Flemming, J.P., **280**, 87
Fliegner, J., **269**, 350
Florenciano, F., **281**, 165
Floriani, C., **270**, 298
Folgado, J.-V., **268**, 145; **279**, 58
Font-Bardia, M., **278**, 43
Foran, G.J., **268**, 231
Ford, P.C., **270**, 345
Fornasiero, S., **275–276**, 515
Forster, G.D., **274**, 243
Forster, L.S., **277**, 211
Fortin, S., **279**, 159
Fragale, C., **272**, 38
Fraley, M.E., **280**, 316
Franciò, G., **277**, 119
Franco, D.W., **267**, 93
Franke, R., **270**, 95
Franolic, J.D., **279**, 7
Fraser, D., **274**, 64
Fraústo da Silva, J.J.R., **280**, 308
Frediani, P., **272**, 141
Freiberg, E., **279**, 206
Freisinger, E., **279**, 152; **283**, 223
Frey, U., **269**, 322
Friebel, C., **268**, 117
Fröhlich, R., **270**, 446
Frohmader, G., **267**, 19
Fronczek, F.R., **282**, 96
Frøystein, N.Å., **273**, 41
Fu, W., **277**, 61
Fuentes, S., **274**, 108
Fuess, H., **278**, 190
Fuhrmann, P., **270**, 527
Fujihara, T., **278**, 15
Fujii, Y., **283**, 51
Fujinami, S., **283**, 124
Fujita, J., **281**, 77, 221

Fukuda, Y., **269**, 260
Fukumoto, A., **270**, 459
Fukumoto, T., **283**, 151
Funabiki, T., **275–276**, 222
Furihata, M., **283**, 268
Fusi, V., **268**, 63; **273**, 326; **275–276**, 168
Fuyuhiro, A., **274**, 210; **278**, 15

Gabor, B., **270**, 555
Gaggelli, E., **275–276**, 274
Gaggelli, N., **275–276**, 274
Gajadhar-Plummer, A.S., **281**, 201
Gajda, T., **268**, 297; **274**, 167; **275–276**, 130
Galindo, A., **272**, 125
Gallert, S., **274**, 111
Galuzina, T.V., **280**, 275
Galvez, N., **281**, 95
Gambardella, M.T.do P., **267**, 209
Gambarotta, S., **280**, 143
Gao, S., **282**, 149
García-Granda, S., **273**, 334
García-Lozano, J., **279**, 58
García-Tasende, M.S., **267**, 169
García-Tojal, J., **278**, 150
García-Vázquez, J.A., **271**, 1; **274**, 131; **279**, 249; **281**, 70
Garland, M.-T., **278**, 108; **281**, 18
Garner, C.D., **273**, 225; **275–276**, 150
Garralda, M.A., **282**, 230
Garribba, E., **272**, 68
Gaspar, M., **278**, 51
Gasparri Fava, G., **269**, 297; **275–276**, 320
Gass, H.J., **281**, 43
Gaswick, D.C., **278**, 241
Gatteschi, D., **271**, 9
Gatto, S., **269**, 235
Gau, H.-M., **278**, 232
Gavioli, G., **280**, 282
Gavrish, S.P., **282**, 142
Gebhardt, P., **281**, 181
Geicke, J., **269**, 157; **272**, 101
Gekhman, A.E., **275–276**, 453
Gelbrich, T., **268**, 93
Gelinsky, M., **272**, 115
Gelman, F., **280**, 21
George, T.A., **281**, 153
Geraghty, M., **277**, 257
Geraldes, C.F.G.C., **268**, 249; **273**, 288; **277**, 193
Gerasko, O., **271**, 222
Gerber, T.I.A., **269**, 235; **278**, 96
Geremia, S., **272**, 74; **273**, 62
Gerfin, T., **270**, 499
Gevert, O., **275–276**, 435
Ghilardi, C.A., **269**, 274
Giannini, L., **270**, 298
Giannoccaro, P., **272**, 38
Gierling, K., **269**, 13
Gili, P., **277**, 139
Gillogly, K.A., **275–276**, 106
Giorgi, C., **273**, 326
Girgsdies, F., **280**, 138
Gladysz, J.A., **269**, 167
Glerup, J., **274**, 155
Głowiak, T., **275–276**, 28
Gluth, M.W., **273**, 54
Gobetto, R., **275–276**, 521
Gockel, P., **272**, 115
Goddard, R., **282**, 61
Godinho, S.S.M.C., **279**, 65
Godovikov, I.A., **280**, 238
Goh, C., **270**, 46
Goh, N.-K., **282**, 10
Goldman, A.S., **270**, 537
Goll, W., **283**, 233
Golovaneva, I.F., **280**, 275
Golub, A.S., **280**, 211
Gomes, L., **271**, 83
Gómez Sal, P., **273**, 270; **280**, 1
Gómez-Tagle, P., **273**, 167
Gonçalves, I.S., **275–276**, 263
Gonzalez, F.A., **271**, 137
González, G., **278**, 61
Gonzàlez-Duarte, P., **278**, 10
Gooden, V.M., **268**, 31
Goodfellow, B.J., **273**, 279
Goodgame, D.M.L., **272**, 131
Goodwin, H.A., **274**, 32
Gordon, N.R., **268**, 31
Görls, H., **269**, 283; **281**, 181
Gormal, C.A., **275–276**, 150
Görner, H., **271**, 160
Gorter, S., **271**, 19
Goßner, U., **269**, 47
Gotoh, T., **270**, 424
Gottschalk-Gaudig, T., **269**, 63
Goubitz, K., **268**, 37
Gourkova, A.Y., **280**, 348
Gouzerh, P., **268**, 199
Graiff, C., **269**, 191; **273**, 320
Gramlich, V., **270**, 499; **274**, 219
Grampp, G., **272**, 291
Grannas, M.J., **279**, 192
Grant, G.J., **274**, 192
Grasso, G., **275–276**, 395
Grätzel, M., **274**, 64
Gravereau, P., **281**, 134
Gray, H.B., **270**, 433; **275–276**, 470
Green, M.L.H., **272**, 261
Gregáň, F., **273**, 101
Grepioni, F., **273**, 116
Grevels, F.-W., **271**, 160; **278**, 83
Großheimann, G., **270**, 60
Großmann, G., **269**, 47
Grove, D.M., **269**, 322
Groy, T.L., **267**, 101
Grünsteudel, H., **275–276**, 334
Grushin, V.V., **280**, 87
Gubin, S.P., **280**, 275
Guedes da Silva, M.F.C., **277**, 83
Guerri, A., **281**, 90
Guesmi, S., **280**, 118
Guilhem, J., **275–276**, 234
Guillon, E., **268**, 13
Gullotti, M., **273**, 339; **282**, 180
Gunnewegh, E.A., **268**, 43
Guo, Z., **273**, 1
Gusev, O.V., **280**, 249
Gutiérrez, E., **273**, 244
Gutiérrez-Rodríguez, A., **273**, 334
Gütlich, P., **273**, 54
Gyurcsik, B., **275–276**, 130

Haarman, H.F., **270**, 34
Haas, M., **275–276**, 334
Haase, W., **278**, 127, 190; **281**, 111
Haasnoot, J.G., **268**, 37; **273**, 259; **279**, 217; **282**, 257
Habtemariam, A., **273**, 1, 8
Hacht, B., **283**, 211
Hadden, C.E., **274**, 192
Hadjikakou, S.K., **271**, 243
Hadjikostas, C.C., **271**, 195
Hadjiliadis, N., **272**, 7; **283**, 24
Haefner, S.C., **271**, 187
Hage, R., **268**, 43
Hagen, J., **281**, 53
Haggerty, B.S., **282**, 136
Hahn, F., **281**, 195
Hahn, F.E., **269**, 347
Haid, R., **274**, 143
Haider, J.J., **279**, 44
Haiduc, I., **271**, 75
Haim, A., **270**, 189
Håkansson, M., **277**, 232
Hammud, H.H., **281**, 153
Hamon, J.-R., **268**, 199
Hamon, P., **268**, 199
Hamza, M.S.A., **279**, 178
Handa, M., **278**, 223, 253; **283**, 275
Hang, Q.W., **277**, 21
Hanton, L.R., **281**, 101
Harada (Nee Noguchi), R., **283**, 44
Harcourt, R.D., **269**, 1
Harman, W.D., **270**, 337
Haroutiunian, S.G., **275–276**, 510
Harrod, J.F., **270**, 330
Hartl, F., **268**, 1
Hartmann, J., **280**, 348
Hartmann, M., **267**, 137; **269**, 91
Hartung, W., **274**, 126
Harvey, I., **275–276**, 150
Harvey, J.N., **273**, 111
Hasan, M., **280**, 21
Hashimoto, M., **281**, 190
Hasnain, S.S., **275–276**, 150
Hathcock, D.J., **282**, 131
Hatzisymeon, C., **281**, 121
Haushalter, R.C., **268**, 257; **275–276**, 122; **277**, 263; **282**, 123
Hayashi, T., **275–276**, 159
Haynes, A., **270**, 382
Hazell, R., **268**, 159
Hefter, G., **275–276**, 37
Hegetschweiler, K., **271**, 222; **279**, 136
Hegmans, A., **279**, 152
Heidari, N., **279**, 186
Heinemann, F.W., **269**, 63; **272**, 211; **280**, 39
Heinemann, O., **281**, 53
Heise, K.H., **273**, 234
Helberg, L.E., **270**, 337
Helliwell, M., **273**, 225; **279**, 249
Henderson, W., **277**, 26; **281**, 101
Henling, L.M., **270**, 467
Hennig, H., **271**, 160
Hennige, A., **280**, 39
Henry, B., **275–276**, 130
Heo, N.H., **267**, 127
Herberhold, M., **267**, 19; **273**, 24
Herdtweck, E., **279**, 44
Hernandez, G., **275–276**, 106
Hernández-Molina, M., **277**, 139
Herr, T., **271**, 40
Herrmann, W.A., **279**, 44
Hervás, M., **272**, 109; **275–276**, 73
Hibino, T., **283**, 124
Hidai, M., **267**, 73; **273**, 238; **280**, 163
Higashi, M., **282**, 247
Higgs, T.C., **273**, 14
Hilgers, H.J., **275–276**, 1, 9
Hill, A.M., **271**, 203
Hill, G.S., **278**, 226
Hill, H.A.O., **272**, 261
Hill, M.N.S., **271**, 57
Hill, S.P.W., **272**, 131
Hillhouse, G.L., **270**, 399
Himeno, S., **281**, 214
Hindmarsh, K., **278**, 32
Hiraki, K., **277**, 8
Hiromi, K., **273**, 204
Hirose, J., **273**, 204
Hirota, H., **283**, 243
Hirota, S., **275–276**, 90
Hirsch-Kuchma, M., **267**, 165, 319; **271**, 191; **279**, 206
Hisada, H., **283**, 51
Hisaeda, Y., **273**, 299
Hitchcock, P.B., **269**, 181; **281**, 165
Hitomi, Y., **275–276**, 159
Ho, Y.-C., **278**, 232
Hodgson, D.J., **274**, 155
Hoff, C.D., **270**, 488
Hoffman, M.Z., **279**, 200
Hofmann, A., **273**, 24
Hofmann, J., **269**, 111
Hofmann, T., **267**, 19
Hogarth, G., **274**, 243
Hoggard, P.E., **282**, 217
Holah, D.G., **278**, 226
Holm, R.H., **270**, 46
Holtcamp, M.W., **270**, 467; **274**, 115
Hommerich, B., **274**, 251
Hong, M., **267**, 313; **271**, 93; **274**, 229; **277**, 171
Hoogenraad, M., **279**, 217
Hopfgartner, G., **271**, 36
Hori, A., **281**, 207
Hori, H., **283**, 124

Hori, T., **281**, 214
Horike, S., **278**, 253; **283**, 275
Hormes, J., **270**, 95
Hosokawa, Y., **283**, 118
Hosomizu, M., **273**, 238
House, D.A., **269**, 229; **274**, 42; **277**, 177; **278**, 24, 32
Howell, F.S., **269**, 342; **271**, 13
Huang, C.F., **279**, 233
Huang, J., **272**, 295; **277**, 130; **282**, 261
Huang, T.-T., **270**, 247
Hübener, E., **279**, 152
Hüber, M., **278**, 127
Hueso-Ureña, F., **268**, 77; **277**, 103
Huffman, J.C., **270**, 261
Hughes, A.N., **278**, 226
Hugi-Cleary, D., **275–276**, 1
Hunter, A.D., **282**, 155
Huntington, K.M., **271**, 49
Hüttermann, J., **273**, 131
Huttner, G., **269**, 143
Hwang, J.-W., **283**, 145
Hwang, T.-Y., **278**, 232
Hyslop, A.G., **275–276**, 427

Iaconi, V., **275–276**, 295
Iakovleva, O., **275–276**, 65
Ianelli, S., **268**, 221; **278**, 66
Ibarlucea, L., **282**, 230
Ichimura, K., **277**, 151
Ide, K., **274**, 224
Ienco, A., **275–276**, 366
Ievlev, M.A., **280**, 249
Ihlo, L., **281**, 160
Ike, U.U., **279**, 95
Ikuzawa, M., **281**, 174
Im, S.-C., **283**, 184
Imhof, L., **269**, 347
Imhof, W., **282**, 111
Imori, Y., **281**, 207
Ingallina, P., **270**, 279
Ino, I., **271**, 145
Inomata, Y., **271**, 13
Inoue, K., **273**, 204
Inoue, M., **269**, 224; **271**, 207
Inoue, M.B., **269**, 224; **271**, 207
Inoue, T., **283**, 184
Ioannidis, E., **272**, 283
Ionidis, A.V., **280**, 302
Ishii, Y., **267**, 73; **280**, 163
Ishimori, K., **272**, 80
Itoh, H., **269**, 260
Itoh, K., **277**, 8
Itoh, T., **283**, 51
Iwabuchi, S., **267**, 143
Iwado, A., **283**, 44
Iwamoto, H., **273**, 204
Iwasaki, H., **275–276**, 289
Iwasawa, K., **274**, 224
Izatt, R.M., **278**, 6
Izotova, Y.A., **277**, 83

Jackman, M., **279**, 24
Jackson, W.G., **268**, 85; **274**, 42
Jacob, J., **270**, 55
Jagner, S., **277**, 232
Jahng, Y., **267**, 265
Jakusch, T., **275–276**, 130
James, B.R., **267**, 209; **270**, 130; **273**, 1; **280**, 75
Janczak, J., **281**, 195
Jandrasics, E., **272**, 153
Jang, M., **270**, 363
Janik, M., **269**, 135
Jansen-Varnum, S.A., **282**, 155
Januszczyk, M., **269**, 326
Jaquinod, L., **283**, 98
Jaryszak, E.M., **282**, 217
Jeffery, J.C., **267**, 1, 239, 323; **278**, 178
Jeliazkova, B.G., **273**, 403
Jessop, P.G., **280**, 75
Jezierska, J., **275–276**, 28
Jeżowska-Bojczuk, M., **275–276**, 546
Jhon, M.S., **272**, 33
Ji, D., **273**, 14
Ji, L., **272**, 295
Ji, L.-N., **283**, 17
Ji, W., **279**, 172
Jiang, F., **274**, 229
Jiang, J., **268**, 49, 141
Jiang, Z.-H., **279**, 69, 76
Jiao, X., **272**, 295
Jiménez, H.R., **275–276**, 73
Jiménez, I., **273**, 270
Jiménez, M.V., **272**, 55
Jiménez-Aparicio, R., **277**, 76; **279**, 30
Jiménez-Pulido, S.B., **268**, 77; **277**, 103
Jitsukawa, K., **283**, 105, 160
Johnson, A.R., **275–276**, 215
Johnson, C.R., **267**, 11
Johnson III, W.L., **278**, 91
Jolly, P.W., **270**, 60
Jonas, K., **270**, 555
Jones, A.G., **267**, 165, 319; **271**, 191; **279**, 206
Jones, P.M., **275–276**, 327
Jones, W.D., **270**, 77
Joosten, M., **271**, 29
Journaux, Y., **278**, 159
Ju, T.D., **270**, 488
Julve, M., **268**, 263; **278**, 159; **279**, 127; **282**, 82, 252
Jun, J.-H., **268**, 271
Jung, J.-S., **268**, 271
Jung, M.-J., **267**, 127
Junker, M., **275–276**, 481
Jurkschat1, K., **275–276**, 203
Jutand, A., **273**, 76

Kısakürek, D., **271**, 19
Kaagman, J.-W.F., **270**, 34
Kabata, N., **278**, 223
Kaczmarek, K., **269**, 279
Kahwa, I.A., **281**, 201
Kai, Y., **283**, 184
Kaizaki, S., **274**, 210; **278**, 15; **281**, 77
Kajikawa, Y., **283**, 61
Kalinin, V.N., **280**, 71
Kamachi, T., **267**, 257
Kamaras, P., **279**, 111
Kaneda, Y., **270**, 459
Kang, S.G., **268**, 271; **274**, 24; **279**, 238
Kanitz, R., **281**, 10
Kankare, J., **267**, 63
Kanno, H., **281**, 221
Kano, T., **282**, 209
Kapinos, L.E., **280**, 50
Kappenstein, C., **281**, 134
Karagiannidis, P., **271**, 243; **281**, 121
Karasawa, M., **283**, 184
Kashiwabara, K., **281**, 77
Kasprzykowski, F., **283**, 1
Kasuga, K., **278**, 223, 253; **283**, 275
Katada, M., **267**, 143; **269**, 310; **283**, 80
Kataoka, K., **275–276**, 289
Kato, M., **279**, 51
Kato, T., **278**, 223
Kato, Y., **269**, 260
Katrusiak, A., **269**, 326
Kaufmann, B., **269**, 101
Kawaguchi, H., **283**, 136
Kawamoto, T., **282**, 71; **283**, 260
Kawase, Y., **267**, 201
Kawata, S., **267**, 143; **283**, 80
Kazankov, G.M., **280**, 57
Keefer, C.E., **282**, 200
Keim, W., **280**, 150
Keita, B., **275–276**, 234
Keitel, D.P., **280**, 150
Kempinger, W., **269**, 111
Kennedy, B.J., **268**, 231
Kennedy, M., **273**, 279
Keppler, B.K., **267**, 137
Kerpen, K., **271**, 160
Kessissoglou, D.P., **272**, 24, 283
Khajehnouri, F., **271**, 231
Khan, M.A., **270**, 227
Khan, M.I., **277**, 69
Kidani, Y., **283**, 251
Kikkawa, M., **273**, 204
Kikuchi, A., **283**, 151
Killat, S., **280**, 150
Kim, D., **279**, 243
Kim, D.-Y., **267**, 127
Kim, E., **278**, 103
Kim, H.H., **268**, 271
Kim, J., **274**, 24
Kim, J.E., **281**, 229
Kim, J.-H., **283**, 145
Kim, K., **279**, 238
Kim, K.M., **279**, 116
Kim, M.-S., **279**, 238
Kim, S.-J., **274**, 24
Kimble, S.B., **267**, 101
Kimoto, H., **281**, 174; **283**, 118
Kimura, E., **270**, 424
King, R.B., **270**, 68; **277**, 202
Kingma, I.E., **269**, 203
Kirchner, K., **268**, 69
Kirmse, R., **271**, 199; **281**, 160
Kirschbaum, K., **274**, 251
Kiss, E., **283**, 202
Kiss, T., **268**, 297; **283**, 202
Kita, M., **281**, 77
Kita, S., **274**, 247
Kitagawa, S., **267**, 143; **271**, 129; **283**, 80
Kitagawa, T., **283**, 184
Kivekas, R., **281**, 95
Kjelsberg, M.A., **279**, 7
Klapötke, T.M., **269**, 1
Klausmeyer, K.K., **270**, 405
Klement, R., **278**, 127
Klesczewski, B.G., **270**, 511
Klinga, M., **281**, 95
Klooster, W.T., **279**, 7
Klumpp, G.W., **269**, 203
Klyagina, A.P., **280**, 275
Kneebone, P.E., **281**, 101
Knight, B.W., **282**, 200
Knoch, F., **272**, 211
Knowles, P.F., **275–276**, 175
Kobayashi, M., **283**, 256
Kobayashi, N., **277**, 31
Kobuke, Y., **283**, 167
Koch, S.A., **279**, 7
Kochubey, D.I., **280**, 339
Koetzle, T.F., **279**, 7
Kögerler, P., **268**, 239
Köhl, G., **270**, 95
Kohzuma, T., **283**, 184
Koikawa, M., **274**, 224; **277**, 134
Koike, T., **270**, 424
Kojima, H., **275–276**, 222
Kojima, M., **271**, 24; **274**, 184; **282**, 209
Kokkou, S., **281**, 121
Kolb, U., **275–276**, 203
Kolnaar, J.J.A., **282**, 257
Kolomiychuk, V.N., **280**, 339
Kolotilov, S.V., **278**, 217
Kolp, B., **272**, 153
Komatsu, E., **281**, 85
Koo, K., **270**, 399
Kooijman, H., **279**, 217
Koskimies, J.K., **268**, 21
Kostikas, A., **283**, 111
Kosugi, E., **275–276**, 90
Kováč, J., **281**, 134
Kovacik, I., **275–276**, 435
Kövari, E., **274**, 251
Kozłowski, H., **275–276**, 546; **283**, 1
Kraatz, H.-B., **278**, 143
Krämer, R., **269**, 117
Kramolowsky, R., **269**, 317; **281**, 64
Kratz, F., **267**, 133
Kravtsov, D.N., **280**, 238
Krebs, B., **274**, 251
Kreher, U., **281**, 181
Kressl, C., **275–276**, 1
Krickemeyer, E., **271**, 9
Kröckel, M., **271**, 9; **283**, 111
Kroneck, P.M.H., **275–276**, 52
Krüger, C., **270**, 273; **281**, 53
Krut'ko, D.P., **280**, 257

Kryukova, G.A., **280**, 339
Krzyminiewski, R., **269**, 326
Ksenofontov, V., **273**, 54
Kubiak, C.P., **270**, 111
Kubiak, R., **281**, 195
Kubota, K., **269**, 310
Kuchiyama, Y., **277**, 31
Kudo, H., **269**, 260
Kudo, K., **277**, 151
Kudryavtsev, A.B., **267**, 293
Kühn, F.E., **279**, 44
Kukushkin, V.Yu., **277**, 83
Kuma, H., **283**, 260
Kumagai, H., **267**, 143; **283**, 80
Kumar, K., **275–276**, 106
Kumar, S.B., **269**, 313
Kumita, H., **283**, 160
Kuribayashi, Y., **281**, 207
Kurihara, H., **282**, 247
Kurk, D.N., **281**, 153
Kuroda, A., **279**, 165
Kuroda-Sowa, T., **268**, 317; **271**, 129, 145; **281**, 116
Kurosawa, H., **270**, 87
Kurzeev, S.A., **280**, 57
Kushi, Y., **282**, 71
Kuzmann, E., **274**, 167
Kuz'mina, L.G., **280**, 257
Kuznetsov, V.F., **280**, 172

Labarre, J.-F., **268**, 307
Labinger, J.A., **270**, 433, 467
Lachicotte, R.J., **270**, 363
Ladogana, S., **267**, 49; **271**, 105; **278**, 202
Lahoz, F.J., **274**, 15
Lahuerta, P., **278**, 61
Lai, Y.-H., **279**, 221
Laikov, D.N., **280**, 328
Lakin, M.T., **267**, 183
Lamb, J.D., **282**, 142
Lambert, F., **275–276**, 234
Lammers, H., **268**, 249; **277**, 193
Lampeka, Y.D., **282**, 142
Lamrani, M., **283**, 145
Lance, M., **279**, 210
Landín, I.O., **269**, 224
Landis, C.R., **270**, 285
Landis, K.G., **282**, 155
Lando, D.Y., **275–276**, 510
Lanfranchi, M., **273**, 320; **275–276**, 500
Lang, J.-P., **283**, 136
Lang, M., **269**, 111
Lang, R.F., **270**, 488
Langes, C., **274**, 143
Łankiewicz, L., **283**, 1
Lappert, M.F., **269**, 181
Larsen, R.W., **271**, 119
Lasagni, B., **270**, 123
Latajka, R., **283**, 1
Latos-Grażyński, L., **273**, 184; **277**, 242
Latour, J.-M., **282**, 243
Latva, M., **267**, 63
Laurent, J.-P., **268**, 125
Law, W.-F., **268**, 49, 141
Layh, M., **269**, 181
Lazardi, K., **275–276**, 528
Le, Q.T.H., **267**, 201
Leadbeater, N.E., **278**, 250
Lebedev, A.M., **273**, 234
Lebreton, C., **272**, 188
Leduc, P., **272**, 4
Lee, B.-W., **267**, 127
Lee, C.-T., **274**, 7
Lee, D.-J., **267**, 127
Lee, E.Y., **269**, 337
Lee, G.H., **279**, 233
Lee, I.S., **279**, 243
Lee, J., **270**, 353
Lee, K.S., **279**, 7
Lee, S.S., **279**, 243; **281**, 229
Lee, Y.-A., **279**, 116
LeGall, J., **273**, 409
Legros, J.-P., **271**, 217
Lehmann, A., **268**, 109; **277**, 219
Leigh, G.J., **280**, 193
Leising, R.A., **273**, 393
Le Maux, P., **273**, 142
Lemenovskii, D.A., **280**, 257
Leong, A.J., **273**, 372
Le Pichon, L., **272**, 188; **280**, 118
Leplawy, M.T., **269**, 279
Leung, Y.C., **272**, 261
Leupold, O., **275–276**, 334
Levason, W., **271**, 203
Levchenko, G., **273**, 54
Lever, A.B.P., **281**, 126
Levitin, I., **270**, 169; **280**, 295
Levitsky, V., **282**, 61
Levy, E.G., **281**, 101
Lewis, J., **274**, 201
Lezama, L., **268**, 239; **278**, 150; **279**, 58
L'Haridon, P., **282**, 173
L'Her, M., **282**, 173
Li, C.-H., **269**, 302
Li, H., **268**, 145; **271**, 99, 174
Li, J., **273**, 255, 310
Li, L., **279**, 121
Li, Q., **270**, 551
Li, S., **277**, 69
Li, S.-L., **268**, 177
Li, W., **282**, 96
Li, Y., **279**, 221
Li, Z., **270**, 279
Liable-Sands, L., **270**, 543
Liang, L.-C., **270**, 353
Lianza, F., **274**, 219
Liao, D.-Z., **279**, 69, 76
Lima, E.C., **267**, 93
Lima, M.J., **273**, 279
Lin, C.-W., **270**, 247
Lin, F.-T., **268**, 287; **271**, 124
Lin, J., **268**, 141
Linati, L., **273**, 339
Lindner, E., **269**, 13
Lindoy, L.F., **273**, 372
Linert, W., **267**, 293; **278**, 190
Liou, L.-S., **269**, 302; **274**, 7
Lippard, S.J., **270**, 527
Lippert, B., **267**, 87; **269**, 135; **273**, 31; **279**, 152; **282**, 237; **283**, 223
Lippolis, V., **267**, 115
Lipponer, K.-G., **267**, 137
Lipunova, G.N., **280**, 30
Lis, T., **277**, 127
List, A.K., **270**, 399
Liu, G., **275–276**, 58
Liu, H., **267**, 313; **271**, 93; **272**, 295; **274**, 229; **277**, 171
Liu, H.-L., **282**, 90
Liu, M.-Y., **273**, 409
Liu, S.-X., **282**, 149
Liu, W., **268**, 49, 141
Lloret, F., **268**, 263; **278**, 159; **279**, 127; **282**, 82, 252
Lobana, T.S., **267**, 169
Lobbia, G.G., **282**, 1
Lock, C.J.L., **274**, 137
Lodeiro, C., **267**, 55
Lolli, S., **270**, 479
Lombardi, A., **275–276**, 301
Lombardi, M.G., **282**, 82, 252
Long, G.S., **279**, 1
Longato, B., **277**, 163
Lo Passo, C., **275–276**, 141, 242
Lopes, A.D., **279**, 44
López, D.M., **269**, 224
López, G., **281**, 165
López, J.A., **274**, 15
Lorenz, I.-P., **269**, 157; **272**, 101
Lorenzo Luis, P.A., **277**, 139
Lough, A.J., **270**, 238
Louro, M.S., **271**, 1
Louro, R.O., **273**, 196
Lozano, C.M., **271**, 137
Lu, J., **267**, 313; **274**, 229; **275–276**, 58, 447
Lu, Q., **270**, 207
Lu, Y., **268**, 257
Lucchini, V., **275–276**, 385; **277**, 247
Luchinat, C., **269**, 125; **275–276**, 373; **283**, 12
Lucken, E.A.C., **271**, 231
Luckert, S., **269**, 43
Lügger, T., **269**, 347
Luo, G.-T., **272**, 295
Lyons, J.E., **270**, 162
Lyssenko, K.A., **280**, 249, 288

Ma, Z., **268**, 103
Maßau, M., **282**, 61
Macías, A., **267**, 55; **282**, 42
Macey, D.J., **267**, 7
MacFarlane, K.S., **267**, 209; **270**, 130
Macinai, R., **269**, 125
Mackie, A.G., **282**, 25
Maćkiewicz, Z., **283**, 1
Maclean, A.L., **268**, 231
Madden, J., **282**, 131
Madden, L.K., **278**, 241
Maeda, Y., **283**, 124
Maejima, N., **283**, 105
Maekawa, M., **268**, 317; **271**, 129, 145; **281**, 116
Maestri, M., **277**, 225
Magafa, V., **272**, 7
Maglio, O., **275–276**, 301
Maguire, J.A., **270**, 77
Maichle-Mössmer, C., **270**, 177
Maisonat, A., **275–276**, 182
Maitlis, P.M., **270**, 382; **280**, 62, 249
Mak, T.C.W., **268**, 177; **270**, 551; **271**, 99; **277**, 21
Malamatari, D.A., **272**, 283
Maldotti, A., **272**, 197
Malik, M.A., **274**, 239
Malinak, S.M., **278**, 241
Malinski, T., **271**, 167
Maloshtan, I.M., **282**, 142
Mamasachlissov, Eu.Sh., **275–276**, 510
Mandon, D., **275–276**, 334
Mangani, S., **275–276**, 295, 349
Mani, F., **273**, 151
Mann, K.L.V., **278**, 178
Mano, H., **274**, 184
Manotti Lanfredi, A.M., **267**, 177; **273**, 259
Manriquez, V., **267**, 151
Mansuy, D., **272**, 4
Manzanero, A., **280**, 1
Manzur, C., **268**, 199
Marchetti, F., **270**, 177
Marchi, A., **272**, 267
Marek, J., **267**, 299
Marek, K., **270**, 151
Maresca, K., **281**, 70
Mari, M.N., **270**, 123
Marić, Lj., **271**, 239
Marques, M.P.M., **273**, 288
Marras, P., **275–276**, 440
Marsich, N., **278**, 237
Martell, A.E., **270**, 207; **278**, 170; **281**, 60
Martín, A., **273**, 270
Martin, D.L., **278**, 91
Martin, R.B., **283**, 30
Martín, S., **280**, 189
Martínez, J., **280**, 62
Martinez, M., **278**, 61
Martini, G., **272**, 274
Martinuzzi, S., **269**, 125
Martín-Zarza, P., **277**, 139
Maruyama, T., **269**, 310
Marvelli, L., **272**, 267; **280**, 302
Marzilli, L.G., **275–276**, 90
Marzotto, A., **275–276**, 410, 419
Masarwa, M., **270**, 151
Mascharak, P.K., **269**, 269
Mashima, K., **270**, 459
Masi, D., **275–276**, 528
Mason, W.R., **278**, 113
Maspoch, S., **278**, 10
Massa, W., **268**, 117
Masuda, H., **275–276**, 222; **283**, 105, 160
Matias, P.M., **273**, 213
Matijašić, I., **282**, 76

Matikainen, J., **268**, 21
Matonic, J.H., **267**, 173
Matović, Z.D., **268**, 221; **278**, 66, 209
Matsubayashi, G., **279**, 165
Matsuhiro, B., **273**, 191
Matsui, M., **267**, 201
Matsumoto, K., **272**, 162; **281**, 174; **283**, 118, 256
Matsushima, H., **274**, 224
Matsushita, R., **283**, 175
Matsuzawa, Y., **277**, 151
Matthews, H.R., **277**, 89
Mattioli, M., **280**, 282
Matzanke, B.F., **275–276**, 334
Maury, O., **279**, 210
Mautner, F., **269**, 313
May, J.A., **275–276**, 327
May, L., **279**, 111
May, P.M., **275–276**, 37
Maye, J.P., **280**, 8
Mayer, J.M., **270**, 202
Mayr, A., **279**, 7
Mays, M.J., **277**, 186
Mazzei, T., **281**, 90
McAuliffe, C.A., **279**, 249; **282**, 25
McCaffrey, L.J., **281**, 101
McCann, M., **277**, 257; **279**, 24
McCann, S., **279**, 24
McCarley, R.E., **271**, 239
McCleverty, J.A., **267**, 323
McCollum, D.G., **270**, 13
McCrea, J., **277**, 257
McFarlane, W., **281**, 121
McGarvey, B.R., **272**, 43
McKee, V., **277**, 257; **279**, 24
McKenna, C.E., **280**, 193
McLeod, J.J., **278**, 241
Mc Mahon, C.N., **280**, 193
McMillin, D.R., **273**, 346
McMorran, D.A., **281**, 101
McPartlin, M., **273**, 372
McPherson, M.J., **275–276**, 175
Mdleleni, M.M., **270**, 345
Mealli, C., **275–276**, 366
Meetsma, A., **271**, 180
Mehlman, S.A., **267**, 159
Meienberger, M., **279**, 136
Meier, B., **275–276**, 65
Meli, A., **272**, 55
Melník, M., **277**, 127
Memon, N., **268**, 211
Menabue, L., **268**, 205; **273**, 397
Menon, M., **275–276**, 551
Merbach, A.E., **275–276**, 15
Mercuri, M.L., **267**, 115; **273**, 175
Mereiter, K., **268**, 69
Meshulam, A., **273**, 266
Messerschmidt, A., **273**, 160
Messori, L., **275–276**, 510; **281**, 90
Mestroni, G., **273**, 62
Meunier, B., **272**, 228
Meyer, G., **273**, 76
Meyer, J., **271**, 9
Meyer, L.A., **275–276**, 122
Meyer, T.J., **270**, 511
Meyer-Klaucke, W., **275–276**, 334
Meyerstein, D., **270**, 440; **273**, 266
Micera, G., **268**, 297; **269**, 279; **272**, 68; **275–276**, 440; **283**, 202, 233
276,Michelin, R.A., **275–276**, 515; **280**, 308
Micheloni, M., **268**, 63; **275–276**, 168
Michelsen, K., **274**, 155
Midollini, S., **269**, 274
Mifune, M., **283**, 44
Mikata, Y., **279**, 51
Miki, E., **267**, 147
Milius, W., **267**, 19; **273**, 24
Millar, S.P., **270**, 363
Miller, J.R., **278**, 229
Minakami, M., **273**, 204
Mini, E., **281**, 90
Minutolo, F., **275–276**, 462
Mironova, S.A., **270**, 221
Mirza, H.A., **271**, 112
Miyaki, Y., **270**, 87
Miyamoto, T.K., **279**, 105
Miyashita, I., **283**, 256
Mizobe, Y., **267**, 73; **273**, 238
Mizutani, M., **283**, 105
Moerman, K.L., **281**, 10
Mogami, K., **279**, 51
Mohamadou, A., **268**, 13
Mohrmann, L., **283**, 251
Moinet, C., **273**, 142
Moïse, C., **268**, 323
Moiseev, I.I., **275–276**, 453; **280**, 339
Moiseev, S.K., **280**, 71
Moiseeva, N.I., **275–276**, 453
Moisidis, G., **272**, 283
Mok, K.F., **279**, 221
Moldrheim, E., **273**, 41
Molina, J., **274**, 53
Molinari, A., **272**, 197
Moliner, N., **274**, 1
Moll, M., **272**, 211
Monacelli, F., **267**, 109
Mondal, J.U., **278**, 1
Moneta, W., **282**, 243
Moneti, S., **272**, 55
Monge, A., **273**, 244
Monkiewicz, J., **270**, 273
Monleón, D., **275–276**, 73
Monsù Scolaro, L., **275–276**, 141, 242
Monzani, E., **273**, 339; **282**, 180
Moore, G.R., **275–276**, 256
Morais, J., **273**, 213
Morales, J.D., **271**, 137
Morandini, F., **282**, 163
Moratal, J.M., **275–276**, 73
Morelli, G., **278**, 76
Moreno, C., **274**, 82
Moreno, J.M., **281**, 95
Moreno-Carretero, M.N., **268**, 77; **277**, 103
Moret, M., **282**, 163
Morgenstern-Badarau, I., **275–276**, 234
Mori, T., **267**, 329
Mori, W., **275–276**, 43; **283**, 118, 268
Morice, C., **273**, 142
Moriguchi, T., **281**, 190
Morikawa, T., **273**, 204
Morikita, T., **269**, 310
Morishima, I., **272**, 80
Morita, K., **283**, 167
Moriwaki, K., **268**, 317
Moro, S., **272**, 62
Morozov, V.F., **275–276**, 510
Morris, R.H., **270**, 238
Motekaitis, R.J., **278**, 170; **281**, 60
Motohashi, N., **283**, 44
Moura, I., **273**, 279, 409
Moura, J.J.G., **273**, 279, 409
Mozzon, M., **275–276**, 515
Mroziński, J., **277**, 127
Muir, M.M., **271**, 137
Mukaida, M., **269**, 342
Mukkala, V.-M., **267**, 63
Mülhaupt, R., **267**, 133
Müller, A., **268**, 239; **271**, 9; **280**, 30
Muller, I., **278**, 197
Muller, J.G., **275–276**, 314
Müller, L., **278**, 127
Muller, R.N., **273**, 92; **275–276**, 15
Munakata, M., **268**, 317; **271**, 129, 145; **281**, 116
Munari, I., **282**, 163
Muñoz, M.C., **268**, 263; **274**, 1
Mura, P., **282**, 17
Murakami, T., **274**, 247
Murakami, Y., **273**, 299
Murayama, K., **281**, 36
Murillo, C.A., **267**, 173
Mürner, H., **271**, 36
Murphy, C.A., **279**, 121
Mynott, R., **270**, 555

Nadjo, L., **275–276**, 234
Nagai, T., **267**, 147
Nagai, Y., **275–276**, 222
Nagao, H., **269**, 342
Nagao, N., **269**, 342
Nagasawa, A., **269**, 260; **283**, 256
Nagata-Tanba, S., **267**, 147
Nagatomo, S., **283**, 184
Nagel, U., **269**, 34
Nagy, L., **274**, 167
Nakajima, K., **274**, 184; **282**, 209
Nakajima, S., **283**, 243
Nakamura, A., **270**, 459; **275–276**, 43; **283**, 91
Nakamura, H., **283**, 243
Nakamura, N., **283**, 260
Nakano, E., **281**, 207
Nakano, H., **270**, 459
Nakano, M., **279**, 165
Nakao, Y., **283**, 118
Nakashima, M., **277**, 134
Nakayama, J., **283**, 256
Nardin, G., **272**, 74; **278**, 237; **282**, 180
Nash, K.L., **269**, 211
Nastri, F., **275–276**, 301
Natile, G., **275–276**, 500
Navarro, A., **271**, 174
Navarro, J.A., **272**, 109; **275–276**, 73
Navarro, J.A.R., **274**, 53
Navarro, M., **275–276**, 528
Navarro, R.E., **269**, 224
Navon, G., **273**, 388
Nayak, S.K., **267**, 49
Nazarenko, A.Y., **282**, 142
Ndifon, P.T., **282**, 25
Nedden, H.G., **269**, 34
Neenan, T.X., **267**, 193
Nefedov, V.I., **273**, 234
Negishi, E., **280**, 8
Németh, S., **270**, 326
Nesper, R., **274**, 64
Nesset, M.J.M., **272**, 204
Nesterenko, P., **280**, 295
Neuburger, M., **275–276**, 359
Neves, A., **281**, 111
Nevodchikov, V.I., **280**, 138
Ng, D.K.P., **268**, 49, 141
Nicasio, M.C., **273**, 244
Nichols, P.J., **279**, 192
Nicholson, T., **267**, 165, 319; **271**, 191; **279**, 206
Niël, J., **268**, 43
Nielsen, J.B., **274**, 232
Nierlich, M., **279**, 210
Niiya, H., **281**, 214
Nishikawa, N., **283**, 91
Nitsche, H., **273**, 234
Nock, B.A., **281**, 148
Noels, A.F., **275–276**, 528
Noguchi, Y., **272**, 162
Nolan, K.B., **268**, 313
Nolting, H.F., **275–276**, 65
Nomura, Y., **267**, 73
Norman, R.E., **273**, 393
Nöth, H., **269**, 7, 101
Novgorodov, B.N., **280**, 339
Novikov, Y.N., **280**, 211
Numan, M., **281**, 121
Nunes, D., **271**, 65
Nyrönen, T., **277**, 55

Oberhauser, W., **274**, 143
O'Brien, P., **274**, 239
Ochertyanova, E.A., **280**, 328
O'Connor, C.J., **268**, 271; **277**, 69
Oda, R., **277**, 8
Odani, A., **278**, 136
Odom, A.L., **278**, 103
Oevering, H., **267**, 183
Ogawa, A., **273**, 299
Ogoshi, H., **275–276**, 159

Oh, C.-E., **267**, 127
Ohba, M., **283**, 72
Ohms, G., **269**, 47
Ohnishi, S., **283**, 136
Ohno, T., **273**, 299
Ojima, I., **270**, 279
Okamura, T., **275–276**, 43; **283**, 91
Ōkawa, H., **283**, 72
Okeya, S., **281**, 174
Okura, I., **267**, 257; **279**, 51
Olejnik, J., **269**, 279
Oliva, G., **267**, 209
Olive, M., **281**, 1
Olk, R.-M., **281**, 160
Olmstead, M.M., **269**, 269; **270**, 28
Ondracek, A.L., **267**, 123; **278**, 245
Ondrejkovičová, I., **277**, 127
Onishi, M., **277**, 8
Onodera, K., **283**, 184
Ooi, B.-L., **274**, 103
Oomura, K., **269**, 342
Ooyama, D., **269**, 342
Orendáč, M., **281**, 134
Orendáčová, A., **281**, 134
Orioli, P., **275–276**, 295, 510; **281**, 90
Orlandini, A., **269**, 274
Orlinkov, A., **280**, 355
Oro, L.A., **274**, 15
Orth, S.D., **270**, 337
Orvig, C., **273**, 47
Osman, A.M., **275–276**, 98
Ostuni, A., **268**, 205
Otter, C.A., **278**, 178
Owens, J.W., **277**, 1; **279**, 226
Ożarowski, A., **275–276**, 28
Ozawa, T., **281**, 207
Ozette, K., **272**, 4

Pacchioni, G., **275–276**, 500
Pacholska, E., **273**, 184
Padmakumar, R., **270**, 392
Paetzold, P., **269**, 43
Pagliarin, R., **282**, 180
Paine, R.T., **269**, 101
Palanca, P., **268**, 145; **271**, 174
Pallavicini, P., **267**, 177; **275–276**, 117
Palma, P., **269**, 191; **275–276**, 168
Palmisano, G., **272**, 18
Pályi, G., **280**, 282
Pampaloni, G., **270**, 177
Panunzi, B., **281**, 141
Panz, C., **269**, 73
Panzalorto, M., **277**, 119
Panzanelli, A., **269**, 279
Paoletti, P., **273**, 326
Paoli, P., **275–276**, 168; **282**, 76, 252
Papadimitriou, C., **267**, 299
Papadopoulos, A.N., **272**, 283
Papadopoulos, M., **281**, 148
Papadopoulos, M.S., **268**, 169
Papaefthymiou, V., **283**, 111
Papánková, B., **278**, 190
Parajón-Costa, B.S., **268**, 239
Parak, F., **275–276**, 65
Parand, A., **268**, 211
Park, J.-B., **273**, 409
Park, J.G., **267**, 265
Parri, L., **275–276**, 349
Paskonova, E., **280**, 295
Pastor, A., **272**, 125
Pastushok, V.N., **278**, 6
Patel, B., **270**, 537
Patrício, L., **271**, 65
Paulo, A., **271**, 65
Paulsen, H., **275–276**, 334
Paulus, H., **278**, 127
Pavlishchuk, V.V., **278**, 217
Pavone, V., **275–276**, 301
Pazderski, L., **273**, 259
Pearson, J.M., **270**, 382
Pearson, R.G., **270**, 252
Pecoraro, V.L., **273**, 85; **283**, 37
Pedone, C., **275–276**, 301; **278**, 76
Pedraz, T., **273**, 270
Pedrosa de Jesus, J.D., **281**, 225
Peeters, O.M., **269**, 332
Pelecanou, M., **268**, 169; **281**, 148
Pelizzi, C., **269**, 297
Pelizzi, G., **278**, 209
Pellacani, G.C., **272**, 252
Pellei, M., **282**, 1
Pellinghelli, M.A., **267**, 115; **273**, 175
Pelosi, G., **268**, 221; **269**, 297; **275–276**, 320; **278**, 66
Penner, N., **280**, 295
Penner-Hahn, J.E., **275–276**, 215
Pepi, F., **275–276**, 192
Perec, M., **281**, 18
Peregudov, A.S., **280**, 238
Pereira, E., **271**, 83
Perera, S.D., **270**, 312; **279**, 95
Peressini, S., **270**, 145; **280**, 322
Pérez, P.J., **273**, 244
Perez-Carreño, E., **275–276**, 366
Pérez-Lourido, P., **271**, 1; **279**, 249
Perils, J., **269**, 235; **278**, 96
Pernice, I., **275–276**, 141, 242
Perry, M., **277**, 1
Pertici, P., **268**, 323
Peruzzini, M., **272**, 1; **275–276**, 366; **280**, 302
Peters, J.A., **268**, 249; **277**, 193
Petroni, D., **270**, 177
Petrova, G.N., **270**, 221
Petrovskii, P.V., **280**, 71, 233, 243, 249
Petrukhina, M.A., **267**, 173
Petter, W., **274**, 219
Pettinari, C., **282**, 1
Pettit, L.D., **283**, 1
Philippakopoulos, P., **272**, 24
Philippopoulos, A.I., **283**, 24
Phillips, D.J., **268**, 135; **271**, 210
Piacenti, F., **272**, 141
Piccioli, M., **283**, 12
Picher, M.T., **268**, 145; **271**, 174
Pichon, R., **282**, 173
Pierattelli, R., **269**, 125
Pietro, W.J., **281**, 126
Pietrzak, J., **269**, 326
Pillinger, M., **279**, 44
Pilloni, G., **275–276**, 401; **277**, 163; **282**, 163
Pilon, M.C., **280**, 87
Pinilla, E., **282**, 230
Pinillos, M.T., **274**, 15
Pinkerton, A.A., **274**, 251
Pintus, G., **273**, 175
Pinzino, C., **270**, 177
Pirmettis, I.C., **268**, 169; **281**, 148
Pisareva, I.V., **280**, 233
Pizarro, J.L., **278**, 150
Pizzotti, M., **270**, 479; **280**, 282
Place, H., **279**, 1
Planinić, P., **271**, 239
Plass, W., **271**, 9; **280**, 30
Platt, A.W.G., **274**, 177
Plotkin, S., **270**, 189
Podberezskaya, N.V., **269**, 292
Poë, A.J., **268**, 189; **274**, 82
Poggi, A., **275–276**, 117
Pogni, R., **275–276**, 274, 557
Poilblanc, R., **275–276**, 182; **283**, 24
Polborn, K., **269**, 73, 157; **272**, 101; **273**, 106
Pollmann, J., **270**, 95
Polyakov, V.A., **267**, 293; **280**, 57
Pombeiro, A.J.L., **271**, 65; **277**, 83; **280**, 308
Pomije, M.K., **269**, 58
Pompe, S., **273**, 234
Pond, A.E., **275–276**, 250
Ponomareva, V.V., **268**, 93
Pontellini, R., **268**, 63; **275–276**, 168
Ponticelli, G., **268**, 221; **278**, 209
Pozdniakova, Yu.A., **280**, 282
Pozzi, G., **272**, 274
Predieri, G., **273**, 320; **275–276**, 320
Prévot, L., **283**, 98
Priego, J.L., **277**, 76; **279**, 30
Primus, J.-L., **275–276**, 98
Pritchard, R.G., **282**, 25
Priuska, E.M., **273**, 85
Prokopchuk, E.M., **278**, 226
Prónayová, N., **273**, 101
Proserpio, D.M., **273**, 255, 310
Protchenko, A.V., **280**, 138
Protzenko, G.A., **280**, 211
Provó Kluit, H.A.J., **267**, 183
Prushan, M.J., **278**, 217
Prytkov, S., **279**, 111
Psillakis, E., **278**, 178
Psomas, G., **272**, 24
Pubanz, D., **275–276**, 15
Pullen, A.E., **282**, 90
Purches, G., **272**, 1
Purrington, S.T., **282**, 200

Quaranta, E., **272**, 38
Que, Jr. , L., **273**, 393
Queiroz, S.L., **267**, 209
Quirós-Olozábal, M., **268**, 77; **277**, 103

Rabe, G.W., **267**, 309
Rabinovich, D., **274**, 232
Rackelmann, G.H., **272**, 211
Radanović, D.D., **268**, 221
Radanović, D.J., **268**, 221; **278**, 66, 209
Rae, A.D., **271**, 210
Raffaelli, A., **275–276**, 462
Rahimian, K., **270**, 330
Raithby, P.R., **274**, 201; **277**, 186
Rakintzis, N.Th., **273**, 201
Ralph, S.F., **281**, 10
Ramachandran, R., **274**, 82
Ramkisoensing, K., **279**, 217
Randaccio, L., **272**, 74; **278**, 237; **282**, 180
Ranford, J.D., **267**, 27
Raper, E.S., **271**, 57, 195; **277**, 14
Rapta, M., **279**, 111
Raptopoulou, C.P., **268**, 169; **272**, 24, 283
Ratajczak-Sitarz, M., **269**, 326
Rath, N.P., **271**, 236
Ratilainen, J., **277**, 55
Re, N., **270**, 298
Read, J.F., **267**, 159; **281**, 43
Real, J.A., **274**, 1
Realo, E., **275–276**, 334
Reedijk, J., **267**, 183, 193; **268**, 37; **271**, 19; **273**, 259; **279**, 217; **282**, 257; **283**, 251
Reetz, M.T., **282**, 61
Refosco, F., **275–276**, 401
Regan, J.J., **275–276**, 470
Rehder, D., **267**, 229
Reibenspies, J.H., **270**, 207, 405; **274**, 115
Reich, T., **273**, 234
Reinerth, W.A., **267**, 335
Ren, L., **268**, 271
Rennie, M.-A., **277**, 186
Rettig, S.J., **267**, 209; **270**, 130
Reutov, O.A., **280**, 257
Rey, L., **273**, 244
Reyes, M.L., **273**, 244
Reynolds, J.R., **282**, 90
Reza, Md Y., **274**, 224
Rheingold, A.L., **267**, 309; **270**, 399, 543, 559; **282**, 136
Ribot, F., **279**, 144
Ricciuto, L., **270**, 238
Richards, J.H., **275–276**, 470
Richards, R.L., **275–276**, 150
Richmond, M.G., **267**, 223
Richter, R., **271**, 40
Rieder, A., **274**, 143
Riera, V., **268**, 131

Rietjens, I.M.C.M., **275–276**, 98
Riley, M.J., **268**, 55
Rillema, D.P., **279**, 200
Rinker, R.G., **270**, 345
Riordan, C.G., **270**, 543
Rissanen, K., **277**, 55
Ristori, S., **272**, 274
Ritter, K., **271**, 160
Rivero, I.A., **274**, 108
Rivière-Baudet, M., **277**, 98
Rizopoulos, A.L., **273**, 72
Rizzarelli, E., **275–276**, 395
Rizzoli, C., **270**, 298
Robert, A., **272**, 228
Robert, F., **268**, 199; **279**, 144
Robertson, N., **279**, 65
Robins, M., **279**, 226
Robinson, R., **279**, 226
Robinson, W.T., **277**, 177
Robl, C., **269**, 47, 83
Rode, B.M., **272**, 89
Rodgers, K.K., **275–276**, 481
Rodríguez, A., **267**, 55; **274**, 131; **281**, 70; **282**, 42
Rodríguez-Blas, T., **267**, 55; **282**, 42
Rodríguez-Cabán, J.L., **271**, 137
Rogers, M.S., **275–276**, 175
Rogers, R.D., **269**, 211
Rojo, T., **268**, 239; **278**, 150
Romão, C.C., **275–276**, 263; **279**, 44
Rombeck, I., **273**, 31
Romeo, A., **275–276**, 242
Romeo, R., **275–276**, 141
Romero, J., **271**, 1; **274**, 131; **279**, 249; **281**, 70
Romero, M.A., **274**, 53
Romero-Isart, N., **278**, 10
Roper, G.C., **270**, 488
Rose, D.J., **268**, 163; **274**, 131; **281**, 70; **282**, 123
Rosini, C., **268**, 323
Rosini, G.P., **270**, 77, 537
Rossi, G., **272**, 168
Rossi, L.M., **281**, 111
Rossi, R., **272**, 267
Rossi, S., **272**, 274
Rothe, J., **270**, 95
Roundhill, D.M., **270**, 216
Rousset, C.J., **280**, 8
Roviello, A., **281**, 141
Rowe, S.M., **270**, 337
Royer, A.C., **268**, 211
Royer, E.C., **277**, 76; **279**, 30
Royo, P., **273**, 270; **280**, 1
Roznyatovsky, V.A., **280**, 328
Rubaijlo, V.L., **275–276**, 453
Rudy, R.I., **280**, 339
Ruffo, F., **281**, 141
Rugge, A., **279**, 200
Ruiz, J., **281**, 165
Ruiz, R., **268**, 263; **278**, 159
Ruiz-Pérez, C., **277**, 139
Ruiz-Ramírez, L., **273**, 167
Ruloff, R., **275–276**, 15
Ruminski, R.R., **281**, 1
Ruppa, K.B.P., **280**, 143
Rusanov, E.B., **268**, 93
Rusnak, F., **273**, 279
Russell, D.R., **274**, 177; **277**, 253
Russo, N., **273**, 72
Rüter, H.D., **275–276**, 334
Ryabov, A.D., **280**, 57
Rybak-Akimova, E.V., **270**, 151
Rybin, L.V., **280**, 243
Rybinskaya, M.I., **280**, 243
Ryu, K., **274**, 24
Rzączyńska, Z., **275–276**, 28

Sabat, M., **267**, 87; **282**, 237
Sadler, G.G., **268**, 31
Sadler, P.J., **272**, 261; **273**, 1, 8
Saito, Y., **283**, 44
Sajadi, S.A.A., **273**, 101; **283**, 193
Sakata, K., **281**, 190
Sakuragi, H., **273**, 204
Sakurai, H., **283**, 175
Sakurai, T., **275–276**, 289
Saladini, M., **268**, 205; **273**, 397
Salah El Fallah, M., **278**, 43
Sala Pala, J., **278**, 108
Salas, J.M., **274**, 53
Salas-Peregrín, J.M., **268**, 77; **277**, 103
Salvadori, P., **268**, 323; **275–276**, 462
Salvini, A., **272**, 141
Samaras, E., **272**, 24
Samuel, E., **279**, 210
Samuleev, P.V., **280**, 57
Sánchez, A., **267**, 169; **277**, 139
Sanchez, C., **279**, 144
Sánchez-Delgado, R.A., **275–276**, 528
Sanmartin, J., **274**, 73
Sanna, D., **268**, 297; **275–276**, 440; **283**, 202, 233
Sannazzaro, A.I., **273**, 367
Sano, K., **281**, 214
Sanseverino, M., **275–276**, 301
Santini, C., **282**, 1
Santos, A.M., **278**, 51
Santos, I., **271**, 65
Santos, R.H.A., **267**, 209
Sanz, V., **268**, 145; **271**, 174
Sapiña, F., **278**, 159; **279**, 58
Sardone, N., **272**, 68
Sarova, G.Ch., **273**, 403
Sartori, D.A., **268**, 311
Sarveswaran, V., **277**, 186
Sasakawa, Y., **283**, 184
Sasaki, T., **283**, 243
Sasaki, Y., **283**, 151
Sato, M., **267**, 329
Satoh, Y., **269**, 342
Sattelberger, A.P., **271**, 187
Saumweber, R., **269**, 83
Sava, G., **273**, 62
Sawluk, J., **269**, 317
Sawusch, S., **268**, 109; **269**, 350; **277**, 219
Sazikova, N., **270**, 169
Schacht, J., **273**, 85
Schaffner, K., **271**, 160
Schebler, P.J., **270**, 543
Scheidt, W.R., **278**, 197
Schejter, A., **273**, 388
Scherk, C., **275–276**, 65
Schettino, A., **273**, 62
Schilde, U., **268**, 109; **269**, 350; **277**, 219
Schilling, P.J., **280**, 211
Schilling, T., **270**, 95
Schimanski, A., **283**, 223
Schleis, T., **270**, 238
Schmid, R., **268**, 69
Schmidpeter, A., **269**, 7
Schmidt, H., **267**, 229
Schmidt, K., **269**, 135
Schmidt, M., **269**, 101; **275–276**, 65
Schmidt, T., **274**, 126
Schmittel, M., **275–276**, 435
Schneider, H., **279**, 186
Schneider, J.J., **281**, 53
Schneider, R., **275–276**, 150
Scholl, H.-J., **273**, 131
Schollmeyer, D., **273**, 54
Scholten, C., **274**, 137
Schoonover, J.R., **275–276**, 380
Schrock, R.R., **270**, 353
Schröder, A., **269**, 23
Schröder, D., **273**, 111
Schröder, G., **269**, 135
Schultz, F.A., **278**, 1
Schulze Tilling, A., **270**, 95
Schumacher, P., **267**, 133
Schumann, H., **280**, 21, 138
Schütte, M.T., **267**, 133
Schwarz, H., **273**, 111
Schwarz, W., **269**, 91
Scodinu, A., **277**, 236
Scott, B., **279**, 1
Scott, R.A., **273**, 409
Scozzafava, A., **275–276**, 295
Scudder, M.L., **274**, 32
Seevogel, K., **270**, 95
Seibig, S., **279**, 37
Seidelmann, O., **271**, 40
Seino, H., **280**, 163
Seiter, M., **273**, 354
Sellés, A., **273**, 270
Sellmann, D., **269**, 63; **272**, 211; **280**, 39
Sen, A., **270**, 8
Serroni, S., **275–276**, 320
Serveiss, D., **281**, 1
Server-Carrió, J., **279**, 58
Sessanta o Santi, A., **273**, 62
Sessoli, R., **271**, 9
Seybert, D.W., **277**, 1
Sgobino, A., **272**, 197
Shahinian, M.S., **275–276**, 510
Shaikh, S.N., **270**, 162
Shang, M., **278**, 197
Shapley, P.A., **267**, 335
Shaw, B.L., **270**, 312; **279**, 95
Shchegolikhina, O.I., **280**, 282
Shea, S.E., **281**, 43
Sheil, M.M., **281**, 10
Sheldrick, W.S., **267**, 39
Shellenbarger-Jones, A., **267**, 319
Shen, H., **267**, 223
Shen, J.-Y., **268**, 69
Shenton, D.J., **270**, 312
Shepherd, R.E., **267**, 11; **268**, 279, 287; **271**, 124; **277**, 46; **279**, 85
Sherry, A.D., **273**, 288
Shestakov, A.F., **280**, 288
Shi, N., **268**, 103
Shi, T., **282**, 55
Shibahara, T., **274**, 103; **282**, 50
Shibata, N., **283**, 184
Shidara, S., **275–276**, 289
Shields, G.P., **274**, 201
Shilov, A.E., **270**, 221; **280**, 288
Shilova, A.K., **270**, 221
Shim, B.Y., **269**, 337
Shiro, M., **270**, 424
Shishin, A.V., **267**, 293
Shivakumar, M., **275–276**, 551
Shklover, V., **274**, 64
Shoemaker, R.K., **281**, 153
Shokhireva, T.Kh., **272**, 204
Shoup, S.S., **274**, 192
Shubina, E.S., **280**, 302
Shubochkin, L.K., **280**, 339
Shur, V.B., **280**, 264
Shuvalova, N.I., **270**, 221
Siasios, G., **269**, 317; **281**, 64
Sidorov, S.N., **280**, 348
Sieler, J., **268**, 93
Sigalas, M.P., **273**, 72
Sigel, H., **273**, 101; **280**, 50; **283**, 193
Sikkink, R., **273**, 279
Silvestru, A., **271**, 75
Simándi, L., **281**, 235
Simándi, L.I., **270**, 326
Simão, J.E.J., **281**, 225
Simard, M.G., **282**, 30
Simões, P., **273**, 213
Simões Gonçalves, M.L.S., **278**, 51
Simon, E., **282**, 173
Simonneaux, G., **273**, 142
Sinclair, L., **278**, 1
Singer, A.L., **277**, 157
Sini, G., **280**, 26
Sinn, E., **278**, 217
Širac, S., **271**, 239
Sironi, A., **282**, 163
Sita, L.R., **270**, 118
Skopenko, V.V., **268**, 93
Slattery, S.J., **282**, 131
Slebodnick, C., **283**, 37
Sletten, E., **273**, 41
Sletten, J., **279**, 127
Sloan, J., **272**, 261
Slovokhotov, Y.L., **280**, 211
Slovokhotov, Yu.L., **280**, 275
Slugovc, C., **268**, 69
Smeets, W.J.J., **270**, 34

Smit, J.P., **267**, 49; **271**, 105; **278**, 202
Smith, B.E., **275–276**, 150
Smith, D.R., **275–276**, 359; **277**, 225
Smith, L.T., **268**, 151
Smith, R., **279**, 226
Smith, S.R., **273**, 316
Snoeck, T.L., **278**, 83
Soghomonian, V., **275–276**, 122
Sohn, Y.S., **279**, 116
Sokolov, M., **271**, 222
Sokolov, M.N., **269**, 292
Sola, M., **272**, 168; **273**, 397
Solan, G.A., **277**, 186
Solans, X., **268**, 263; **277**, 139; **278**, 43
Solari, E., **270**, 298
Söllner, R., **275–276**, 435
Solomon, E.I., **275–276**, 327
Soltek, R., **269**, 143
Somoza, F., **282**, 96
Son, H.L., **272**, 89
Son, M.-S., **272**, 33
Son, S.U., **281**, 229
Song, B., **273**, 101; **280**, 50; **283**, 193
Song, N.W., **279**, 243
Song, R., **272**, 228
Sono, M., **275–276**, 250
Sordo, J., **267**, 169
Soriano, A., **283**, 12
Soubra, C., **272**, 95
Soucek, M.D., **277**, 253
Sournies, F., **268**, 307
Sousa, A., **271**, 1; **274**, 131; **279**, 249; **281**, 70
Sóvágó, I., **275–276**, 440; **283**, 233
Spek, A.L., **267**, 183; **270**, 34; **279**, 217
Spiccia, L., **279**, 192
Spiro, T.G., **270**, 392
Spyroulias, G.A., **275–276**, 182
St. Pierre, T.G., **267**, 7
Stålhandske, C.M.V., **273**, 409
Stampfl, T., **274**, 143
Stankevich, I.V., **280**, 219
Staples, R.J., **271**, 49
Starikova, Z., **270**, 169
Stark, G.A., **269**, 167
Starke, I., **268**, 109
Stassinopoulou, C.I., **268**, 169; **281**, 148
Stavropoulos, G., **272**, 7
Steel, P.J., **269**, 229
Steimann, M., **269**, 13
Steinforth, W., **274**, 251
Steinhuebel, D.P., **270**, 527
Stelzer, N.A., **280**, 243
Stephens, A.K.W., **273**, 47
Stergiou, A.C., **269**, 332
Sterns, M., **271**, 151
Steuber, J., **275–276**, 52
Steunou, N., **279**, 144
Stirba, S.E., **278**, 61
Stöckigt, D., **282**, 61
Stockland, Jr., R.A., **271**, 236
Stolarov, I.P., **275–276**, 453
Stone, K., **282**, 131
Stoppioni, P., **273**, 151
Strähle, J., **270**, 177
Stramm, C., **269**, 111
Strange, R.W., **275–276**, 150
Strauch, P., **271**, 199; **278**, 118
Streib, W.E., **280**, 125
Strinna Erre, L., **272**, 68
Strotmann, M., **280**, 71
Strouse, G.F., **275–276**, 380
Struchkov, Y., **270**, 169
Struchkov, Yu.T., **280**, 243
Struck, G.E., **274**, 115
Stufkens, D.J., **278**, 83
Su, W., **267**, 313; **274**, 229; **277**, 171
Suarez-Varela, J., **281**, 95
Sue, R.E., **273**, 8
Suenaga, Y., **271**, 129, 145; **281**, 116
Sugimori, T., **278**, 223, 253; **283**, 275
Sugimura, Y., **283**, 184
Suh, M.P., **269**, 337
Süli-Vargha, H., **283**, 233
Sülü, M., **270**, 499
Sun, B.-W., **279**, 69, 76
Sun, S., **268**, 103
Sun, Y., **281**, 60
Sundberg, M.R., **267**, 249; **268**, 21
Sung, Y.K., **272**, 33
Sünkel, K., **269**, 111
Supuran, C.T., **273**, 334; **275–276**, 295
Suranna, G.-P., **280**, 150
Surdy, P., **268**, 297
Surville-Barland, C., **278**, 159
Suwannachot, Y., **272**, 89
Suzuki, A., **275–276**, 159
Suzuki, E., **283**, 184
Suzuki, M., **283**, 124
Suzuki, N., **280**, 8
Suzuki, S., **275–276**, 289; **283**, 118, 260
Suzuki, T., **281**, 77
Svensson, J., **278**, 24
Svoboda, I., **278**, 127
Sykes, A.G., **271**, 228
Szulbinski, W.S., **269**, 253
Szulc, A., **270**, 440

Tabbi, G., **282**, 180
Tachikawa, H., **281**, 85
Tadsanaprasittipol, A., **278**, 143
Tafesse, F., **269**, 287
Tagawa, Y., **283**, 260
Tajima, K., **283**, 61
Takada, J., **283**, 175
Takagi, H.D., **277**, 31
Takahashi, K., **281**, 174; **283**, 72
Takahashi, T., **280**, 8
Takalo, H., **267**, 63
Takamizawa, S., **283**, 118, 268
Takats, J., **277**, 61
Takeda, S., **283**, 268
Takeuchi, T., **271**, 13
Takimura, T., **275–276**, 159
Taler, G., **273**, 388
Talman, E.G., **283**, 251
Tamburini, S., **272**, 235
Tamura, H., **279**, 165; **283**, 175
Tan, X.S., **277**, 21
Tanabe, Y., **280**, 163
Tanaka, M., **267**, 147; **270**, 459
Tanaka, T., **275–276**, 222
Tang, W., **275–276**, 58
Tang, W.X., **277**, 21
Tang, Z., **267**, 73
Tangoulis, V., **272**, 24
Tani, K., **270**, 459
Tanner, D.B., **282**, 90
Tarasconi, P., **269**, 297
Tasić-Stojanović, S., **278**, 66
Tatsumi, K., **283**, 136
Tauzher, G., **272**, 74
Tavagnacco, C., **270**, 145; **280**, 322
Tavares, P., **273**, 409
Taylor, M.J., **277**, 26
Tedesco, E., **273**, 116
Tejero, R., **275–276**, 73
Teleshev, A.T., **267**, 293
Temkin, O.N., **280**, 202
Terzis, A., **272**, 283
Tesche, B., **270**, 95
Teterin, A.Yu., **273**, 234
Teterin, Yu.A., **273**, 234
Teuben, J.H., **271**, 180
Teunis, C., **275–276**, 98
Thaler, A., **279**, 186
Theophanides, T.M., **273**, 201
Theriault, A., **281**, 43
Therien, M.J., **275–276**, 427
Thiel, R., **269**, 23
Thompson, L.K., **279**, 121
Thorburn, I.S., **270**, 130
Thornton-Pett, M., **270**, 312; **279**, 95
Thorp, H.H., **273**, 316
Tian, X., **272**, 295
Tiekink, E.R.T., **269**, 317; **274**, 53; **279**, 192; **281**, 64
Tiripicchio, A., **269**, 191; **273**, 320; **275–276**, 500
Tisato, F., **275–276**, 401
Tittsworth, R.C., **280**, 211
Tjan, F.S.B., **268**, 43
Tokii, T., **274**, 224; **277**, 134
Tollari, S., **272**, 18
Tomasin, P., **272**, 235
Tomizawa, H., **267**, 147
Tommasi, I., **272**, 38
Tondello, E., **275–276**, 340
Torres, J.L., **278**, 10
Torres, M.R., **282**, 230
Toscano, M., **273**, 72
Touchard, D., **272**, 188; **280**, 118
Toupet, L., **272**, 188
Tozer, R., **275–276**, 203
Traeger, J.C., **267**, 281
Traldi, P., **282**, 119
Trautwein, A.X., **271**, 9; **275–276**, 334; **283**, 111
Tribbitt, S.A., **268**, 329
Trifonova, O.I., **280**, 328
Trifunović, S.R., *278*, 209
Tripepi, G., **270**, 177
Trogu, E.F., **267**, 115; **273**, 175
Troyanov, S.I., **271**, 180
Trylus, K.-H., **269**, 23
Tsai, T.Y.R., **267**, 81
Tsang, S.C., **272**, 261
Tsiamis, C., **269**, 332
Tsikalova, M., **270**, 169; **280**, 295
Tsiveriotis, P., **272**, 7
Tsuge, A., **281**, 190
Tsumura, S., **283**, 275
Tuchagues, J.-P., **279**, 69, 76
Tunice, C., **272**, 18
Tunik, S.P., **268**, 189
Turner, D.L., **273**, 196
Turpin, R., **271**, 217
Tušek-Božić, L., **282**, 76
Tweedle, M.F., **275–276**, 106
Tylli, H., **268**, 21

Ubeda, M.A., **278**, 61
Ueda, T., **281**, 214
Uehara, A., **283**, 124
Ueki, M., **283**, 243
Ueyama, N., **275–276**, 43; **283**, 91
Ugo, R., **270**, 479; **280**, 282
Ugozzoli, F., **267**, 177; **273**, 259
Uguagliati, P., **275–276**, 385; **277**, 247
Uhlemann, E., **268**, 109; **269**, 350; **277**, 219
Uhrhammer, D., **278**, 1
Ujimoto, K., **282**, 247
Ulstrup, J., **272**, 109
Umakoshi, K., **283**, 151
Umetani, S., **267**, 201
Underhill, A.E., **279**, 65
Unger, C., **267**, 133
Unoura, K., **269**, 260
Urbanos, F.A., **277**, 76; **279**, 30
Urbina, J.A., **275–276**, 528
Ushiyama, M., **283**, 243
Ustynyuk, N.A., **280**, 249, 328
Ustynyuk, Yu.A., **280**, 328
Usui, Y., **283**, 51
Utsuno, S., **281**, 221
Uytterhoeven, M.G., **274**, 90

Vahrenkamp, H., **272**, 115; **282**, 193
Vaid, T.P., **270**, 414
Valencia, L., **282**, 42
Valensin, G., **275–276**, 274
Valente, E.J., **268**, 151
Valentin, B., **277**, 98
Valetsky, P.M., **280**, 348
Valko, M., **278**, 127
Valle, G., **275–276**, 410, 419; **282**, 1
Valliant, J.F., **274**, 137

Valls, E., **275–276**, 521
Valtancoli, B., **268**, 63; **273**, 326
van Bekkum, H., **268**, 249; **277**, 193
Van Calcar, P.M., **270**, 28
Vančová, V., **277**, 127
van der Baan, J.L., **269**, 203
Vander Elst, L., **273**, 92
van der Heijden, A.M., **268**, 249; **277**, 193
van der Plas, J.L., **267**, 193
VanDerveer, D.G., **274**, 192
van Eldik, R., **278**, 32; **279**, 37
Van Engen, D., **279**, 7
Van Haandel, M.J.H., **275–276**, 98
van Koningsbruggen, P.J., **268**, 37; **273**, 54; **274**, 1
van Koten, G., **269**, 322
Vargaftik, M.N., **275–276**, 453; **280**, 339
Vargas, F., **275–276**, 528
Vargas, J., **267**, 151
Várnagy, K., **275–276**, 440; **283**, 233
Vašák, M., **272**, 150
Vázquez de Miguel, A., **273**, 270
Vecchio, G., **275–276**, 395
Veeger, C., **275–276**, 98
Veghini, D., **280**, 226
Velders, A.H., **273**, 259
Veldman, N., **267**, 183
Veltsistas, P., **267**, 299
Venanzi, L.M., **270**, 499
Vencato, I., **281**, 111
Vernaglione, G., **275–276**, 295
Vértes, A., **274**, 167
Vicente, R., **269**, 313; **278**, 43
Vicentini, G., **274**, 122, 236
Vickers, P.W., **270**, 382
Viezzoli, M.S., **269**, 125; **273**, 131
Vigato, P.A., **272**, 235
Vignali, C., **275–276**, 320
Vila, A.J., **273**, 367
Villagrán, M., **273**, 191
Vincent, J.B., **268**, 211
Virovets, A., **271**, 222
Virovets, A.A., **271**, 228
Virovets, A.V., **269**, 292
Visentin, F., **275–276**, 385; **277**, 247
Vitt, S., **280**, 355
Vittadini, A., **275–276**, 340
Vitulli, G., **268**, 323
Vivas-Mejía, P.E., **271**, 137
Vivo, A., **281**, 141
Vizza, F., **272**, 55
Vlček Jr., A., **278**, 83
Vogel, W., **270**, 95
Vogler, R., **272**, 115
Voigt, A., **271**, 199
Vol'pin, M.E., **280**, 264
von Zelewsky, A., **271**, 36; **272**, 153
Vorontsov, E.V., **278**, 122; **280**, 302
Vorontzov, E.V., **280**, 233
Vrieze, K., **270**, 34
Vuano, S., **272**, 74

Wada, Y., **272**, 80
Wadepohl, H., **274**, 201
Wagner, T.R., **282**, 155
Walcher, D., **273**, 54
Walker, F.A., **272**, 204
Walsh, K.G., **280**, 30
Walther, D., **281**, 181
Walton, P.H., **269**, 241
Walton, R.A., **267**, 123; **278**, 245
Wan, X., **275–276**, 106
Wang, G.-L., **279**, 69, 76
Wang, H.H., **273**, 175
Wang, J.-C., **269**, 302; **274**, 7
Wang, K., **270**, 537
Wang, R., **280**, 150
Wang, Y., **279**, 233
Wang, Z.-Q., **280**, 62
Ward, M.D., **267**, 1, 239, 323; **278**, 178
Warda, S.A., **268**, 117
Warren, C.J., **282**, 123
Wasfi, S.H., **268**, 329; **278**, 91; **282**, 136
Wasgestian, F., **274**, 126
Washington, J., **270**, 111
Watanabe, H., **283**, 175
Waters, J.M., **267**, 27; **277**, 37
Watson, K.J., **267**, 159
Wayland, B.B., **270**, 197
Webb, J., **267**, 7; **275–276**, 37
Webster, L., **271**, 151
Webster, M., **271**, 203
Wei, H.H., **279**, 233
Weigand, W., **269**, 83; **273**, 106
Weiß, A., **269**, 47
Weiss, R., **275–276**, 334; **283**, 98
Weitzel, M., **268**, 211
Wen, R., **282**, 96
Wendt, O.F., **277**, 236
Werner, H., **275–276**, 435
Werner, R., **278**, 190; **281**, 111
Werner, W.L.J., **269**, 203
Westerhausen, M., **269**, 91
Weston, W.S., **280**, 99
Wever, R., **273**, 160
Weyhermüller, T., **268**, 43; **274**, 111
Whang, D., **279**, 238
Wharf, I., **282**, 30
Wheatley, N., **278**, 229
Wheeler, D.E., **271**, 167
White, A.J.P., **281**, 201
White, P.S., **269**, 1; **270**, 511
Wiede, P., **268**, 69
Wieghardt, K., **268**, 43; **274**, 111
Wilke, G., **270**, 273
Willett, R.D., **279**, 1
Williams, A.F., **275–276**, 279
Williams, D.J., **272**, 131; **281**, 201
Williams, D.S., **270**, 511
Williams, J.M., **273**, 175
Williams, P.A.M., **268**, 239
Wilson, K., **273**, 213
Wilson, L.J., **275–276**, 106
Winkler, H., **275–276**, 334; **283**, 111
Winkler, J.R., **275–276**, 470
Wittrig, R.E., **270**, 111
Wocadlo, S., **268**, 117
Wojakowska, H., **269**, 326
Wojcinski II, L.M., **270**, 8
Wolczanski, P.T., **270**, 414
Wolny, J.A., **272**, 153
Wolowiec, S., **269**, 203; **277**, 242
Wong, K.-Y., **272**, 176
Wong, W.T., **270**, 247; **279**, 172
Wong, W.-Y., **279**, 172
Woollins, J.D., **267**, 299
Woska, D.C., **270**, 197
Wu, B., **271**, 99
Wu, B.-M., **277**, 21
Wu, D., **267**, 313; **271**, 93
Wu, J.-Z., **283**, 17
Wu, L.P., **268**, 317
Wu, M., **267**, 81
Wu, P., **268**, 103
Wünsch, R., **273**, 106

Xavier, A.V., **272**, 261; **273**, 196
Xi, R., **270**, 118
Xiang, D.F., **277**, 21
Xin, X., **279**, 172
Xu, L., **277**, 130; **282**, 261
Xu, Q., **274**, 103
Xu, Y., **282**, 10; **283**, 17

Yablokov, Y.V., **275–276**, 28
Yagi, M., **281**, 221
Yajima, T., **278**, 136
Yamada, Y., **275–276**, 43; **283**, 91
Yamagishi, T., **267**, 329
Yamaguchi, K., **275–276**, 289; **283**, 260, 268
Yamaguchi, M., **267**, 329
Yamamoto, M., **268**, 317; **271**, 129
Yamamoto, T., **269**, 310
Yamamura, T., **283**, 243
Yamane, H., **283**, 118
Yamashita, J., **281**, 190
Yamauchi, O., **275–276**, 90; **278**, 136
Yamazaki, A., **269**, 260
Yamazaki, Y., **281**, 214
Yan, S., **273**, 393
Yan, S.G., **271**, 145
Yandulov, D.V., **280**, 125
Yang, G., **283**, 17
Yang, J., **271**, 75
Yang, W.-K., **274**, 7
Yano, S., **279**, 51
Yanovsky, A., **270**, 169
Yanovsky, A.I., **268**, 313; **280**, 233, 243
Yao, W., **280**, 26
Yap, G.P.A., **267**, 309; **270**, 559; **280**, 143, 172
Yasui, H., **283**, 175
Yatsimirsky, A.K., **273**, 167
Yeh, A., **267**, 81
Yoder, J.C., **274**, 115
Yokochi, A., **275–276**, 447, 541
Yokoyama, K., **282**, 209
Yokoyama, M., **279**, 51
Yonemura, M., **283**, 72
Yordanov, A.T., **270**, 216
Yordanov, N.D., **272**, 291
Yoshida, H., **281**, 207
Yoshida, N., **272**, 162
Yoshida, S., **275–276**, 222
Yoshikawa, Y., **271**, 24
Yoshino, Y., **282**, 38
Yoshizaki, F., **283**, 184
You, X.-Z., **271**, 99
Yuen, T., **273**, 310
Yuge, H., **279**, 105

Záhonyi-Budó, É., **281**, 235
Zakeeruddin, S.M., **274**, 64
Zakrzewski, J., **278**, 101
Zamora, F., **267**, 87; **282**, 237
Zanello, P., **273**, 24; **281**, 53
Zangrando, E., **277**, 111; **279**, 152
Zani, A., **275–276**, 117
Zanobini, F., **272**, 1
Zanotti-Gerosa, A., **270**, 298
Zehnder, M., **275–276**, 359
Zeiske, T., **269**, 47
Zeng, D., **279**, 172
Zevaco, T.A., **269**, 283
Zhang, B., **282**, 10
Zhang, F., **278**, 136
Zhang, J., **269**, 211
Zhang, Q., **271**, 93; **277**, 171
Zhang, W., **271**, 93
Zhang, X.X., **278**, 6
Zhang, Y., **277**, 263
Zhang, Z.-Z., **268**, 177
Zhao, D., **269**, 162
Zhdanov, A.A., **280**, 282
Zhen, Y., **271**, 93
Zhong, Z., **280**, 183
Zhong, Z.J., **271**, 99
Zhou, J.-Y., **283**, 17
Zhou, X., **270**, 551
Zhu, D., **268**, 103; **275–276**, 58
Zickgraf, A., **275–276**, 203
Zilverentant, C.L., **282**, 257
Zonnedda, G., **273**, 175
Zoppellaro, G., **282**, 180
Zou, X., **267**, 305
Zsolnai, L., **269**, 143
Zubavichus, Y.V., **280**, 211
Zubavichus, Ya.V., **280**, 275
Zubieta, J., **268**, 163, 257; **271**, 1; **274**, 131; **275–276**, 122; **277**, 263; **281**, 70; **282**, 123
Zubkowski, J.D., **268**, 151
Zucchi, C., **280**, 282

ELSEVIER

Inorganica Chimica Acta 283 (1998) 293–364

Inorganica Chimica Acta

Cumulative Subject Index of Volumes 267–283

Ab initio calculations
five-coordinate platinum(II) alkyne complexes: synthesis, ab initio calculations and crystal and molecular structure of $[PtI_2(Me_2phen)(\eta^2\text{-}PhC{\equiv}CPh)]\cdot CHCl_3$, **275–276**, 500
Ab initio computations
the cations $[ICNI]^+$ and $[H_3N\text{–}NH_3]^{2+}$ — natural bond orbital analysis and some valence bond considerations, **269**, 1
Ab initio MO
structures and electronic states of the electron donor–acceptor (EDA) complexes: trimethylamine–halogen complex $(CH_3)_3N\cdots F_2$, **281**, 85
Acetate complexes
weak intermolecular antiferromagnetic exchange in $\{[Ru_2(O_2CCH_3)_4(L)]X\}_n$ polymers (L = 4,4′-dipyridine and 1,4-diazabicyclooctane, $X = PF_6^-$ or BPh_4^-), **279**, 121
solid-state synthesis of cluster compound $[MoCu_3OS_3(PPh_3)_3(CH_3COO)]$ with its structure, reactivity and optical non-linearities, **279**, 172
synthesis, structure and characterization of dinuclear nickel(II) complexes containing the derivatives of *N,N,N′,N′*-tetrakis[(2-benzimidazolyl)methyl]-2-hydroxy-1,3-diaminopropane and an acetate or a propionate ion, **283**, 118
Acetato-bridged complexes
preparation and properties of a series of (μ-oxo)di(μ-acetato) diruthenium(III) complexes, $[Ru_2(\mu\text{-}O)(\mu\text{-}CH_3COO)_2(pyridine)_4(L)_2]^{2+}$ and $[Ru_2(\mu\text{-}O)(\mu\text{-}CH_3COO)_2(AA)_2(L)_2]^{2+}$, where AA is 2,2′-bipyridine or 1,10-phenanthroline, and L is a monodentate ligand, **283**, 151
Acetylacetonate complexes
synthesis, characterization and application of Ru(BINAP)(Acac)(MNAA)(MeOH), a highly effective catalyst for the asymmetric hydrogenation of 2-(6′-methoxynaphth-2′-yl)acrylic acid, **270**, 247
Acetylacetonato complexes
redox behavior of a binuclear ruthenium complex having a di-μ-nitrosyl ligand, $[\{Ru(acac)_2\}_2(\mu\text{-}NO)_2]$ (acac = acetylacetonato), **269**, 342
Acetyl complexes
rhodium aryloxide complexes containing terdentate nitrogen ligands, C–O bond formation, hydrogen bonding with phenol and oxidative addition reactions. Molecular structure of an Rh(III)–acetyl complex, **270**, 34
Acetylene reduction
catalytic reduction of acetylene in the presence of molybdenum and iron clusters, including FeMo cofactor of nitrogenase, **270**, 221
Acidity constants
metal ion-coordinating properties of imidazole and derivatives in aqueous solution: interrelation between complex stability and ligand basicity, **280**, 50
Acids and bases
metal ion stabilities correlate with electron affinity rather than hardness or softness, **283**, 30
Acridine complexes
intercalator-linked cisplatin: synthesis and antitumor activity of *cis*-dichloroplatinum(II) complexes connected to acridine and phenylquinolines by one methylene chain, **279**, 51
Acrylate anion complexes
synthesis, characterization and application of Ru(BINAP)(Acac)(MNAA)(MeOH), a highly effective catalyst for the asymmetric hydrogenation of 2-(6′-methoxynaphth-2′-yl)acrylic acid, **270**, 247
Actinide complexes
calculation of NMR paramagnetic shift temperature dependence in actinide methyl borohydrides, **272**, 43
functionalized acyclic Schiff bases and related complexes with d- and f-metal ions, **272**, 235
Active carbon supported complexes
effect of ruthenium precursor on hydrogen-treated active carbon supported ruthenium catalysts for ammonia synthesis, **280**, 183
Adenine
electrochemistry of dioxygen activation in the presence of adenine, **280**, 322
A-frame complexes
an unusual cyanide-bridged extended A-frame complex: the molecular structure of $[\{Pd_2Br_2(\mu\text{-}PPh_2)(\mu\text{-}dppm)\}_2(\mu\text{-}CN)_2]$, **271**, 236
Aggregation phenomena
interaction of organometallic cationic complex ions containing terpyridine ligands with nucleic acids: an investigation on aggregative phenomena, **275–276**, 242
Alakane oxidation
azide promotion of alkane oxidations catalyzed by metal complexes in solution, **270**, 162
Alizarin complexes
linkage isomers of alizarin-bis(bipyridine)ruthenium(II), **281**, 126
Alkali metal compounds
metal cluster topology. 19. Beyond the auride ion: triangulated gold networks and ethane-like structural units in binary and ternary alkali metal gold intermetallics, **277**, 202
Alkali metal ion complexes
thermodynamic study of complexation of alkali and alkaline-earth metal ions with benzene-containing cryptands, **278**, 6
Alkaline complexes
NMR conformational study of diamagnetic complexes of some triaza-triacetate macrocycles, **273**, 288
Alkaline cytochrome *c*
effect of pyridine on the alkaline isomerization of cytochrome *c*, **275–276**, 58
Alkaline-earth complexes
NMR conformational study of diamagnetic complexes of some triaza-triacetate macrocycles, **273**, 288
Alkaline-earth metal ion complexes
thermodynamic study of complexation of alkali and alkaline-earth metal ions with benzene-containing cryptands, **278**, 6
Alkanes

Elsevier Science S.A.

unprecedented facile reactions of propane and cycloalkanes with elemental sulfur leading to dialkylsulfides and dicycloalkylsulfides, **280**, 355

Alkene complexes
the kinetics and mechanism of ligand exchange in photogenerated (η^1-bromobenzene)$Cr(CO)_5$/alkene systems, **267**, 49
coordination of a disubstituted alkene in a chelated d^0-metal–alkyl–alkene complex. Evidence for equilibrium between complexes with coordinated and free alkenes, **270**, 5
reaction of the dinuclear cobalt complex $[Co_2\{\mu\text{-}PPh_2CHCHC(O)\text{-}CHCHPPh_2\}(CO)_4]$ with two-electron donor ligands; the stabilisation of cobalt–alkene bonds by incorporation of alkenes into a bridging diphosphine ligand, **277**, 186

Alkoxide-bridged complexes
crystal structure, spectral and magnetic properties of a new (μ-acetate)(μ-alkoxide) dicopper(II) complex as a model for tyrosinase, **281**, 111

Alkoxide complexes
synthesis and single crystal X-ray structure of a novel trinuclear copper(II) methoxide complex, **269**, 241
elimination of benzene from a benzyloxy ligand to form a carbonyl group on a tungsten-phosphine center, **270**, 202
the propensity of alkoxide and aryloxide derivatives of tungsten carbonyls to aggregate in solution. Synthesis and X-ray structures of dinuclear, trinuclear and tetranuclear complexes derived from the $MeOW(CO)_5^-$ anion, **270**, 405
reactions of $Ti(O\text{–}i\text{-}Pr)Cl_3$ with 2-aminophenol and the crystal structure of the zwitterionic complex $[Ti(O\text{–}i\text{-}Pr)Cl_3(THF)\text{-}(^-OC_6H_4NH_3^+)]\cdot THF$, **278**, 232

Alkoxy compounds
alkoxymesitylchlorogermanes and aryloxymesitylchlorogermanes, **277**, 98

Alkylcobalamins
Co–C force constants from resonance Raman spectra of alkylcobalamins: insensitivity to dimethylbenzylimidazole coordination, **270**, 392

Alkyl complexes
coordination of a disubstituted alkene in a chelated d^0-metal–alkyl–alkene complex. Evidence for equilibrium between complexes with coordinated and free alkenes, **270**, 5
palladium(II)-catalyzed alternating copolymerization of alkenes with sulfur dioxide, **270**, 8
titanium–carbon functionalities on an oxo surface defined by a calix[4]arene moiety and its redox chemistry, **270**, 298
extracting absolute titanium–alkyl and –hydride bond enthalpies from relative $D(TiR(H))$ in $(silox)_2(^tBu_3SiNH)TiR$: electronegativity and *ECT* models, **270**, 414
synthesis of chloro and methyl imido cyclopentadienyl molybdenum and tungsten complexes. X-ray molecular structures of $[WCp^*Cl_3(N^tBu)]$, $[MoCp^*ClMe_2(N^tBu)]$ and $[WCp^*ClMe_2(N^tBu)]$, **273**, 270
tetravalent niobium alkyls and enolate stabilized by anionic organic amides, **280**, 143
preparation of doubly-silylene-bridged zirconocene alkyl complexes, $(Me_2Si)_2\{\eta^5\text{-}C_5H_2\text{-}4\text{-}CHMe_2\}\{\eta^5\text{-}C_5H\text{-}3,5\text{-}(CHMe_2)_2\}ZrR_2$ (R = CH_3, CH_2Ph) and investigations of their activity in 1-pentene polymerization. Molecular structure of $(Me_2Si)_2\{\eta^5\text{-}C_5H_2\text{-}4\text{-}CHMe_2\}\{\eta^5\text{-}C_5H\text{-}3,5\text{-}(CHMe_2)_2\}Zr(CH_2Ph)_2$, **280**, 226
electrophoretic study of cationic cobalt(III) complexes with $[N_2O]$ and $[N_2O_2]$ Schiff base ligands, **280**, 295

Alkyl halide complexes
chelated alkyl halide manganese complexes $(\eta^5\text{:}\eta^1\text{-}C_5H_4CH_2CH_2X)\text{-}Mn(CO)_2$ from photolysis of $(\eta^5\text{-}C_5H_4CH_2CH_2X)Mn(CO)_3$, **280**, 316

Alkylidene complexes
protonation of Fischer-type alkylidyne carbonyltungsten complexes. Structural comparison of alkylidyne and alkylidene metal complexes, including a neutron diffraction study of $[W(CHCH_3)\text{-}Cl_2(CO)(PMe_3)_2]$, **279**, 7

Alkylidenephosphorane complexes
comparison of the neutral benzyldiphenylphosphine complexes $C_6H_6Cr(CO)_2PPh_2Bz$ and $C_5H_5Mn(CO)_2PPh_2Bz$ with the isoelectronic manganiobenzyldiphenylphosphonium salts $[C_5R_5Mn(CO)(NO)PPh_2Bz]BF_4$, **272**, 101

Alkyne complexes
preparation of trimethylsilyl alkyne complexes of bis(pentamethylcyclopentadienyl)zirconium, $(\eta^5\text{-}C_5Me_5)_2Zr(RC{\equiv}CSiMe_3)$, and their regioselective reactions with nitrous oxide, **270**, 399
five-coordinate platinum(II) alkyne complexes: synthesis, ab initio calculations and crystal and molecular structure of $[PtI_2(Me_2phen)(\eta^2\text{-}PhC{\equiv}CPh)]\cdot CHCl_3$, **275–276**, 500
reactions of platinum(II)–alkyne and –alkynyl complexes with the hydroxy compounds $HO(CH_2)_2X$ (X = Br, I, OH), **275–276**, 515
the synthesis, molecular structure and fluxional properties of the 1-phenylpropyne complex $[WI_2(CO)\{Ph_2P(CH_2)PPh_2\}(\eta^2\text{-}MeC_2Ph)]$, **279**, 65

Alkyne hydrogenation
solution and crystal structure of the dihydrogen complex $[Ru(H_2)(H)(PMe_2Ph)_4]PF_6$, an active alkyne hydrogenation catalyst, **270**, 238

Alkyne trimerization
synthesis, structure, spectroelectrochemical and theoretical investigations of $[(\eta^5\text{-}Me_5Cp)\underline{Co(\eta^4\text{-}MeC{=}CMe\text{–}CMe{=}CMe)Sn}\text{-}\{CH(SiMe_3)_2\}_2]$. Formation of a cobalt complexed stannacycle via a mixed alkyne–stannylene cycloaddition reaction, **281**, 53

Alkynyl complexes
mono- and bis-alkynyl ruthenium(II) complexes containing the ferrocenyl moiety; crystal structure of *trans*-$[Ru(C{\equiv}CC_5H_4FeC_5H_5)_2\text{-}(Ph_2PCH_2CH_2PPh_2)_2]$ and electrochemical studies, **272**, 188
reactions of platinum(II)–alkyne and –alkynyl complexes with the hydroxy compounds $HO(CH_2)_2X$ (X = Br, I, OH), **275–276**, 515
ammonia ruthenium complexes for the access to new hydrido, carbonyl and isocyanide ruthenium-alkynyl derivatives, **280**, 118

Allenylidene complexes
study of the relative π-acceptor ability of unsaturated ligands $:C(=C)_nPh_2$ (n = 0, 1, 2 and 4) based on cyclic voltammetry of complexes *trans*-$[RhCl(L)(P^iPr_3)_2]$ (L = $:C(=C)_nPh_2$, C_2H_4 and CO), **275–276**, 435

Allyl complexes
η^3-allyl–cobalt(II) complexes, **270**, 60
bis(η^3-allyl)(η^8-pentalene)zirconium: preparation, structure and structural dynamics, **270**, 555
novel palladium(II) allyl complexes with nitrogen–sulfur donor bidentate ligands. Mechanism of allyl amination of $[Pd(\eta 3\text{-allyl})\text{-}(N\text{-}SR)]ClO_4$ (allyl = C_3H_5; N-SR = C_5H_4N-2-CH_2SR, R = C_6H_5, C_2H_5) in the presence of activated olefins. X-ray structure determination and fluxional behavior, **275–276**, 385
crystal structure and fluxional behavior of novel Pd(II) allyl complexes containing a potentially tridentate S,N,S ligand: an example of selective *syn–anti* isomerism, **277**, 247

Aluminium complexes
fixation of carbon dioxide with 1,2-epoxypropane catalysed by tetra-t-butylphthalocyaninatoaluminium(III) acetate, tetra-t-butylphthalocyaninatoaluminium(III) trimethylsilanolate and tetra-t-butylphthalocyaninatoaluminium(III) triphenylsilanolate, **278**, 223

Aluminophosphate complexes
LMU-3: a new cobalt aluminophosphate with exclusively five-coordinated aluminium and octahedrally coordinated cobalt, **269**, 73

Aluminum alkyls
alkyl exchange between aluminum trialkyls and zirconocene dichloride complexes — a measure of electron densities at the Zr center, **270**, 376

Alumoxane complexes

zirconocene–alumoxane (1:1) – a catalyst for the selective dimerization of α-olefins, **270**, 20

Amide complexes
synthesis of *mer*-trichlorotris(tetrahydrofuran)tungsten(III) by intermetal chlorine atom transfer, **278**, 103
tetravalent niobium alkyls and enolate stabilized by anionic organic amides, **280**, 143
binuclear nickel(II) and copper(II) complexes of amide-containing two-armed and macrocyclic ligands, **282**, 222

Amide ligand complexes
synthesis, structure, and properties of {*N,N'*-bis[2-(1-pyrazolyl)-ethyl]pyridine-2,6-dicarboxamido}copper(II), **269**, 269

Amidic acid complexes
synthesis and rearrangements of the pentaamminecobalt(III) linkage isomers of some amidic acids, **268**, 85

Amidocarbenes
mononuclear and binuclear *o*-xylylene derivatives of Pd(II), **269**, 191

Amine complexes
second sphere coordination behavior of aquo and amine ligands bound to a η^6-benzeneruthenium(II) cation, **270**, 87
reversible β-hydride elimination of amine ligands forming stable *cis*-η^2-iminium hydride Os^{II} complexes, **270**, 337
structures and electronic states of the electron donor–acceptor (EDA) complexes: trimethylamine–halogen complex $(CH_3)_3N\cdots F_2$, **281**, 85
the phenomenon of conglomerate crystallization. Part 51. The crystallization behavior of nickel dinitro amine complexes. *Trans*-[Ni(3,2,3-tet)$(NO_2)_2$]·H_2O (**I**), *mer*-[Ni(N5-Me-1,5,9-triazanonane)$(NO_2)_2$ (**II**), *cis*-[Ni(tren)$(NO_2)_2$]·H_2O (**IiI**) and *cis*-[Ni(edda)$(H_2O)_2$]·H_2O (**IV**) and control of the bonding mode of the nitrite ligand by the choice of the amine, **282**, 96
can Pt(IV)–amine complexes act as 'prodrugs'?, **283**, 251

Amine-phosphine complexes
the reactivity of five-coordinate Ru(II)(1,4-bis(diphenylphosphino)-butane) complexes with the N-donor ligands: ammonia, pyridine, 4-substituted pyridines, 2,2′-bipyridine, bis(*o*-pyridyl)amine, 1,10-phenanthroline, 4,7-diphenylphenanthroline and ethylenediamine, **267**, 209

Amine-phosphonato complexes
the dissociation kinetics of the ethylenediaminetetrakis(methylenephosphonato) complex of europium(III), **278**, 185

Amino-acetate ligand complexes
structural information on *trans*-1,2-diaminocyclohexane-*N,N'*-tetraacetateferrate(III) in the solid and aqueous phase, **279**, 37

Amino acid complexes
novel trigonal-bipyramidal copper(II) complexes containing one discrete amino acid chelate: crystal structures of [Cu(L-val)(pmdt)]ClO_4 and [Cu(aiba)(pmdt)]$ClO_4\cdot H_2O$ (L-val = L-valinate ion, aiba = α-aminoisobutyrate ion, and pmdt = *N,N,N′,N″,N″*-pentamethyldiethylenetriamine), **274**, 247
crystal structure, EPR and magnetic susceptibility studies of tetrakis[μ-(β-alanine)-*O,O′*]dichlorodicopper(II) dichloromonohydrate, **275–276**, 28
L- and D-cysteine derivatives of β-cyclodextrin: different molecular recognition properties of their copper(II) complexes for amino acids, **275–276**, 395
electrostatic ligand–ligand interactions in ternary copper(II) complexes with 3,5-diiodo-L-tyrosine and polar amino acids, **278**, 136
structures and electrochemical properties of the Co(III) ternary complexes containing NO_3-type tripodal tetradentate ligands and amino acids: effect of the outer coordination sphere on the electrochemical properties, **283**, 160

Amino acid derivative complexes
potentiometric and spectroscopic studies of transition metal complexes of bis(imidazolyl) and bis(pyridyl) derivatives of amino acids, **283**, 233

Amino acid esters
enantioselective cleavage of activated amino acid esters promoted by chiral palladacycles, **280**, 57

Amino acid porphyrin complexes
the synthesis and spectroscopic characterization of a chiral zinc(II) porphyrin dimer containing an amino acid bridge: model complexes for the induced circular dichroism of porphyrin protein supramolecules, **272**, 295

Amino acids
salt-induced peptide formation from amino acids in the presence of clays and related catalysts, **272**, 89
homology predicted structure and comparison with the secondary structure from NMR data for plastocyanin from the cyanobacterium *Synechocystis* sp. PCC 6803, **275–276**, 73

Aminocyclodextrin complexes
synthesis and antitumor activity of platinum(II) complexes of aminocyclodextrin, **272**, 162

Aminodiphosphine ligands
concomitant coordination of vinylidene and carbene ligands made possible at ruthenium(II), **272**, 1

Amino ester complexes
oxidation of amino esters to imine complexes with chiral high-valent oxoruthenium porphyrins, **273**, 142

Aminoglycoside toxicity
NMR studies of iron–gentamicin complexes and the implications for aminoglycoside toxicity, **273**, 85

Aminoglycosidic antibiotic
kanamycin revisited: a combined potentiometric and spectroscopic study of copper(II) binding to kanamycin B, **275–276**, 546

Aminolysis
an amazing example of yin–yang aminolysis in cyclophosphazenes, **268**, 307

Aminomethylpyridine complexes
bis(2-aminomethylpyridine) complexes of Cr(III) and Co(III), **277**, 177

Aminophosphine complexes
chelate ring-opening ruthenium complexes: X-ray crystal structure and solution studies of *cis*, *trans*-bis(2-dimethyl-aminoethyl)diphenylphosphino(dichloro)ruthenium(II), **273**, 1

Aminopolycarboxylate ligand complexes
exchange interactions through hydrogen-bond bridges. Crystal structure, spectroscopic characterisation and magnetic properties of the complex [{Cu(en)}$_2$(μ-egta)]·$4H_2O$ (H_4egta = 3,12-bis-(carboxymethyl)-6,9-dioxa-3,12-diazatetradecanedioic acid), **279**, 58

Aminosugar complexes
copper(II) complexes with aminosugar derived Schiff bases as ligands, **273**, 191

Ammine complexes
monofunctional *trans*-$Pt^{II}(NH_3)_2$ modification of pyrimidine-rich deoxyoligonucleotides: direct platination and use of a protective group, **269**, 135
hydrolysis of nerve agent simulants by synergistic effects of tetraminecobalt(III) and microemulsions, **269**, 287
kinetics and mechanism of the oxidation of ammineruthenium(II) complexes by bromine, **270**, 189
anionopentaaminecobalt(III) complexes with polyamine ligands. 29. Relative substitution lability in structurally related chloropentaaminecobalt(III) complexes, **274**, 42
influence of *cis*- and *trans*-diamminedichloroplatinum(II) binding on the helix-coil transition of DNAs with different GC content, **275–276**, 510
kinetics and mechanism for reduction of *trans*-dichloro-tetracyanoplatinate(IV) by tetraammineplatinum(II) and bis(ethylenediamine)platinum(II), **282**, 55
the effect of second-sphere coordination. 7. Isolation of 18-crown-6 ether adducts of ruthenium–ammine complexes, **282**, 247

Ammonia complexes

ammonia ruthenium complexes for the access to new hydrido, carbonyl and isocyanide ruthenium-alkynyl derivatives, **280**, 118
transformations of molecular nitrogen into aromatic amines under the action of titanium compounds, **280**, 264
Ammonia synthesis
effect of ruthenium precursor on hydrogen-treated active carbon supported ruthenium catalysts for ammonia synthesis, **280**, 183
Amphiphilic ion pairs
artificial ion channels of amphiphilic ion pairs composed of oligoether–ammonium and hydrophobic carboxylate or phosphates, **283**, 167
Amphiphilic ligands
synthesis and coordination properties of amphiphilic 3-oxodithiocarboxylic esters, **269**, 83
Angular overlap model
the effects of d–s mixing in low symmetry transition metal complexes, **268**, 55
Anion binding
anions binding to bilirubin oxidase from *Trachyderma tsunodae* K-2593, **273**, 204
Anion complexation
a large cavity reinforced cryptand for the binding of metal cations and anions, **273**, 326
Anion complexes
highly reduced organometallics. 42. A new method for the syntheses of $[V(CO)_6]^-$ and $[V(PF_3)_6]^-$ involving anthracenide mediated reductions of $VCl_3(THF)_3$, **269**, 58
model reactions of a carbonylation catalyst: phosphite induced migratory CO insertion in $[MeIr(CO)_2I_3]^-$, **270**, 382
syntheses and structures of some Ru_2^{5+} complexes containing weakly coordinating anions as axial ligands, **275–276**, 447
Anion exchange
a catalytic activity of glass beads, silica gels and anion-exchange resins modified with metal-porphines in oxidative reactions of ascorbic acid, **283**, 44
Anionic complexes
the coordination mode of 3-hydroxypicolinic acid in oxorhenium(V) complexes. Crystal and molecular structures of $(n\text{-}Bu_4N)$-$[ReOCl_3(HO\text{–}C_5H_3N\text{–}CO_2)]$ and $[ReOCl(HO\text{–}C_5H_3N\text{–}CO_2)_2]$, **269**, 235
Ansa-titanocene complexes
the influence of the catalyst preparation protocol and silane structure on the rate and enantioselectivity of ansa-titanocene catalysed hydrosilation of prochiral ketones, **270**, 330
Ansa-zirconocene
cyclomagnesation of dienes catalyzed by a chiral ansa-zirconocene, **280**, 189
Anthraquinone complexes
properties and kinetics of dihydroxy- and diaminoanthraquinone ruthenium bipyridyl dimers, **268**, 31
Anticancer complexes
pK_a values of aqua ligands of platinum(II) anticancer complexes: [^{1}H, ^{15}N] and ^{195}Pt NMR studies of *cis*- and *trans*-$[PtCl_2(NH_3)$-(cyclohexylamine)], **273**, 8
Anticancer drug complexes
molybdenum complexes of the anticancer drug 6-mercaptopurine, **279**, 136
Anticancer properties
dinuclear complexes of first transition series metals with 4,4′-dipyrazolylmethane: characterisation, DNA binding and anticancer properties, **271**, 151
Antiferromagnetic exchange
exchange interactions through hydrogen-bond bridges. Crystal structure, spectroscopic characterisation and magnetic properties of the complex $[\{Cu(en)\}_2(\mu\text{-egta})]\cdot 4H_2O$ (H_4egta = 3,12-bis-(carboxymethyl)-6,9-dioxa-3,12-diazatetradecanedioic acid), **279**, 58
weak intermolecular antiferromagnetic exchange in $\{[Ru_2(O_2CCH_3)_4(L)]X\}_n$ polymers (L = 4,4′-dipyridine and 1,4-diazabicyclooctane, $X = PF_6^-$ or BPh_4^-), **279**, 121
Antiferromagnetism
homodinuclear lanthanide complexes: Ln_2L_3 (H_2L = tetradentate Schiff bases). Magnetic properties (solid state) and spectroscopic studies (solution), **268**, 125
general synthesis of 'salicylaldehyde half-unit complexes': structural determination and use as synthon for the synthesis of dimetallic or trimetallic complexes and of 'self-assembling ligand complexes', **274**, 73
Antifungal activity
solvent-induced reaction of rhodium(III) halide with N-heterocyclic ligands. Crystal structure of *trans*-dichlorotetrakis(*N*-methylimidazole)rhodium(III) chloride dihydrate, **278**, 209
Antimony-arene complexes
metal–metal multiply-bonded complexes of technetium. 7. Oxidative decomposition of tetrachlorotetrakis(dimethylphenylphosphino)-ditechnetium(II) by an aminium hexachloroantimonate(V), **271**, 187
Antimony complexes
nitrosamine complexes of $SbCl_5$ and $CuCl_2$, **269**, 7
Antisense technology
monofunctional *trans*-$Pt^{II}(NH_3)_2$ modification of pyrimidine-rich deoxyoligonucleotides: direct platination and use of a protective group, **269**, 135
Antitumour activity
synthesis and antitumor activity of platinum(II) complexes of aminocyclodextrin, **272**, 162
intercalator-linked cisplatin: synthesis and antitumor activity of *cis*-dichloroplatinum(II) complexes connected to acridine and phenylquinolines by one methylene chain, **279**, 51
Antitumour complexes
antitumour copper(II) salicylaldehyde benzoylhydrazone (H_2sb) complexes: physicochemical properties and the single-crystal X-ray structures of $[\{Cu(H_2sb)(CCl_3CO_2)_2\}_2]$ and $[\{Cu(Hsb)(ClO_4)(C_2H_5OH)\}_2]$ and the related salicylaldehyde acetylhydrazone (H_2sa) complex, $[Cu(Hsa)Cl(H_2O)]\cdot H_2O$, **267**, 27
Antitumour compounds
rhodium(III) analogues of antitumour-active ruthenium(III) compounds: the crystal structure of [ImH][*trans*-$RhCl_4(Im)_2$] (Im = imidazole), **273**, 62
synthesis, characterization and crystal structure of *trans*-aquatrichlorobis(5,7-dimethyl[1,2,4]triazolo[1,5-*a*]pyrimidine-N^3)-ruthenium(III) monohydrate, **273**, 259
Apo enzyme
X-ray structures of apo and tungstate derivatives of vanadium chloroperoxidase from the fungus *Curvularia inaequalis*, **273**, 160
Aqua complexes
fast water exchange on a Pt(II) center: variable temperature and pressure ^{17}O NMR study at 14.1 tesla of the ionic arylplatinum species $[Pt\{C_6H_3(CH_2NMe_2)_2\text{-}2,6\}(OH_2)]^+(OSO_2CF_3)$, **269**, 322
Aqua form
synthesis of the first sulfido-bridged octahedral rhenium(III) aqua ion $[Re_6S_8(H_2O)_6]$, **271**, 228
Aqua ligand complexes
pK_a values of aqua ligands of platinum(II) anticancer complexes: [^{1}H, ^{15}N] and ^{195}Pt NMR studies of *cis*- and *trans*-$[PtCl_2(NH_3)$-(cyclohexylamine)], **273**, 8
Aquo complexes
second sphere coordination behavior of aquo and amine ligands bound to a η^6-benzeneruthenium(II) cation, **270**, 87
Arenechromium tricarbonyls
ricochet inter-ring haptotropic rearrangement of σ-methyl-(η^5-indenyl)chromium tricarbonyls. Experimental kinetic and theoretical DFT study, **280**, 328

Arene complexes
high pressure liquid chromatography resolution of racemic mixtures of chiral tricarbonyl arene chromium(0) and cyclopentadienyl-titanium(IV) complexes, **268**, 323
bonding modes of arenes in photogenerated (arene)$Cr(CO)_5$ complexes, **271**, 105
electrophilic reactivity of arene ligands coordinated to ruthenium carbido cluster complexes, **274**, 201
structures, dynamic behaviour in solution and cross-coupling reactions of the α-palladiated (η^6-alkylarene)tricarbonylchromium complexes containing a palladium–chromium bond, **280**, 71
synthesis and characterization of arene, mono- and dihydrido-arene, monohydrido-cyclopentadienyl, and phosphite complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$, **281**, 153
synthesis of manganese tricarbonyl cationic complexes of ferrocenyl substituted arenes via a manganese tricarbonyl cation transfer reaction, **281**, 229
Arene coordination
trimethylphosphine halide complexes of titanium, zirconium and hafnium in oxidation states IV, III and II. Crystal structure of $MI_4(PMe_3)_n$ (M = Zr, n = 3; M = Hf, n = 2.5), $M_2X_6(PMe_3)_4$ (M = Hf, Ti, X = I; M = Zr, X = Br) and (μ-η^6,η^6-C_6H_6)-$Hf_2I_4(PMe_3)_4 \cdot C_6H_6$, **271**, 180
Arene oxidation
Lewis base adducts of halogenorhenium(VII) oxides: ^{17}O NMR spectroscopy, structural aspects and catalysis, **279**, 44
Arenethiolate complexes
structure and properties of tetraphenylporphinate iron(III) complexes with an intramolecular NH···S benzenethiolate or NH···O phenolate hydrogen bond, **283**, 91
Arenolate complexes
structure and properties of tetraphenylporphinate iron(III) complexes with an intramolecular NH···S benzenethiolate or NH···O phenolate hydrogen bond, **283**, 91
Arm-off reactions
arm-off reactions of the tripodal polyphosphane $MeC(CH_2PPh_2)_3$ in octahedral rhodium(III) complexes of heterocyclic thioamides, **267**, 39
Aromatic diamines
structures and properties of ternary copper(II) complexes with 4,5-dihydroxyphenylalanine or 2,4,5-trihydroxyphenylalanine and aromatic diamines, **283**, 260
Aromatic heterocyclic complexes
protonation of the ground states of ruthenium(II) photosensitizers, **279**, 200
Aromatic ring stacking
ternary complexes in solution. Intramolecular stacking interactions in mixed ligand complexes formed by copper(II), 2,2′-bipyridyl or 1,10-phenanthroline and a pyrimidine-nucleoside 5′-diphosphate (CDP^{3-}, UDP^{3-}, $dTDP^{3-}$), **283**, 193
Arsines
application of electrospray mass spectrometry to the characterisation of tertiary arsine ligands, **281**, 101
Artificial holoenzyme
hydrophobic vitamin B_{12}. Part 15. Carbon-skeleton rearrangement reactions mediated by hydrophobic vitamin B_{12} covalently bound to an anionic lipid species in aqueous media, **273**, 299
Artificial ion channels
artificial ion channels of amphiphilic ion pairs composed of oligo-ether–ammonium and hydrophobic carboxylate or phosphates, **283**, 167
Arylamines
transformations of molecular nitrogen into aromatic amines under the action of titanium compounds, **280**, 264
Aryl complexes
fast water exchange on a Pt(II) center: variable temperature and pressure ^{17}O NMR study at 14.1 tesla of the ionic arylplatinum species $[Pt\{C_6H_3(CH_2NMe_2)_2\text{-}2,6\}(OH_2)]^+(OSO_2CF_3)$, **269**, 322
titanium–carbon functionalities on an oxo surface defined by a calix[4]arene moiety and its redox chemistry, **270**, 298
a mechanistic study of thermal C–H reductive elimination from a six-coordinate d^6 iridium complex, **270**, 537
synthesis of rhodium porphyrin aryls via intermolecular arene carbon–hydrogen bond activation, **270**, 551
triphenyltellurium(IV) dichalcogenoimidodiphosphinates containing novel six-membered TeXYP$_2$N chelate rings. X-ray structures of $Ph_3Te[(SPPh_2)(OPPh_2)N]$ and $Ph_3Te[(OPPh_2)_2N]$, **271**, 75
pentamethylphenylcopper(I): a square-planar tetranuclear cluster, **277**, 232
the *trans* influence of F, Cl, Br and I ligands in a series of square-planar Pd(II) complexes: relative affinities of halide anions for the metal centre in *trans*-$[(Ph_3P)_2Pd(Ph)X]^+$, **280**, 87
crystal and molecular structures of some tri-aryltin isothiocyanates (Ar_3SnNCS; Ar = C_6H_5, *p*-$CH_3C_6H_4$, *o*-$CH_3C_6H_4$, *o*-$CH_3OC_6H_4$) and of the adduct hexamethylphosphoramide(*N*-thiocyanato)-tris(*o*-tolyl)tin, **282**, 30
Aryl halides
unexpected bell-shaped effect of the ligand on the rate of the oxidative addition to palladium(0) complexes generated in situ from mixtures of $Pd(dba)_2$ and *para*-substituted triarylphosphines, **273**, 76
Aryloxide complexes
rhodium aryloxide complexes containing terdentate nitrogen ligands, C–O bond formation, hydrogen bonding with phenol and oxidative addition reactions. Molecular structure of an Rh(III)–acetyl complex, **270**, 34
the propensity of alkoxide and aryloxide derivatives of tungsten carbonyls to aggregate in solution. Synthesis and X-ray structures of dinuclear, trinuclear and tetranuclear complexes derived from the $MeOW(CO)_5^-$ anion, **270**, 405
crystal structures of aryloxide complexes of zinc embracing sodium aryl group interactions: $Na[Zn(2,6\text{-diphenylphenoxide})_3(H_2O)]$ and $Na[Zn_2(2,6\text{-diisopropylphenoxide})_4Cl] \cdot 3THF$, **274**, 115
reactions of Ti(O–i-Pr)Cl_3 with 2-aminophenol and the crystal structure of the zwitterionic complex $[Ti(O\text{–}i\text{-}Pr)Cl_3(THF)(^-OC_6H_4NH_3^+)] \cdot THF$, **278**, 232
preparation, structure and properties of triphenylphosphine rhodium aryloxides, **280**, 172
Aryloxy compounds
alkoxymesitylchlorogermanes and aryloxymesitylchlorogermanes, **277**, 98
Ascorbic acid
a catalytic activity of glass beads, silica gels and anion-exchange resins modified with metal-porphines in oxidative reactions of ascorbic acid, **283**, 44
Asparagine glutamine derivatives
palladium(II) complexes of *N*-sulfonyl-asparagine and glutamine. Evidence for metal coordination of the deprotonated amide nitrogen of the side-chain, **273**, 397
Asymmetric hydrogenation
synthesis, characterization and application of Ru(BINAP)(Acac)-(MNAA)(MeOH), a highly effective catalyst for the asymmetric hydrogenation of 2-(6′-methoxynaphth-2′-yl)acrylic acid, **270**, 247
probing the nature of H_2 activation in catalytic asymmetric hydrogenation, **270**, 285
Atom transfer
synthesis of *mer*-trichlorotris(tetrahydrofuran)tungsten(III) by inter-metal chlorine atom transfer, **278**, 103
ATP complexes

a ^{31}P and ^{1}H NMR relaxometric study of the interaction between adenosine triphosphate (ATP) and paramagnetic ions (Gd^{3+} and Mn^{2+}), **273**, 92

Autolysis
reductive changes to polynuclear iron(III) clusters in iron-loaded human spleen tissue, **267**, 7

Autoxidation
structure and reactivity of reconstituted myoglobins: interaction between protein and polar side chain of chemically modified hemin, **275–276**, 159

Axial ligands
the orientation of the iron axial ligands in the low-potential cytochrome c_{549} from *Synechocystis sp.* PCC 6803 studied by NMR, **273**, 196

1-Aza-allyl complexes
synthesis and structures of tin(II) and lead(II) 1-aza-allyls; the $[N(SiMe_3)C(Ph)C(SiMe_3)_2]^-$ ligand, **269**, 181

Aza-crown ether complexes
stability constants of copper(I) and silver(I) complexes with aza-crown ethers and cryptands in dimethylsulfoxide, **279**, 186

Azacyclam complexes
new cyclam-type copper(II) complexes with amide molecular padlock: synthesis, properties and crystal structure, **282**, 142

Azadiboriridines
clusters $N_2B_4R_6$ from the fusion of rings NB_2R_3: mechanism, **269**, 43

Azaferrocene
a simple synthesis of 2,5-dimethylazaferrocene, **278**, 101

Azamacrocycle complexes
synthesis and properties of nickel(III) complexes of hexaazamacrocyclic ligands, **269**, 337
rhenium(VII) and (V) imido and oxo complexes containing 1,4,7-triazacyclononane: synthesis, electrochemistry and crystal structures, **272**, 176
the oxidation of nickel^{II}(1,4,8,11-tetraazacyclotetradecane) by hydroperoxyl radicals: a pulse radiolysis study, **273**, 266
a theoretical study of the methyl ligation in tetraaza macrocycle nickel complexes which model the acetyl-CoA synthase active site, **273**, 379
electrochemical and photophysical properties of two-component coordination compounds containing a metallocyclam and an $Re^{I}(bipy)(CO)_3Cl$ subunit, **275–276**, 117
binding properties and crystal structures of azamacrocycles containing nitrophenol side arms, **275–276**, 168
kinetic studies using Co(III) and Cr(III) complexes of 1,4,7,11-tetraazaundecane (2,2,3-tet) and the crystal structure of *trans*-$[CrCl_2(2,2,3\text{-tetH})(OH_2)]Cl\cdot ClO_4$, **278**, 24
preparation and spectroscopic properties of five 7,16-bis(azophenyl)-tetraaza[14]annulene nickel(II) complexes, **281**, 190

Azamacrocyclic ligand complexes
NMR spectra of dinuclear manganese and iron compounds containing 1,4,7-triazacyclononane and 1,4,7-trimethyl-1,4,7-triazacyclononane, **268**, 43

Azauridine complexes
migration of $[Ru^{II}(hedta)]^-$ among three binding sites on 6-azauridine, **277**, 46

Azide-bridged complexes
synthesis, structure and magnetic properties of one-dimensional azide-bridged manganese(III) uniform chain complex $Mn(salpn)N_3$, **271**, 99

Azide complexes
azide promotion of alkane oxidations catalyzed by metal complexes in solution, **270**, 162
a ternary complex of carbonic anhydrase: X-ray crystallographic structure of the adduct of human carbonic anhydrase II with the activator phenylalanine and the inhibitor azide, **275–276**, 295
crystal structures and magnetic properties of uniform and alternating azido-bridged (2,2′-bipyridyl)copper(II) chains, **282**, 82

Azido-bridged complexes
synthesis, structural characterization and magnetic behaviour of two new $(\mu_{1,3}\text{-}N_3)_2$ dinuclear nickel compounds: study of the magnetic transition at 235 K for the *cis*-$(\mu_{1,3}\text{-}N_3)_2[Ni(dl\text{-}cth)]_2(ClO_4)_2$ compound, **278**, 43
synthesis, crystal structure and magnetic properties of the first single azido-bridged copper(II) chain $[Cu(bpym)(N_3)_2]_n$ (bpym = 2,2′-bipyrimidine), **282**, 252

Azido complexes
transition metal complexes with sulfur ligands. Part CXXVII. Azido, halido and nitrido ruthenium complexes with sulfur-rich coordination spheres, **269**, 63
comparative time-resolved IR and UV spectroscopic study of monophosphine and diphosphine platinum(II) azido complexes, **271**, 160

Azine phosphine complexes
chemistry of the azine phosphine ligand *Z,E*-$PPh_2CH_2C(Bu^t)$=N–N=CMe($C_6H_4NO_2$-4): crystal structure of $[Mo(CO)_4\{PPh_2CH_2C(Bu^t)=N\text{–}N=CMe(C_6H_4NO_2\text{-}4)\}]$, **270**, 312

Azobenzene complexes
molecular structure and reactions of azobenzene complexes with iron–lithium compounds, **280**, 288

Azurin
electron tunneling in Ru-modified His46Asp azurin. Coupling through the Cu ligands, **275–276**, 470

B_{12} model complexes
the Co–CH_3 bond in Schiff base B_{12} models: influence of the *trans* and equatorial ligands as assessed by Fourier transform Raman spectroscopy, **275–276**, 90

Barium complexes
synthesis, structure and characterization of zinc(II), copper(II), zinc(II)barium(II) and copper(II)barium(II) complexes of macrocyclic heteronucleating ligands based on isothiosemicarbazide, **282**, 61

Barium compounds
a new form of decavanadate: hydrothermal synthesis and structural characterization of $Ba_{5.33}(H_3O)_{0.33}[V_{10}O_{30}]$, **277**, 263

Benzamidinato complexes
complexes of the bis(trimethylsilyl)-benzamidinato ligand '*siam*': synthesis and X-ray structures of (*siam*)$_2$M, (*siam*)(*siam*H)MX (M = Ni, Pd), (*siam*)$_2$MnI and (*siam*)ReO_3, and their reactivity towards CO_2, **281**, 181

Benzamidine complexes
synthesis and spectroscopic characterization of copper, nickel and cobalt complexes with *N*-(*N*″,*N*″-disubstituted-aminothiocarbonyl)-*N*′-substituted benzamidines, **268**, 13

Benzene complexes
second sphere coordination behavior of aquo and amine ligands bound to a η^6-benzeneruthenium(II) cation, **270**, 87

Benzimidazole-type ligands
metal ion-coordinating properties of imidazole and derivatives in aqueous solution: interrelation between complex stability and ligand basicity, **280**, 50

Benzochlorins
photophysical properties of porphyrins, phthalocyanines, and benzochlorins, **279**, 226

Benzodiazepine complexes
technetium and rhenium carbonyl complexes with 1,4-benzodiazepine derivatives: X-ray crystal structure, **272**, 267

Benzolamide complexes
copper complexes modelling the interaction between benzolamide and Cu-substituted carbonic anhydrase. Crystal structure of Cu(bz)-$(NH_3)_4$ complex, **273**, 334

Benzoporphyrin complexes
preparation of soluble tetrabenzoporphyrins with substituents at the peripheral positions, **277**, 151

Benzothiazole complexes
cytotoxic anionic tribromo platinum(II) complexes containing benzothiazole and benzoxazole donors: synthesis, characterization, and structure–activity correlation, **271**, 137
Benzoxazole complexes
cytotoxic anionic tribromo platinum(II) complexes containing benzothiazole and benzoxazole donors: synthesis, characterization, and structure–activity correlation, **271**, 137
Benzyloxide complexes
elimination of benzene from a benzyloxy ligand to form a carbonyl group on a tungsten-phosphine center, **270**, 202
Bidentate amine complexes
effect of hydrogen bonding on coordination in *trans*-di(*Z*-aminobenzoato-*O*)di(1,3-diaminopropane-*N*,*N'*)copper(II) ($Z=2$, 3 or 4) complex units, **268**, 21
Bidentate diamine complexes
synthesis of binuclear platinum(II) organometallic complexes as potential cross-linking reagents, **275–276**, 141
Bidentate ligand complexes
synthesis and crystal structure of a seven-coordinate dinuclear iron(III) complex containing only bidentate ligands, **274**, 224
synthesis and characterization of mono-, bi- and tetra-ruthenium(II)-phenanthroline complexes with dipyrido(2,3-*a*:2′,3′-*h*)phenazine (dpop), **281**, 1
preparation and properties of a series of (μ-oxo)di(μ-acetato) diruthenium(III) complexes, $[Ru_2(\mu\text{-}O)(\mu\text{-}CH_3COO)_2(pyridine)_4(L)_2]^{2+}$ and $[Ru_2(\mu\text{-}O)(\mu\text{-}CH_3COO)_2(AA)_2(L)_2]^{2+}$, where AA is 2,2′-bipyridine or 1,10-phenanthroline, and L is a monodentate ligand, **283**, 151
Bidentate nitrogen ligand complexes
synthesis and crystal structure of two ternary dicopper(I) complexes having the unsymmetrical coordination arrangement bridged by 1,8-naphthyridine (napy). $[Cu_2(napy)_2(Me_2CO)](PF_6)_2 \cdot 2Me_2CO$ and $[Cu_2(napy)_2(dppm)(CH_3CN)](PF_6)_2$, **271**, 129
Bidentate razoxane ligands
the preparation and crystal and molecular structure of the complex *cis*-$PtCl_2$(razoxane), **268**, 313
Bidentate sulfur ligand complexes
synthesis and crystal structure of a dinuclear platinum(II) bis(*N*,*N*-diethylamino)carbeniumdithiocarboxylate complex $Pt_2\{\mu\text{-}(Et_2N)_2C_2S_2\}_2Cl_4 \cdot 0.5CH_2Cl_2$: structural study of the coordinated inner salt, **283**, 256
Bimetallic complexes
cooperative bimetallic reactivity, **270**, 13
preparation and properties of ferrocenyl bimetallic compounds for non-linear optics, **279**, 243
bimetallic cyclooligosiloxanolate complexes of copper and nickel, **280**, 282
Bimolecular nucleophilic attack
nucleophilic attack within Ge, Sn and Pb complexes containing $Me_2N(CH_2)_3$– as a potential intramolecular donor ligand, **275–276**, 203
BINAP complexes
synthesis, characterization and application of Ru(BINAP)(Acac)-(MNAA)(MeOH), a highly effective catalyst for the asymmetric hydrogenation of 2-(6′-methoxynaphth-2′-yl)acrylic acid, **270**, 247
Binuclear complexes
crystal structure, spectroscopic, magnetochemical, thermoanalytical and electrochemical properties of binuclear copper(II) complexes of Suprofen, **268**, 239
mononuclear and binuclear *o*-xylylene derivatives of Pd(II), **269**, 191
the cyanide ligand as an efficient bridge in mixed-valence complexes, **269**, 246
X-ray diffraction and EPR studies of binuclear complex $[CuCl(C_6H_9N_3O_3)_2 \cdot H_2O]_2Cl_2$ crystals, **269**, 326
substitution of inosine for chloride in $[Pt_2(hdta)Cl_2]^{2-}$ ($hdta^{4-}$ = 1,6-hexanediamine-*N*,*N*,*N'*,*N'*-tetraacetate), **271**, 124
oxidative addition of methyl iodide and iodine to new binuclear rhodium(I) and iridium(I) compounds containing diaminoanthraquinonate-bridging ligands. Crystal structure of $[Rh_2(\mu\text{-}1,4\text{-}DA)(CO)_2(PPh_3)_2]$ (1,4-H_2DA = 1,4-diaminoanthraquinone), **274**, 15
crystal structure, EPR and magnetic susceptibility studies of tetrakis[μ-(β-alanine)-*O*,*O'*]dichlorodicopper(II) dichloromonohydrate, **275–276**, 28
complexes of heterocyclic thiones and group twelve metals. Part 1. Preparation and characterisation of 1:1 complexes of mercury(II) halides with 1-methylimidazoline-2(3H)-thione: the crystal structure of [(μ$_2$-dibromo) bis(*trans*{(bromo)(1-methyl-imidazoline-2(3H)-thione)}mercury(II))] at 160 K, **277**, 14
crystal structure and characterization of binuclear copper(II) complex with *N*-(2-hydroxyphenyl)-3-methoxycarbonyl-4,6-dimethylsalicylamide, **277**, 134
synthesis and characterisation of mononuclear and binuclear iron(II) complexes of pentadentate and bis(pentadentate) ligands derived from 1,4,7-triazacyclononane, **279**, 192
spectroscopic characterization of coaxially and π–π stacked binuclear copper(II) complexes of ω,ω′-bis(pyridine-2-carboxamido)-alkanes in acidic solution, **283**, 61
Binucleating ligand complexes
cooperative bimetallic reactivity, **270**, 13
Bioactivity
Cu^{II}–herbicide complexes: structure and bioactivity, **272**, 24
Bioinorganic chemistry
Mössbauer spectroscopy with synchrotron radiation: a new technique entering biological inorganic chemistry, **275–276**, 334
Biosensors
tris(4,4′-dimethoxy-2,2′-bipyridine)osmium(II), amperometric properties and crystal structure, **274**, 64
Biphosphinemonoxide complexes
synthesis and characterization of homologous nickel(II) and palladium(II) complexes with biphosphine monoxide ligands $Ph_2P(CH_2)_nP(O)Ph_2$ (n = 1–3) and $pTol_2P(CH_2)P(O)pTol_2$, **280**, 150
Bipyridine complexes
the stereochemistry of the oxidation of diamines and of the reduction of diimines in the coordination sphere of ruthenium(II), **272**, 153
tris(4,4′-dimethoxy-2,2′-bipyridine)osmium(II), amperometric properties and crystal structure, **274**, 64
ground and electronically excited states of $Cr(CO)_4$(bipyridine): energy factored force field analysis of CO stretching vibrations and resonance Raman study, **278**, 83
the chemistry of dinitrato-2,2′-bipyridinecopper(II): preparation and characterization of binuclear complexes having a $Cu(\mu\text{-}OH)_2Cu$ and $Cu(\mu\text{-}N_3)_2Cu$ core, **278**, 143
syntheses and X-ray crystal structures of *cis*-$[Mn(bipy)_2Cl_2] \cdot 2H_2O \cdot EtOH$ and *cis*-$[Mn(phen)_2Cl_2]$ (bipy = 2,2′-bipyridine; phen = 1,10-phenanthroline); catalysts for the disproportionation of hydrogen peroxide, **279**, 24
linkage isomers of alizarin-bis(bipyridine)ruthenium(II), **281**, 126
Bipyridyl complexes
properties and kinetics of dihydroxy- and diaminoanthraquinone ruthenium bipyridyl dimers, **268**, 31
crystal structures and magnetic properties of uniform and alternating azido-bridged (2,2′-bipyridyl)copper(II) chains, **282**, 82
Bipyrimidine complexes
mixed ligand copper(II) coordination polymers constructed by Cu–bpm–Cu dimer unit (bpm = 2,2′-bipyrimidine) as a building block. Crystal structures and magnetic properties of $\{[Cu(bpm)(SO_4)](H_2O)\}_n$, $[Cu_2(bpm)(suc)_{0.5}(ClO_4)_2(OH)(H_2O)_2]_n$ and $\{[Cu(bpm)_{1.5}(suc)_{0.5}](ClO_4)(H_2O)_2\}_n$ (suc = succinate), **283**, 80
Bismuth catalysts

selective oxidation of glucose on bismuth-promoted Pd–Pt/C catalysts prepared from $NOct_4Cl$-stabilized Pd–Pt colloids, **270**, 95

Bismuth complexes
synthesis, X-ray structure and bonding of tris(2,2-6,6-tetramethylheptane-3,5-dionato) bismuth(III), **275–276**, 340
mixed valence, tetradisordered structure of iodine-doped bismuth(III) diphthalocyanine, **281**, 195

Block copolymer complexes
interaction of metal compounds with 'double-hydrophilic' block copolymers in aqueous medium and metal colloid formation, **280**, 348

Blue copper proteins
probing the metal site in *Rhus vernicifera* stellacyanin by Ni(II) substitution and paramagnetic NMR spectroscopy, **273**, 367

Bond enthalpies
extracting absolute titanium–alkyl and –hydride bond enthalpies from relative $D(TiR(H))$ in $(silox)_2(^tBu_3SiNH)TiR$: electronegativity and *ECT* models, **270**, 414

Bonding complexes
synthesis, X-ray structure and bonding of tris(2,2-6,6-tetramethylheptane-3,5-dionato) bismuth(III), **275–276**, 340

Borate complexes
a bimetallic main group oxide cluster of the oxovanadium borate system: $(H_3NCH_2CH_2NH_3)_4[(VO)_{12}O_4\{B_8O_{17}(OH)_4\}_2\{Mn(H_2O)_2\}_2]\cdot H_2O$, **282**, 123

Borate ion
^1H and ^{11}B NMR evidence for specific binding of borate ion to cytochrome *c*, **273**, 388

Borohydride adduct
the 1,8-bis(2′-pyridyl)-3,6-dithiaoctane complex of nickel(II): X-ray crystal structure and borohydride adduct formation, **278**, 217

Borohydride complexes
calculation of NMR paramagnetic shift temperature dependence in actinide methyl borohydrides, **272**, 43

Borylated cobaloxime
synthesis and structure of borylated organocobaloximes containing neutral nitrile ligands, **272**, 74

Bromide complexes
$(C_5H_7N_2O)Cu_3Br_4$: template synthesis of a two-dimensional copper(I) bromide lattice, **279**, 1

Bromination
electrophilic substitution of metal-coordinated pyrazole: nitration, sulfonation and bromination of $[Co(NH_3)_5(pyzH)]^{3+}$ (pyzH = pyrazole), **277**, 89

C_{70} complexes
exohedral and endohedral complexation energies of complexes $X@C_{70}$ between higher fullerene and rare-gas atoms (X = He, Ne, Ar, Kr, Xe), **272**, 33

C_{60}-fullerinide complexes
C_{60}-fullerenides of ytterbium and lutetium, **280**, 138

Cadmium complexes
synthesis and crystal structures of complexes containing an Fe→Cd donor–acceptor bond formed by reaction of *trans*-$(CO)_3Fe(Ph_2Ppym)_2$ (Ph_2Ppym = 2-(diphenylphospino)pyrimidine) with CdX_2 (X = Cl, Br, I, SCN, ClO_4), **268**, 177
synthesis and crystal structure of $[Cd\{(c\text{-}C_6H_{11})_2P(Se)C(S)NPh\}_2]$, **269**, 317
dinuclear metal complexes of Cd(II), Zn(II) and Fe(II) with triple-helical structure and predetermined chirality, **271**, 36
some novel carbamato chemistry of cadmium and zinc, **274**, 239
synthesis and structural studies on metal nitrate complexes with 1,3-dimethyllumazine and 1,3,6,7-tetramethyllumazine: crystal structure of two new cobalt(II) and copper(II) three-dimensionally hydrogen-bonded complexes and a cadmium(II) complex with unusual geometry, **277**, 103
structure and magnetic properties of a novel chloro-bridged polymeric cadmium(II) complex with pyridyl-substituted nitronyl nitroxide, **279**, 233

Calcium complexes
electron spectroscopy for chemical analysis investigation of the interaction of uranyl and calcium ions with humic acids, **273**, 234

Calixarene complexes
electronic absorption spectroscopy for the investigation of the solution binding of gold, palladium, mercury and silver salts to the lower rim substituted calix[4]arenes 25,26,27,28-(2-*N*,*N*-dimethyldithiocarbamoylethoxy)calix[4]arene and 25,26,27,28-(2-methylthioethoxy)calix[4]arene, **270**, 216

Calix[4]arene complexes
titanium–carbon functionalities on an oxo surface defined by a calix[4]arene moiety and its redox chemistry, **270**, 298

Calix[6]arene complexes
cation-π interactions between potassium ions and aromatic rings. Crystal structures of three potassium complexes of calix[6]arene, **281**, 36

Carbamate complexes
palladium-catalyzed seven-membered N,O-heterocycle ring formation, **270**, 123

Carbamato complexes
eight-coordinate chelate complexes of zirconium(IV) and niobium(IV): X-ray diffractometric and EPR investigations, **270**, 177
some novel carbamato chemistry of cadmium and zinc, **274**, 239

Carbene complexes
concomitant coordination of vinylidene and carbene ligands made possible at ruthenium(II), **272**, 1
study of the relative π-acceptor ability of unsaturated ligands $:C(=C)_nPh_2$ (n = 0, 1, 2 and 4) based on cyclic voltammetry of complexes *trans*-$[RhCl(L)(P^iPr_3)_2]$ (L = $:C(=C)_nPh_2$, C_2H_4 and CO), **275–276**, 435
reactions of platinum(II)–alkyne and –alkynyl complexes with the hydroxy compounds $HO(CH_2)_2X$ (X = Br, I, OH), **275–276**, 515

Carbide complexes
chiroptical properties of heterotrimetallic 'wing-tip' butterflies; synthesis and crystal structure of $(\mu\text{-}H)Ru_2Fe_2PdC(CO)_{12}(\eta^3\text{-}\beta\text{-}C_{10}H_{15})$, **280**, 275

Carbido complexes
electrophilic reactivity of arene ligands coordinated to ruthenium carbido cluster complexes, **274**, 201
the use of photogenerated intermediates in the study of cluster build-up reactions: the generation of $Ru_6C(CO)_{17}$ from $Ru_3(CO)_{12}$, **278**, 250

Carbon dioxide
mechanistic details for metathetical exchange between XCO (X = O and RN) and the tin(II) dimer, $\{Sn[N(SiMe_3)_2](\mu\text{-}OBu^t)\}_2$, **270**, 118

Carbon dioxide activation
the reaction of $Ru(CO)_2(MeCO_2)_2(PR_3)_2$ or $Ru_2(CO)_4(MeCO_2)_2(PR_3)_2$ [R = Bu, Ph] with H_2 in the presence of Na_2CO_3: a stoichiometric carbon dioxide activation, **272**, 141

Carbon dioxide complexes
infrared spectroelectrochemical and electrochemical kinetics studies of the reaction of nickel cluster radicals $[Ni_3(\mu_2\text{-}dppm)_3(\mu_3\text{-}L)(\mu_3\text{-}I)]^+$(L = CNR, R = CH_3, i-C_3H_7, C_6H_{11}, $CH_2C_6H_5$, t-C_4H_9, 2,6-$Me_2C_6H_3$; L = CO) with carbon dioxide, **270**, 111

Carbon dioxide fixation
fixation of carbon dioxide with 1,2-epoxypropane catalysed by tetra-t-butylphthalocyaninatoaluminium(III) acetate, tetra-t-butylphthalocyaninatoaluminium(III) trimethylsilanolate and tetra-t-butylphthalocyaninatoaluminium(III) triphenylsilanolate, **278**, 223

Carbon dioxide reduction
bioinorganic chemistry of nickel and carbon dioxide: an Ni complex behaving as a model system for carbon monoxide dehydrogenase enzyme, **272**, 38

Carbon disulphide disproportionation
nucleophilic attack at the five-coordinate anion $[Mn(CO)_3(3,5\text{-di-tert-butyl-catecholate})]^-$ controlled by electronic and steric effects, **268**, 1
Carbon–hydrogen activation
intramolecular and intermolecular C–H activation at a cationic Pt^{II} center, **270**, 467
Carbon monoxide binding
carbonyl bonding on ZnO(0001), CuCl(111) and $Cu_2O(111)$ $3d^{10}$ metal ion surfaces: photoelectron spectroscopy of $CO/Cu_2O(111)$ and electronic structure calculations, **275–276**, 327
Carbon monoxide complexes
molecular structure of the methoxy-iron(III) derivative of 5,15-(*o,o'*(2-methyl-2'-hydroxy-3,3'-diamidobiphenyl)-diphenylporphyrin and CO binding properties of iron(II)–pyridine complexes of 2,2'-substituted biphenyl strapped porphyrins, **283**, 98
Carbon monoxide dehydrogenase
bioinorganic chemistry of nickel and carbon dioxide: an Ni complex behaving as a model system for carbon monoxide dehydrogenase enzyme, **272**, 38
Carbonylation
the dicarbonylation of diiodomethane to malonate esters catalysed by triethylphosphine complexes of rhodium, **280**, 99
Carbonylation catalyst
model reactions of a carbonylation catalyst: phosphite induced migratory CO insertion in $[MeIr(CO)_2I_3]^-$, **270**, 382
Carbonyl complexes
the kinetics and mechanism of ligand exchange in photogenerated $(\eta^1\text{-bromobenzene})Cr(CO)_5$/alkene systems, **267**, 49
site-selective PMe_3 addition to the donor–acceptor complex $Ru_2(CO)_6[(Z)\text{-}Ph_2PCH{=}CHPPh_2]$. X-ray diffraction structure of $Ru_2(CO)_5(PMe_3)[(Z)\text{-}Ph_2PCH{=}CHPPh_2]$, **267**, 223
dissociative kinetics of 'lightly bonded ligands' *L* from the clusters $Rh_6(CO)_{15}L$ (*L* = DMSO, NCMe, cyclooctene, THF and EtOH), **268**, 189
highly reduced organometallics. 42. A new method for the syntheses of $[V(CO)_6]^-$ and $[V(PF_3)_6]^-$ involving anthracenide mediated reductions of $VCl_3(THF)_3$, **269**, 58
ligand connected metal clusters. The molecular structures and solid state packing of $\{Ru_3(CO)_{11}\}_2$(bis(diphenylphosphino)ethane) and $\{Ru_3(CO)_{11}\}_2$(1,4-bis(diphenylphosphino)benzene), **270**, 28
η^3-allyl–cobalt(II) complexes, **270**, 60
model reactions of a carbonylation catalyst: phosphite induced migratory CO insertion in $[MeIr(CO)_2I_3]^-$, **270**, 382
the propensity of alkoxide and aryloxide derivatives of tungsten carbonyls to aggregate in solution. Synthesis and X-ray structures of dinuclear, trinuclear and tetranuclear complexes derived from the $MeOW(CO)_5^-$ anion, **270**, 405
monovalent copper as a potential catalyst for formation of acetaldehyde via the migration of methyl radicals to the coordinated carbonyl in the complex $(CO)Cu^{II}\text{–}CH_3^+$, **270**, 440
first- and second-order mechanisms for oxidative addition for bound methyl disulfide in the complex $W(CO)_3(phen)(MeSSMe)$, **270**, 488
bonding modes of arenes in photogenerated $(arene)Cr(CO)_5$ complexes, **271**, 105
synthesis, characterization and reactivity of bis(diphenylphosphino)-amine nickel complexes, **271**, 112
the reaction of $Ru(CO)_2(MeCO_2)_2(PR_3)_2$ or $Ru_2(CO)_4$-$(MeCO_2)_2(PR_3)_2$ [R = Bu, Ph] with H_2 in the presence of Na_2CO_3: a stoichiometric carbon dioxide activation, **272**, 141
steric limitations in associative substitution reactions of $Os_3(CO)_9(\mu\text{-}C_4Ph_4)$, **274**, 82
NMR characterization of four new isomers of $H(\mu\text{-}H)Os_3(CO)_{10}$-(phosphine), **275–276**, 521
Fourier transform IR study of the interaction of $[(\eta^5\text{-}C_5H_5)_2$-$FeRu(CO)_4]$ with hydroxylated alumina surfaces, **277**, 61
ground and electronically excited states of $Cr(CO)_4$(bipyridine): energy factored force field analysis of CO stretching vibrations and resonance Raman study, **278**, 83
bonding of halogenated arenes in photogenerated $(arene)M(CO)_5$ complexes (M = Cr, Mo, W), **278**, 202
the use of photogenerated intermediates in the study of cluster build-up reactions: the generation of $Ru_6C(CO)_{17}$ from $Ru_3(CO)_{12}$, **278**, 250
the synthesis, molecular structure and fluxional properties of the 1-phenylpropyne complex $[WI_2(CO)\{Ph_2P(CH_2)PPh_2\}(\eta^2\text{-}MeC_2Ph)]$, **279**, 65
some chlorocarbonylruthenium(II) complexes of P,N-donor ligands: crystal structures of $[RuCl(CO)\{PPh_2CH_2C(Bu^t){=}NNH_2\}_2]Cl$ and *fac,cis*-$[RuCl_2(CO)\{PPh_2CH_2C(Bu^t){=}N\text{–}N{=}C(Bu^t)\text{-}CH_2PPh_2\}]$, **279**, 95
binuclear and tetranuclear complexes prepared by reactions of dodecacarbonyltriruthenium with the N-substituted cinnamic acid amides. Crystal and molecular structures of $[Ru_2(\mu\text{-}H)(\mu\text{-}\eta^3\text{-}PhC{=}CHC(O)N(CH_2)_4)(CO)_6]$ and $[Ru_4(\mu_3\text{-}H)_2(\mu\text{-}\eta^3\text{-}PhC{=}CHC(O)N(CH_2)_4)_2(CO)_{10}]$, **280**, 243
ferrocenyl-substituted imine ligands and their reactivity towards $Fe_2(CO)_9$; crystal structure of $Fc\text{–}C(H){=}C(H)\text{–}C(H){=}N\text{–}C_6H_{11}$ and $(\eta^4\text{-}R^1\text{–}C(H){=}C(H)\text{–}C(H){=}N\text{–}R^2)Fe(CO)_3$ ($R^1 = C_6H_5$, $R^2 = Fc$; $R^1 = Fc$, $R^2 = C_6H_{11}$, $R^1 = R^2 = Fc$; $Fc = (\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4)$), **282**, 111
Carborane complexes
synthesis, solution behavior and molecular structure of 16-electron *closo*-2-(Ph_3P)-1-N,2-[μ-($\eta^2\text{-}CH_2CH{=}CH_2$)]-1-N-($\sigma\text{-}CH_2CH{=}CH_2$)-2,1-$RhCB_{10}H_{10}$, the first η^2-olefinic monocarbon metallacarborane, **280**, 233
Carboxylate-bridged complexes
syntheses and structures of some Ru_2^{5+} complexes containing weakly coordinating anions as axial ligands, **275–276**, 447
synthesis, characterization and kinetics of the metallation of the intermediate compound $[Rh_2(\mu\text{-}O_2CCH_3)_3(O_2CCH_3)\{\eta^2\text{-}\underline{P}(2\text{-}(CH_3\underline{O})C_6H_4)Ph_2\}]\cdot H_2O$ to produce compound $[Rh_2(\mu\text{-}O_2CCH_3)_3\text{-}\{\mu\text{-}(\underline{C}\underline{C}_5H_4)\underline{P}(2\text{-}(CH_3O)C_6H_4)Ph\}]\cdot(H_2O)$ from Rh(II) tetraacetate, **278**, 61
Carboxylate complexes
three dinuclear copper(II) carboxylates with the paddle-wheel cage structure as intermediates in copper(II) catalyzed oxidations of carboxylic acids. X-ray crystal structures of [tetrakis(diphenyl acetato-μ-*O,O'*)bis(acetonitrile-*N*)dicopper(II)] tetrakis-(acetonitrile), [tetrakis(diphenylacetato-μ-*O,O'*)bis-(acetone-*O*)dicopper(II)] and [tetrakis(1-phenyl-1-carboxylato-μ-*O,O'-cyclo*-pentane)-bis(ethanol-*O*)dicopper(II)] bis(ethanol), **267**, 183
effect of substitution on packing in *trans*-di(*Z*-chlorobenzoato-*O*)-di(1,3-diaminopropane-*N,N'*)nickel(II) (*Z* = 2, 3 or 4) complexes, **267**, 249
synthesis, spectral and structural characterization of the zinc carboxylate [Zn(2-quinolinecarboxylato)$_2$(1-methylimidazole)$_2$], **269**, 283
synthesis, characterization, and reactivity of organometallic Zr(IV) carboxylate complexes, **270**, 527
the reaction of $Ru(CO)_2(MeCO_2)_2(PR_3)_2$ or $Ru_2(CO)_4$-$(MeCO_2)_2(PR_3)_2$ [R = Bu, Ph] with H_2 in the presence of Na_2CO_3: a stoichiometric carbon dioxide activation, **272**, 141
polymeric and dimeric magnetic properties of square planar Cu(II) species controlled by hydrogen bond networks: $[Cu^{II}\{OCO\text{-}2,6\text{-}(CH_3CONH)_2C_6H_3\}_2(H_2O)_2]\cdot nH_2O$ ($n = 1, 4$), **275–276**, 43
liquid secondary ion mass spectrometric study of diruthenium(II,III) complexes, **277**, 76
synthesis and X-ray crystal determination of four novel lanthanide(III) oxydiacetate complexes, **281**, 18
Carboxylate ionization

the effect of surface charge on the reduction potential and heme propionate ionization of a monoheme cytochrome: spectroscopic and potentiometric studies of trifluoroacetylated cytochrome c_{551} from *Pseudomonas aeruginosa*, **275–276**, 256

Carboxylato-bridged complexes
structural and spectroscopic models of the manganese catalase active site. Isolation and structures of the asymmetric $[(H_2O)Mn^{III}(\mu\text{-}O)(\mu\text{-}O_2CR)_2Mn^{III}(L)]$ ($L = Cr_2O_7^{2-}$, CH_3OH) cores: analogs of a substrate-bound catalase active site intermediate, **281**, 25

Carboxylato complexes
synthesis of tetracarboxylatediruthenium(II) compounds: reduction and disproportionation reactions of diruthenium(II,III) units. Crystal structure of $Ru_2(\mu\text{-}O_2CCOPh)_4(thf)_2$, **279**, 30

Catalysis
three dinuclear copper(II) carboxylates with the paddle-wheel cage structure as intermediates in copper(II) catalyzed oxidations of carboxylic acids. X-ray crystal structures of [tetrakis(diphenyl acetato-μ-*O*,*O*′)bis(acetonitrile-*N*)dicopper(II)] tetrakis-(acetonitrile), [tetrakis(diphenylacetato-μ-*O*,*O*′)bis(acetone-*O*)dicopper(II)] and [tetrakis(1-phenyl-1-carboxylato-μ-*O*,*O*′-*cyclo*-pentane)-bis(ethanol-*O*)dicopper(II)] bis(ethanol), **267**, 183
zirconocene–alumoxane (1:1) – a catalyst for the selective dimerization of α-olefins, **270**, 20
selective C–H bond activation of arenes catalyzed by methylrhenium trioxide, **270**, 55
azide promotion of alkane oxidations catalyzed by metal complexes in solution, **270**, 162
catalytic reduction of acetylene in the presence of molybdenum and iron clusters, including FeMo cofactor of nitrogenase, **270**, 221
conductive polymeric films of *meso*-tetrakis[2,2]-*para*-cyclophanylmetalloporphyrins — spectroelectrochemistry and catalytic properties, **271**, 167
preparation and characterization of the catalytically active heteropolyoxotungstate anion $[Co^{3+}ZnW_{11}O_{40}H_2]^{7-}$, **278**, 91
the dicarbonylation of diiodomethane to malonate esters catalysed by triethylphosphine complexes of rhodium, **280**, 99
effect of ruthenium precursor on hydrogen-treated active carbon supported ruthenium catalysts for ammonia synthesis, **280**, 183
cyclomagnesation of dienes catalyzed by a chiral ansa-zirconocene, **280**, 189

Catalytic activity
synthesis and catalytic activity of manganese(II) complexes of pentanedioic acid; X-ray crystal structure of $[Mn(phen)_2(H_2O)_2]\cdot[Mn(O_2C(CH_2)_3CO_2)(phen)_2H_2O](O_2C(CH_2)_3CO_2)\cdot 12H_2O$ (phen = 1,10-phenanthroline), **277**, 257

Catalytic dimerization
synthesis and reactivity of Ru(II) complexes containing the phosphino-amine $Ph_2PCH_2CH_2NMe_2$, **268**, 69

Catalytic disproportionation
syntheses and X-ray crystal structures of *cis*-$[Mn(bipy)_2Cl_2]\cdot 2H_2O\cdot EtOH$ and *cis*-$[Mn(phen)_2Cl_2]$ (bipy = 2,2′-bipyridine; phen = 1,10-phenanthroline); catalysts for the disproportionation of hydrogen peroxide, **279**, 24

Catalytic hydrogenation
synthesis, immobilization and catalytic activity of some silylated cyclopentadienyl rhodium(I) complexes, **280**, 21

Catalytic insertion
thermal and photochemical substitution reactions of $CpRe(PPh_3)_2H_2$ and $CpRe(PPh_3)H_4$. Catalytic insertion of ethylene into the C–H bond of benzene, **270**, 77

Catalytic oxidations
synthesis, characterization and stereoselective catalytic oxidations of chelated deuterohaemin-glycyl-L-histidine complexes, **273**, 339
oxidations in the H_2O_2/V^V/AcOH system. Evidence of vanadium(V) complexes with superoxide anion and singlet dioxygen molecule as ligands, **275–276**, 453
synthetic models for biological trinuclear copper clusters. Trinuclear and binuclear complexes derived from an octadentate tetraaminetetrabenzimidazole ligand, **282**, 180

Catalytic oxygenation
catalytic oxygenation of olefin with dioxygen and tetra-t-butylphthalocyanine complexes in the presence of sodium borohydride, **283**, 275

Catecholate complexes
nucleophilic attack at the five-coordinate anion $[Mn(CO)_3$(3,5-di-tert-butyl-catecholate)$]^-$ controlled by electronic and steric effects, **268**, 1
X-ray crystallographic and absorption spectroscopic analyses of structures of catecholato(pyridine)iron chloride complexes in relevance to functional model complexes for catechol 1,2-dioxygenases, **275–276**, 222
preparation and electrochemistry of six-coordinate monooxo molybdenum(VI) complexes containing bidentate catecholate and tridentate NOS-donor Schiff base ligands, **278**, 1

Catechol dioxygenase
X-ray crystallographic and absorption spectroscopic analyses of structures of catecholato(pyridine)iron chloride complexes in relevance to functional model complexes for catechol 1,2-dioxygenases, **275–276**, 222

Cationic clusters
reactivity studies on cationic non-hydridic triruthenium carbonyl clusters. Reactions of $[Ru_3(\mu_3\text{-ampy})(CO)_{10}][BF_4]$ with hydrogen, triethylsilane and triphenylstannane, **268**, 131

Cationic complexes
intramolecular and intermolecular C–H activation at a cationic Pt^{II} center, **270**, 467

CF_3-substituted phosphine complexes
the synthesis of the tripodal phosphine $CH_3C\{CH_2P(m\text{-}CF_3C_6H_4)_2\}_3$ and its rhodium coordination chemistry, **270**, 499

Chalcogenide complexes
generation of mixed iron chalcogenide clusters $MFeX_n^+$ (M = V, Fe; X = O, S; n = 2, 3) in the gas phase, **273**, 111
variable coordination modes for 1,1′-bis(diphenylphosphino)-ferrocene (dppf) mono- and di-chalcogenides or -oxides (dppfE and $dppfE_2$; E = O, S or Se) in copper(I) complexes. Crystal structure of $[\{Cu(dppf)\}_2(\mu\text{-}dppfS_2)](BF_4)_2$ and $[Cu(dppfSe)_2]BF_4$, **277**, 163
synthesis, cyclic voltammetry and $^{31}P\{H\}$ NMR spectra of $[M_3E_2L_3]^{2+}$ clusters; M = Ni, Pd or Pt; E = S, Se or Te; L = bis(diphenylphosphino)methane, 1,2-bis(diphenylphosphino)-ethane, 1,3-bis(diphenylphosphino)propane, or 1,4-bis-(diphenylphosphino)butane, **281**, 174

Chalcogenoimidodiphosphinate complexes
triphenyltellurium(IV) dichalcogenoimidodiphosphinates containing novel six-membered $TeXYP_2N$ chelate rings. X-ray structures of $Ph_3Te[(SPPh_2)(OPPh_2)N]$ and $Ph_3Te[(OPPh_2)_2N]$, **271**, 75

Chelate complexes
evaluation of solution structures of highly luminescent europium(III) chelates by using laser induced excitation of the $^7F_0 \rightarrow {}^5D_0$ transition, **267**, 63
synthesis and crystal structure of the *trans*-dichloro(*N*,*N*′-bis-[2(*S*)-pyrrolidylmethyl]-ethane-1,2-diamine)cobalt(III) tetrachlorocobaltate(II) complex, **267**, 127
synthesis and coordination properties of amphiphilic 3-oxodithiocarboxylic esters, **269**, 83
synthesis, structure, and properties of {*N*,*N*′-bis[2-(1-pyrazolyl)-ethyl]pyridine-2,6-dicarboxamido}copper(II), **269**, 269
axial–equatorial interactions in chelates of β-diones. Crystal and molecular structure of bis(3-nitro-2,4-pentanedionato)copper(II), **269**, 332
coordination of a disubstituted alkene in a chelated d^0-metal–alkyl–alkene complex. Evidence for equilibrium between complexes with coordinated and free alkenes, **270**, 5

syntheses and properties of bivalent metal complexes with *N*-(2-hydroxyethyl)ethylenediamine-*N*,*N*′,*N*′-triacetic acid and the crystal structure of *N*-(2-hydroxyethyl)ethylenediamine-*N*,*N*′,*N*′-triacetato-copper(II) monohydrate, **271**, 13
the effect of end-group substitutions and use of a mixed solvent system on β-diketones and their iron complexes, **278**, 170
chelated alkyl halide manganese complexes $(\eta^5{:}\eta^1\text{-}C_5H_4CH_2CH_2X)$-$Mn(CO)_2$ from photolysis of $(\eta^5\text{-}C_5H_4CH_2CH_2X)Mn(CO)_3$, **280**, 316

Chelate ligand complexes
zinc thiolate complexes with chelating nitrogen ligands, **282**, 193

Chelate phosphane complexes
the synthesis of substituted bis[(diarylphosphinomethyl)cyclopentadienyl]zirconocene dichloride complexes for the preparation of heterodimetallic complexes containing early/late transition metal combinations, **270**, 446

Chelate ring complexes
chelate ring-opening ruthenium complexes: X-ray crystal structure and solution studies of *cis*,*trans*-bis(2-dimethyl-aminoethyl)diphenylphosphino(dichloro)ruthenium(II), **273**, 1
tetrahedral coordination by a seven-membered chelate ring, **275–276**, 279

Chelating diamine complexes
effect of substitution on packing in *trans*-di(*Z*-chlorobenzoato-*O*)-di(1,3-diaminopropane-*N*,*N*′)nickel(II) (*Z* = 2, 3 or 4) complexes, **267**, 249

Chemical evolution
salt-induced peptide formation from amino acids in the presence of clays and related catalysts, **272**, 89

Chemoimmunoconjugates
synthesis of two maleimide derivatives of *cis*-configurated platinum(II) complexes for the preparation of chemoimmunoconjugates, **267**, 133

Chemotherapy
toward a novel metal based chemotherapy against tropical diseases. 4. Synthesis and characterization of new metal-clotrimazole complexes and evaluation of their activity against *Trypanosoma cruzi*, **275–276**, 528

Chiral amine complexes
binuclear oxo-bridged iron(III) complexes of chiral tetradentate ligands, **274**, 155

Chiral azaphosphole
transition metal complexes. IX. Nickel complexes with a chiral azaphosphole ligand, **270**, 273

Chiral complexes
high pressure liquid chromatography resolution of racemic mixtures of chiral tricarbonyl arene chromium(0) and cyclopentadienyl-titanium(IV) complexes, **268**, 323
additions of carbon nucleophiles to acyclic imine complexes of the chiral rhenium Lewis acid $[(\eta^5\text{-}C_5H_5)Re(NO)(PPh_3)]^+$: controlling factors in 1,3-asymmetric induction and syntheses of non-racemic organic amines, **269**, 167
isomerization and racemization of (acetylacetonato)bis(4,4′-dimethyl-2,2′-bipyridine 1,1′-dioxide)chromium(III), (acetylacetonato)bis(4,4′-dimethoxy-2,2′-bipyridine 1,1′-dioxide)-chromium(III) and (acetylacetonato)bis(4,4′-diethoxy-2,2′-bipyridine 1,1′-dioxide)chromium(III) complexes, **281**, 221

Chiral diphosphine complexes
pentamethylcyclopentadienyl ruthenium(II) complexes containing chiral diphosphines: synthesis, characterisation and electrochemical behaviour. X-ray structure of $(\eta^5\text{-}C_5Me_5)Ru\{(S,S)\text{-}Ph_2PCH(CH_3)CH(CH_3)PPh_2\}Cl$, **282**, 163

Chiral P,N-ligand complexes
palladium(II) and platinum(II) complexes containing new phosphinoxy chiral P,N-ligands, **277**, 119

Chiral polydentate ligand complexes
stereoselectivity in reactions of metal complexes. Part XXI. Kinetics and mechanism of electron transfer between *rac*-$[Co(L)H_2O]^+$ (L = *N*,*N*′-[(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]) and optically active iron(II) complexes, **275–276**, 1

Chiral polydentate ligands
stereoselectivity in reactions of metal complexes. Part XXII. Kinetics and mechanism of electron-transfer reactions between *rac*-$[Co(L)X]^+$ (L = [(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]; X = pyridine, imidazole, 1-methylimidazole and pyrazine) and optically active iron(II) complexes, **275–276**, 9

Chiral separations
high pressure liquid chromatography resolution of racemic mixtures of chiral tricarbonyl arene chromium(0) and cyclopentadienyl-titanium(IV) complexes, **268**, 323

Chloranilate bridged complexes
exchange coupling across multiple-atom bridges: crystal structure and magnetic properties of a chloranilate bridged dicopper(II) complex, **274**, 111

Chloride complexes
the synthesis and characterization of *trans*-$ReCl_2(dppe)_2$ and α-$Re_2Cl_4(dppe)_2$, **268**, 151
complexes of niobium and tantalum with chloride, sulfide and oxide ligands: synthesis and crystal structure of the (tetrachloro)bis-$(\eta^2\text{-disulfido})(\mu\text{-oxo})(\mu\text{-}\eta^2{:}\eta^2\text{-disulfido})$ bismetallates $[Cl_2(S_2)Nb(\mu\text{-}O)(\mu\text{-}S_2)Nb(S_2)Cl_2]^{2-}$ and $[Cl_2(S_2)Ta(\mu\text{-}O)$-$(\mu\text{-}S_2)Ta(S_2)Cl_2]^{2-}$ as their tetraphenylphosphonium salts, **274**, 251
chloride ion transfer during the U(IV)/U(III) reduction of UCl_4 in tetrahydrofuran (THF), studied by cyclic voltammetry; synthesis and molecular structure of $[NBu_4][buCl_5(THF)]$, **279**, 210
kinetics of the photolysis of bis(triethylphosphine)dichloro-platinum(II) in chloroform, **282**, 217

Chloride transfer
chloride ion transfer during the U(IV)/U(III) reduction of UCl_4 in tetrahydrofuran (THF), studied by cyclic voltammetry; synthesis and molecular structure of $[NBu_4][buCl_5(THF)]$, **279**, 210

N-Chloroacetylglycine complexes
investigation on coordination ability of *N*-chloroacetylglycine towards Cu(II) in solid and solution state, **268**, 205

Chloroalkanes
thermal and photochemical reactions of bis(8-quinolinolato)-copper(II), bis(8-quinolinethiolato)copper(II) and (8-quinolinolato,8-quinolinethiolato)copper(II) with chloroalkanes, **273**, 403

Chlorocadmate compounds
organic–inorganic composite materials: structural archetypes of linear polymeric chlorocadmates(II), **272**, 252

Chlorophenolate complexes
transition metal compounds of chlorophenols: synthesis, characterization and single-crystal X-ray structures of (*N*,*N*,*N*′,*N*′-tetramethylethylenediamine)bis(2,4,6-trichlorophenolato)cobalt(II) and bis(*N*-methylimidazole)bis(2,6-dichlorophenolato)copper(II), **271**, 19

Chromium complexes
the kinetics and mechanism of ligand exchange in photogenerated $(\eta^1\text{-bromobenzene})Cr(CO)_5$/alkene systems, **267**, 49
DNA nicking by a trinuclear chromium complex, **268**, 211
high pressure liquid chromatography resolution of racemic mixtures of chiral tricarbonyl arene chromium(0) and cyclopentadienyl-titanium(IV) complexes, **268**, 323
the first X-ray structures of chromium(III) sulfoxide complexes, **269**, 229
syntheses and structural characterization of chromium(III) complexes containing hydrotris(1-pyrazolyl)borate, **269**, 302
bonding modes of arenes in photogenerated (arene)$Cr(CO)_5$ complexes, **271**, 105
comparison of the neutral benzyldiphenylphosphine complexes $C_6H_6Cr(CO)_2PPh_2Bz$ and $C_5H_5Mn(CO)_2PPh_2Bz$ with the

isoelectronic manganiobenzyldiphenylphosphonium salts $[C_5R_5Mn(CO)(NO)PPh_2Bz]BF_4$, **272**, 101

first solid state synthesis of the racemic compound and enantiomer of dinuclear Λ,Λ-di-μ-hydroxo-bis[bis((1*S*,2*S*)-1,2-*trans*-cyclohexanediamine)chromium(III)] and Δ,Δ-di-μ-hydroxo-bis-[bis((1*R*,2*R*)-1,2-*trans*-cyclohexanediamine)chromium(III)] bromides, **274**, 210

bis(2-aminomethylpyridine) complexes of Cr(III) and Co(III), **277**, 177

a series of $[Cr_2(\mu\text{-}OH)_n(\mu\text{-}RCO_2)_{3-n}(bispma)_2]$ type chromium(III) dinuclear complexes ($n = 1, 2$) with tridentate bis(2-pyridylmethyl)amine (bispma): synthesis, crystal structure, and magnetic properties, **278**, 15

kinetic studies using Co(III) and Cr(III) complexes of 1,4,7,11-tetraazaundecane (2,2,3-tet) and the crystal structure of *trans*-$[CrCl_2(2,2,3\text{-tetH})(OH_2)]Cl\cdot ClO_4$, **278**, 24

ground and electronically excited states of $Cr(CO)_4$(bipyridine): energy factored force field analysis of CO stretching vibrations and resonance Raman study, **278**, 83

bonding of halogenated arenes in photogenerated (arene)$M(CO)_5$ complexes (M = Cr, Mo, W), **278**, 202

structures, dynamic behaviour in solution and cross-coupling reactions of the α-palladiated (η^6-alkylarene)tricarbonylchromium complexes containing a palladium–chromium bond, **280**, 71

nephelauxetic series of in-plane ligands for square-pyramidal nitridochromium(V) complexes as studied by means of ESR spectroscopy, **281**, 207

isomerization and racemization of (acetylacetonato)bis(4,4′-dimethyl-2,2′-bipyridine 1,1′-dioxide)chromium(III), (acetylacetonato)bis(4,4′-dimethoxy-2,2′-bipyridine 1,1′-dioxide)-chromium(III) and (acetylacetonato)bis(4,4′-diethoxy-2,2′-bipyridine 1,1′-dioxide)chromium(III) complexes, **281**, 221

Cinnamic acid amide complexes

binuclear and tetranuclear complexes prepared by reactions of dodecacarbonyltriruthenium with the N-substituted cinnamic acid amides. Crystal and molecular structures of $[Ru_2(\mu\text{-}H)(\mu\text{-}\eta^3\text{-}PhC{=}CHC(O)N(CH_2)_4)(CO)_6]$ and $[Ru_4(\mu_3\text{-}H)_2(\mu\text{-}\eta^3\text{-}PhC{=}CHC(O)N(CH_2)_4)_2(CO)_{10}]$, **280**, 243

Cisplatin

synthesis of two maleimide derivatives of *cis*-configurated platinum(II) complexes for the preparation of chemoimmunoconjugates, **267**, 133

imidazole release from the antitumor-active ruthenium complex imidazolium *trans*-tetrachlorobis(imidazole)ruthenate(III) by biologically occurring nucleophiles, **267**, 137

Clay-catalysed

salt-induced peptide formation from amino acids in the presence of clays and related catalysts, **272**, 89

Clotrimazole complexes

toward a novel metal based chemotherapy against tropical diseases. 4. Synthesis and characterization of new metal-clotrimazole complexes and evaluation of their activity against *Trypanosoma cruzi*, **275–276**, 528

Cluster complexes

new efficient synthesis of the triangular cluster $[W_3Se_4(H_2O)_9]^{4+}$ and X-ray structure of the first mixed-metal seleno-bridged cubane-type tungsten–tin cluster $(Me_2NH_2)_6[(SCN)_9W_3Se_4SnCl_3]\cdot 0.5H_2O$, **269**, 292

infrared spectroelectrochemical and electrochemical kinetics studies of the reaction of nickel cluster radicals $[Ni_3(\mu_2\text{-}dppm)_3(\mu_3\text{-}L)(\mu_3\text{-}I)]^+$(L = CNR, R = CH_3, i-C_3H_7, C_6H_{11}, $CH_2C_6H_5$, t-C_4H_9, 2,6-$Me_2C_6H_3$; L = CO) with carbon dioxide, **270**, 111

catalytic reduction of acetylene in the presence of molybdenum and iron clusters, including FeMo cofactor of nitrogenase, **270**, 221

synthesis of a new series of Nb(IV) β-diketonates with the $Nb_2S_4^{4+}$ core: crystal structure of $Nb_2S_4(CF_3COCHCOCH_3)_4$, **271**, 222

synthesis and properties of $[M_6X_{12}(C_3H_7NO)_6]X_2$ (M = Nb, Ta; X = Cl, Br), **271**, 239

steric limitations in associative substitution reactions of $Os_3(CO)_9(\mu\text{-}C_4Ph_4)$, **274**, 82

electrophilic reactivity of arene ligands coordinated to ruthenium carbido cluster complexes, **274**, 201

polynuclear cobalt cluster compounds formed via sulfur–carbon bond cleavage of 2,2-dicyano-1,1-ethylenedithiolate, **274**, 229

NMR characterization of four new isomers of $H(\mu\text{-}H)Os_3(CO)_{10}$-(phosphine), **275–276**, 521

synthesis and characterization of the tungsten–gallium heterometallic cluster $Na_2[W_3GaO_4(O_2CEt)_8]_2$, **277**, 130

pentamethylphenylcopper(I): a square-planar tetranuclear cluster, **277**, 232

electronic absorption and MCD spectra for the $Pt(PPh_3)(AuPPh_3)_6^{2+}$ and $Pt(H)(PPh_3)(AuPPh_3)_7^{2+}$ cluster complexes, **278**, 113

preparation of tetrabutylammonium octa-μ_3-bromohexa(trifluoromethanesulfonato)octahedro-hexamolybdate(2−), $(Bu_4N)_2[Mo_6Br_8^i(CF_3SO_3)_6^a]$ and other derivatives containing the $Mo_6Br_8^{4+}$ core, **278**, 241

the use of photogenerated intermediates in the study of cluster build-up reactions: the generation of $Ru_6C(CO)_{17}$ from $Ru_3(CO)_{12}$, **278**, 250

chiroptical properties of heterotrimetallic 'wing-tip' butterflies; synthesis and crystal structure of $(\mu\text{-}H)Ru_2Fe_2PdC(CO)_{12}(\eta^3\text{-}\beta\text{-}C_{10}H_{15})$, **280**, 275

synthesis, cyclic voltammetry and $^{31}P\{H\}$ NMR spectra of $[M_2E_2L_3]^{2+}$ clusters; M = Ni, Pd or Pt; E = S, Se or Te; L = bis(diphenylphosphino)methane, 1,2-bis(diphenylphosphino)-ethane, 1,3-bis(diphenylphosphino)propane, or 1,4-bis-(diphenylphosphino)butane, **281**, 174

chemistry of rare earth hypocoordinate oxocluster cations: formation of novel oxoacetate clusters, **281**, 201

preparation and characterization of twin-cubane-type clusters with molybdenum–copper–sulfur cores: X-ray structure determination of $[\{(NH_3)_6(H_2O)(OH)Mo_3CuS_4(O)\}_2]Cl_4\cdot 8H_2O$, **282**, 50

a bimetallic main group oxide cluster of the oxovanadium borate system: $(H_3NCH_2CH_2NH_3)_4[(VO)_{12}O_4\{B_8O_{17}(OH)_4\}_2\{Mn(H_2O)_2\}_2]\cdot H_2O$, **282**, 123

Cluster compounds

stepwise substituting magnetic centres in a giant cluster host system, **271**, 9

Clusters

generation of mixed iron chalcogenide clusters $MFeX_n^+$ (M = V, Fe; X = O, S; $n = 2, 3$) in the gas phase, **273**, 111

Cobalamin complexes

the crystal structure of trifluoromethylcobalamin, a coenzyme B_{12} analog, **267**, 305

Cobalt complexes

synthesis and crystal structure of the *trans*-dichloro(*N*,*N*′-bis-[2(*S*)-pyrrolidylmethyl]-ethane-1,2-diamine)cobalt(III) tetrachlorocobaltate(II) complex, **267**, 127

synthesis and rearrangements of the pentaamminecobalt(III) linkage isomers of some amidic acids, **268**, 85

syntheses and structure of dinuclear cobalt(II) complexes of the polypyridine ligand 6,6′-bis[bis(2-pyridylmethyl)aminomethyl]-2,2′-bipyridine, **268**, 159

LMU-3: a new cobalt aluminophosphate with exclusively five-coordinated aluminium and octahedrally coordinated cobalt, **269**, 73

intramolecularly alkylated Cosalen complexes: thermolysis and photolysis of coenzyme B_{12} models with a cobalt-to-ligand carbon bridge, **269**, 203

η^3-allyl–cobalt(II) complexes, **270**, 60

the dynamics of formation of the O_2–Co^{II} bond in the cobalt(II) cyclidene complexes, **270**, 151

rate constants and activation parameters for organo–cobalt porphyrin bond homolysis from NMR relaxation times, **270**, 197

on the mechanism of silylformylation catalyzed by Rh–Co mixed metal complexes, **270**, 279
transition metal compounds of chlorophenols: synthesis, characterization and single-crystal X-ray structures of (*N,N,N′,N′*-tetramethylethylenediamine)bis(2,4,6-trichlorophenolato)cobalt(II) and bis(*N*-methylimidazole)bis(2,6-dichlorophenolato)copper(II), **271**, 19
alternative reaction path for the photochemical reaction of tris(oxalato)cobaltate(III) in various aprotic solvents in the presence of oxygen, **271**, 29
metal complexes of Imazapyr, a herbicide provided with efficient metal-chelating ability: crystal structure of the cobalt(III) and manganese(II) complexes, **272**, 68
synthesis and coordinating properties of the functionalized macrocyclic ligand 1,4-bis(1-methylimidazol-2-ylmethyl)-7-carboxymethyl-1,4,7-triazacyclononane. Crystal structures of cobalt(III), nickel(II) and zinc(II) complexes, **273**, 151
hydrophobic vitamin B_{12}. Part 15. Carbon-skeleton rearrangement reactions mediated by hydrophobic vitamin B_{12} covalently bound to an anionic lipid species in aqueous media, **273**, 299
structural and electronic studies of the coordination of 6-(thiazol-2-yl)-2,2′-bipyridine and related systems to Fe(II), Co(II) and Ni(II), **274**, 32
anionopentaaminecobalt(III) complexes with polyamine ligands. 29. Relative substitution lability in structurally related chloropentaaminecobalt(III) complexes, **274**, 42
cobalt(III) complexes of crown thioethers: the crystal structure of (1,4,7,11,14,17-hexathiacycloeicosane)cobalt(III) tetrafluoroborate dihydrate, **274**, 192
polynuclear cobalt cluster compounds formed via sulfur–carbon bond cleavage of 2,2-dicyano-1,1-ethylenedithiolate, **274**, 229
stereoselectivity in reactions of metal complexes. Part XXI. Kinetics and mechanism of electron transfer between *rac*-$[Co(L)H_2O]^+$ (L = *N,N′*-[(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]) and optically active iron(II) complexes, **275–276**, 1
stereoselectivity in reactions of metal complexes. Part XXII. Kinetics and mechanism of electron-transfer reactions between *rac*-$[Co(L)X]^+$ (L = [(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]; X = pyridine, imidazole, 1-methylimidazole and pyrazine) and optically active iron(II) complexes, **275–276**, 9
4′-(4-fluorophenyl)-2,2′:6′,2″-terpyridine – an aryl-2,2′:6′,2″-terpyridine with an electron-releasing aryl substituent, **275–276**, 359
adducts of nitric oxide with cobaltous tetraphenylporphyrin and phthalocyanines: potential nitric oxide sorbents, **277**, 1
a linear open-chain piperazine–pyridine ligand and its *meso*-helical Co complex, **277**, 55
electrophilic substitution of metal-coordinated pyrazole: nitration, sulfonation and bromination of $[Co(NH_3)_5(pyzH)]^{3+}$ (pyzH = pyrazole), **277**, 89
synthesis and structural studies on metal nitrate complexes with 1,3-dimethyllumazine and 1,3,6,7-tetramethyllumazine: crystal structure of two new cobalt(II) and copper(II) three-dimensionally hydrogen-bonded complexes and a cadmium(II) complex with unusual geometry, **277**, 103
bis(2-aminomethylpyridine) complexes of Cr(III) and Co(III), **277**, 177
reaction of the dinuclear cobalt complex $[Co_2\{\mu\text{-}PPh_2CHCHC(O)CHCHPPh_2\}(CO)_4]$ with two-electron donor ligands; the stabilisation of cobalt–alkene bonds by incorporation of alkenes into a bridging diphosphine ligand, **277**, 186
metal binding properties of three Cys_2X_2 (X = His, Asp) metallothionein-related peptides, **278**, 10
kinetic studies using Co(III) and Cr(III) complexes of 1,4,7,11-tetraazaundecane (2,2,3-tet) and the crystal structure of *trans*-$[CrCl_2(2,2,3\text{-tetH})(OH_2)]Cl\cdot ClO_4$, **278**, 24
spectroscopic and magnetic properties and structure of a five-coordinate, O_2-binding cobalt(II) Schiff base complex and of the copper(II) analogue, **278**, 127
complexes of a new bidentate chelating pyridyl/sulfonamide ligand with copper(II), cobalt(II) and palladium(II): crystal structures and spectroscopic properties, **278**, 178
electrophoretic study of cationic cobalt(III) complexes with $[N_2O]$ and $[N_2O_2]$ Schiff base ligands, **280**, 295
synthesis, structure, spectroelectrochemical and theoretical investigations of $[(\eta^5\text{-}Me_5Cp)\overline{Co(\eta^4\text{-}MeC{=}CMe\text{–}CMe{=}CMe)Sn}\{CH(SiMe_3)_2\}_2]$. Formation of a cobalt complexed stannacycle via a mixed alkyne–stannylene cycloaddition reaction, **281**, 53
syntheses and crystal structures of geometrical isomeric pairs: *trans*- and *cis*-$(PPh_4)[Co(acac)_2(CN)_2]$ and *trans*- and *cis*-$[Co(acac)_2(PMe_3 \text{ or } PEt_3)_2]PF_6$ (acac = pentane-2,4-dionate), **281**, 77
the reaction of cobalt powder with tetraiodo{1,2-bis(dibenzylphosphino)ethane} to form 1,2-bis(dibenzylphosphino)-ethane cobalt diiodide, $Co(dBzP_2)I_2$; and the X-ray crystal structure of the diphosphinodioxide complex, $Co\{dBzP_2(O)_2\}I_2$, **282**, 25
electrochemical, spectrochemical characterization and crystal structure of (1,4,8,11-tetrazacyclotetradecane)(acetylacetonato)cobalt(III) complexes, **282**, 173
structures and electrochemical properties of the Co(III) ternary complexes containing NO_3-type tripodal tetradentate ligands and amino acids: effect of the outer coordination sphere on the electrochemical properties, **283**, 160

Cobalt(II) substitution
spectroscopic distinction between two Co(II) ions substituted for types 1 and 2 Cu in nitrite reductase, **275–276**, 289

Cobalt(III) complexes
hydrolysis of nerve agent simulants by synergistic effects of tetraminecobalt(III) and microemulsions, **269**, 287
kinetics and mechanism of the axial substitution reactions of (ligand)bis(4,5-dichloro-1,2-benzosemiquinonediiminato)-cobalt(III) complexes, **270**, 326

Cobalt powder
the reaction of cobalt powder with tetraiodo{1,2-bis(dibenzylphosphino)ethane} to form 1,2-bis(dibenzylphosphino)-ethane cobalt diiodide, $Co(dBzP_2)I_2$; and the X-ray crystal structure of the diphosphinodioxide complex, $Co\{dBzP_2(O)_2\}I_2$, **282**, 25

Cobinamide complexes
thermodynamics of the binding of nitrogenous ligands to β-trifluoromethylcobinamide and β-cyanomethylcobinamide, **279**, 178

Co–CH_3 stretching mode
the Co–CH_3 bond in Schiff base B_{12} models: influence of the *trans* and equatorial ligands as assessed by Fourier transform Raman spectroscopy, **275–276**, 90

Coenzyme B_{12} models
intramolecularly alkylated Cosalen complexes: thermolysis and photolysis of coenzyme B_{12} models with a cobalt-to-ligand carbon bridge, **269**, 203

Complex stabilities
solution behaviour and zinc complexation of tripeptides with cysteine and/or histidine at both termini, **272**, 115

Composite materials
organic–inorganic composite materials: structural archetypes of linear polymeric chlorocadmates(II), **272**, 252

Conformation
tetrahedral coordination by a seven-membered chelate ring, **275–276**, 279

Conformational isomerization
isomerization and racemization of (acetylacetonato)bis(4,4′-dimethyl-2,2′-bipyridine 1,1′-dioxide)chromium(III), (acetylacetonato)bis(4,4′-dimethoxy-2,2′-bipyridine 1,1′-dioxide)-

chromium(III) and (acetylacetonato)bis(4,4′-diethoxy-2,2′-bipyridine 1,1′-dioxide)chromium(III) complexes, **281**, 221

Conglomerate crystallization
the phenomenon of conglomerate crystallization. Part 51. The crystallization behavior of nickel dinitro amine complexes. *Trans*-[Ni(3,2,3-tet)$(NO_2)_2$]·H_2O (**I**), *mer*-[Ni(N5-Me-1,5,9-triazanonane)$(NO_2)_2$ (**II**), *cis*-[Ni(tren)$(NO_2)_2$]·H_2O (**III**) and *cis*-[Ni(edda)$(H_2O)_2$]·H_2O (**IV**) and control of the bonding mode of the nitrite ligand by the choice of the amine, **282**, 96

π-Conjugated polymers
new type of polymer containing 1,1′-ferrocenediyl unit in π-conjugated main chain. Preparation and redox behavior, **269**, 310

η^2-Coordinated pyrazines
an asymmetrically-coordinated 2,3-dimethylpyrazine bridging ligand in the (II,II) binuclear ion, $\{[Ru^{II}(hedta)]_2(2,3\text{-}Me_2pz)\}^{2-}$ ($hedta^{3-}$ = *N*-hydroxyethylethylenediaminetriacetate), **268**, 279

Coordination polymer complexes
the structural characterization of the novel ribbon sheet, $[Cu_2Cl_2(pyz)]_n$ (pyz = pyrazine), **267**, 143
mixed ligand copper(II) coordination polymers constructed by Cu–bpm–Cu dimer unit (bpm = 2,2′-bipyrimidine) as a building block. Crystal structures and magnetic properties of $\{[Cu(bpm)(SO_4)]\cdot(H_2O)\}_n$, $[Cu_2(bpm)(suc)_{0.5}(ClO_4)_2(OH)(H_2O)_2]_n$ and $\{[Cu(bpm)_{1.5}(suc)_{0.5}](ClO_4)(H_2O)_2\}_n$ (suc = succinate), **283**, 80

Copolymerization
palladium(II)-catalyzed alternating copolymerization of alkenes with sulfur dioxide, **270**, 8

Copper-63 NQR
crystal structure and ^{63}Cu NQR of bis[1-aza-2-(1-pyrrolidinyl)]-copper(I) chloride and bromide, **271**, 231

Copper complexes
antitumour copper(II) salicylaldehyde benzoylhydrazone (H_2sb) complexes: physicochemical properties and the single-crystal X-ray structures of $[\{Cu(H_2sb)(CCl_3CO_2)_2\}_2]$ and $[\{Cu(Hsb)(ClO_4)(C_2H_5OH)\}_2]$ and the related salicylaldehyde acetylhydrazone (H_2sa) complex, $[Cu(Hsa)Cl(H_2O)]\cdot H_2O$, **267**, 27
the structural characterization of the novel ribbon sheet, $[Cu_2Cl_2(pyz)]_n$ (pyz = pyrazine), **267**, 143
planar or tetrahedral coordination in copper(II) complexes induced by bromo substitution, **267**, 151
three dinuclear copper(II) carboxylates with the paddle-wheel cage structure as intermediates in copper(II) catalyzed oxidations of carboxylic acids. X-ray crystal structures of [tetrakis(diphenyl acetato-μ-*O*,*O*′)bis(acetonitrile-*N*)dicopper(II)] tetrakis-(acetonitrile), [tetrakis(diphenylacetato-μ-*O*,*O*′)bis(acetone-*O*)dicopper(II)] and [tetrakis(1-phenyl-1-carboxylato-μ-*O*,*O*′-*cyclo*-pentane)-bis(ethanol-*O*)dicopper(II)] bis(ethanol), **267**, 183
coordination compounds of the rigid rod ligand 1-(2-pyridyl)-2-(2-thienyl)ethyne (pythio). Single crystal X-ray structures of $[Cu(pythio)_2(NO_3)_2]$ and $[Zn(pythio)_2Cl_2]$, **267**, 193
complexes of the terdentate N-donor ligand 6-(2-aminophenyl)-2,2′-bipyridine (L): crystal structures of mononuclear $[ZnL_2][PF_6]_2$ and tetranuclear $[\{CuL(MeCN)\}_4(\mu^4\text{-}PO_4)][PF_6]_5$ containing an unusual μ^4-bridging phosphate ion, **267**, 239
effect of hydrogen bonding on coordination in *trans*-di(*Z*-amino-benzoato-*O*)di(1,3-diaminopropane-*N*,*N*′)copper(II) (*Z* = 2, 3 or 4) complex units, **268**, 21
synthesis, X-ray structure, spectroscopic and magnetic properties of [bis(4-amino-3,5-bis(pyridin-2-yl)-1,2,4-triazole-*N*′,*N*1)-(aqua)copper(II)]bis(hydrogensulfate), **268**, 37
X-ray structures of $Cu(poph)Cl_2$, $[Cu(pxo)Cl_2]_2$ and $[Cu(pxo)(pxo\text{–}H)Cl]_2$ (poph = 2-pyridinecarboxaldehyde 1-oxide 2′-pyridinylhydrazone; pxo = 2-acetylpyridine 1-oxide oxime). One monomeric structure and two *N*-oxide-bridged dimeric structures, involving multidentate pyridine *N*-oxide ligands, **268**, 135
investigation on coordination ability of *N*-chloro-acetylglycine towards Cu(II) in solid and solution state, **268**, 205
crystal structure, spectroscopic, magnetochemical, thermoanalytical and electrochemical properties of binuclear copper(II) complexes of Suprofen, **268**, 239
structure and magnetic properties of a linear oximato-bridged tetranuclear copper(II) complex, **268**, 263
assembly of a mixed-valence Cu(I/II) system coupled by multiple hydrogen bonding through tetrahydroxybenzoquinone, **268**, 317
nitrosamine complexes of $SbCl_5$ and $CuCl_2$, **269**, 7
electronic spectra and structures of Cu^{2+} and Ni^{2+} complexes with dioxotetraazacycloalkanediacetates and their diester derivatives, **269**, 224
synthesis and single crystal X-ray structure of a novel trinuclear copper(II) methoxide complex, **269**, 241
synthesis, structure, and properties of {*N*,*N*′-bis[2-(1-pyrazolyl)-ethyl]pyridine-2,6-dicarboxamido}copper(II), **269**, 269
synthetic, spectroscopic and X-ray crystallographic studies on copper(II) complexes with pyruvic acid and pyridoxal thiosemicarbazones, **269**, 297
X-ray diffraction and EPR studies of binuclear complex $[CuCl(C_6H_9N_3O_3)_2\cdot H_2O]_2Cl_2$ crystals, **269**, 326
interaction of mono- and triphosphate anions with a protonated polyaza macrocycle and its Cu(II) complexes, **270**, 207
monovalent copper as a potential catalyst for formation of acetaldehyde via the migration of methyl radicals to the coordinated carbonyl in the complex $(CO)Cu^{II}\text{–}CH_3^+$, **270**, 440
complexes of copper(I), silver(I), nickel(II) and cadmium(II) with 6-tert-butyldimethylsilylpyridine-2-thionate (6-Bu^{t}SidmepyS$^-$): crystal structure of $[Cu_2(6\text{-}Bu^tSidmepyS)_2dppm_2]$, **271**, 1
transition metal compounds of chlorophenols: synthesis, characterization and single-crystal X-ray structures of (*N*,*N*,*N*′,*N*′-tetramethylethylenediamine)bis(2,4,6-trichlorophenolato)cobalt(II) and bis(*N*-methylimidazole)bis(2,6-dichlorophenolato)copper(II), **271**, 19
[cyclo{tetrakis-$\mu_3(\eta^2\text{-}S,\eta^1\text{-}N)$-(1,3-thiazolidine-2-thionato)-bis-(η^1-S)-(1,3-thiazolidine-2-thione)tetracopper(I)}]: electrochemical synthesis, characterisation and crystal structure at 160 K, **271**, 57
synthesis and crystal structure of two ternary dicopper(I) complexes having the unsymmetrical coordination arrangement bridged by 1,8-naphthyridine (napy). $[Cu_2(napy)_2(Me_2CO)](PF_6)_2\cdot 2Me_2CO$ and $[Cu_2(napy)_2(dppm)(CH_3CN)](PF_6)_2$, **271**, 129
ligand behaviour of a bis-pyrazinecarboxamide copper(II) complex towards a copper(II) β-diketonate. X-ray crystal structures of $[Cu(bpzpn)(H_2O)]\cdot 3H_2O$ and $[Cu(bpzpn)Cu(hfacac)_2]_2$ (H_2bpzpn = *N*,*N*′-bis(2′-pyrazinecarboxamide)-1,3-propane, Hhfacac = hexafluoroacetylacetone), **271**, 210
synthesis and physical studies of copper(I) complexes with the ambidentate ligand (1-piperidyl-2-diphenylphosphino)ethane. X-ray structural characterization of a copper(I) dimer, **271**, 217
crystal structure and ^{63}Cu NQR of bis[1-aza-2-(1-pyrrolidinyl)]-copper(I) chloride and bromide, **271**, 231
synthesis and characterization of copper(I) complexes with triphenylphosphine and heterocyclic thione ligands: the crystal structure of (thiazolidine-2-thione)(bis-triphenylphosphine)copper(I) chloride, **271**, 243
Cu^{II}–herbicide complexes: structure and bioactivity, **272**, 24
Ni(II) and Cu(II) Schiff base complexes with an extended H-bond network, **272**, 283
stopped-flow spectrophotometric study on the reactions between copper(II) ions and some dithiophosphates, **272**, 291
a new class of biomimetically relevant 'Scorpionate' ligands. III. The bis(pyrazolyl)methane(phen-2′-ol)s: synthesis and structural

characterization of mono and dinuclear copper(II) complexes, **273**, 14
stability of binary and ternary copper(II) complexes of the diphosphate analogue, methylphosphonylphosphate, in aqueous solution, **273**, 101
copper(II) complexes with aminosugar derived Schiff bases as ligands, **273**, 191
a reversible Cu^{II}/Cu^{I} electron transfer centre based on a flexible di-imidazole ligand, **273**, 225
copper complexes modelling the interaction between benzolamide and Cu-substituted carbonic anhydrase. Crystal structure of Cu(bz)-$(NH_3)_4$ complex, **273**, 334
thermal and photochemical reactions of bis(8-quinolinolato)-copper(II), bis(8-quinolinethiolato)copper(II) and (8-quinolinolato,8-quinolinethiolato)copper(II) with chloroalkanes, **273**, 403
synthesis and properties of a new tetraaza macrocyclic ligand containing one N-acetic acid pendant arm and its copper(II) and nickel(II) complexes, **274**, 24
ternary copper(II) complexes with the versatile 4,7-dihydro-5-methyl-7-oxo-[1,2,4]triazolo[1,5-*a*]pyrimidine ligand, **274**, 53
general synthesis of 'salicylaldehyde half-unit complexes': structural determination and use as synthon for the synthesis of dimetallic or trimetallic complexes and of 'self-assembling ligand complexes', **274**, 73
exchange coupling across multiple-atom bridges: crystal structure and magnetic properties of a chloranilate bridged dicopper(II) complex, **274**, 111
novel trigonal-bipyramidal copper(II) complexes containing one discrete amino acid chelate: crystal structures of [Cu(L-val)-(pmdt)]ClO_4 and [Cu(aiba)(pmdt)]$ClO_4 \cdot H_2O$ (L-val = L-valinate ion, aiba = α-aminoisobutyrate ion, and pmdt = *N,N,N′,N″,N″*-pentamethyldiethylenetriamine), **274**, 247
crystal structure, EPR and magnetic susceptibility studies of tetrakis[μ-(β-alanine)-*O,O′*]dichlorodicopper(II) dichloromonohydrate, **275–276**, 28
polymeric and dimeric magnetic properties of square planar Cu(II) species controlled by hydrogen bond networks: $[Cu^{II}\{OCO\text{-}2,6\text{-}(CH_3CONH)_2C_6H_3\}_2(H_2O)_2] \cdot nH_2O$ ($n = 1, 4$), **275–276**, 43
coordination chemistry of polyhydroxy acids: role of the hydroxy groups, **275–276**, 130
binding properties and crystal structures of azamacrocycles containing nitrophenol side arms, **275–276**, 168
ESR and NMR investigation of structure and dynamics of the Cu(II) complex of melanostatin in DMSO solution, **275–276**, 274
metallodrug complexes that mediate DNA and lipid damage via sulfite autoxidation: copper(II) famotidine and iron(III) bis(salicylglycine), **275–276**, 314
dynamic properties of bovine Cu,Zn superoxide dismutase from crystallographic data, **275–276**, 349
L- and D-cysteine derivatives of β-cyclodextrin: different molecular recognition properties of their copper(II) complexes for amino acids, **275–276**, 395
copper(I) and copper(II) complexes of polydentate phosphinoamines, **275–276**, 401
potentiometric and spectroscopic studies on the copper(II) and nickel(II) complexes of tripeptides of methionine, **275–276**, 440
toward a novel metal based chemotherapy against tropical diseases. 4. Synthesis and characterization of new metal-clotrimazole complexes and evaluation of their activity against *Trypanosoma cruzi*, **275–276**, 528
kanamycin revisited: a combined potentiometric and spectroscopic study of copper(II) binding to kanamycin B, **275–276**, 546
EPR characterization of a series of mono- and bis-thiosemicarbazone copper(II) complexes, **275–276**, 557
structural study and electron exchange kinetics of the bis(2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline)copper(II)/(I) couple in acetonitrile, **277**, 31
synthesis and structural studies on metal nitrate complexes with 1,3-dimethyllumazine and 1,3,6,7-tetramethyllumazine: crystal structure of two new cobalt(II) and copper(II) three-dimensionally hydrogen-bonded complexes and a cadmium(II) complex with unusual geometry, **277**, 103
crystal structure and characterization of binuclear copper(II) complex with *N*-(2-hydroxyphenyl)-3-methoxycarbonyl-4,6-dimethylsalicylamide, **277**, 134
variable coordination modes for 1,1′-bis(diphenylphosphino)-ferrocene (dppf) mono- and di-chalcogenides or -oxides (dppfE and $dppfE_2$; E = O, S or Se) in copper(I) complexes. Crystal structure of $[\{Cu(dppf)\}_2(\mu\text{-}dppfS_2)](BF_4)_2$ and $[Cu(dppfSe)_2]BF_4$, **277**, 163
pentamethylphenylcopper(I): a square-planar tetranuclear cluster, **277**, 232
bis(1,2-diselenooxalato)nickelate(II) and -cuprate(II) — the first diselenooxalate complexes, **278**, 118
electrostatic ligand–ligand interactions in ternary copper(II) complexes with 3,5-diiodo-L-tyrosine and polar amino acids, **278**, 136
the chemistry of dinitrato-2,2′-bipyridinecopper(II): preparation and characterization of binuclear complexes having a $Cu(\mu\text{-}OH)_2Cu$ and $Cu(\mu\text{-}N_3)_2Cu$ core, **278**, 143
tuning up the T_C in Mn(II)Cu(II) bimetallic planes and design of molecular-based magnets, **278**, 159
complexes of a new bidentate chelating pyridyl/sulfonamide ligand with copper(II), cobalt(II) and palladium(II): crystal structures and spectroscopic properties, **278**, 178
$(C_5H_7N_2O)Cu_3Br_4$: template synthesis of a two-dimensional copper(I) bromide lattice, **279**, 1
exchange interactions through hydrogen-bond bridges. Crystal structure, spectroscopic characterisation and magnetic properties of the complex $[\{Cu(en)\}_2(\mu\text{-egta})] \cdot 4H_2O$ (H_4egta = 3,12-bis-(carboxymethyl)-6,9-dioxa-3,12-diazatetradecanedioic acid), **279**, 58
transition metal complexes with nitronyl nitroxides. Part 1. Crystal structure and properties of square-planar *trans* radical–copper(II)–radical complexes, **279**, 69
solid-state synthesis of cluster compound $[MoCu_3OS_3(PPh_3)_3\text{-}(CH_3COO)]$ with its structure, reactivity and optical non-linearities, **279**, 172
synthesis and metal complexes of a chelating ligand containing thiomorpholine and pyridine: 2,6-bis(thiomorpholinomethyl)pyridine, **279**, 221
bimetallic cyclooligosiloxanolate complexes of copper and nickel, **280**, 282
synthesis of polynuclear copper(II) complexes by controlling the conformation (*cis* or *trans*) of *N,N′*-bis(3-aminopropyl)oxamidate and the coordination mode of 2,2′-bipyrimidine, **281**, 95
crystal structure, spectral and magnetic properties of a new (μ-acetate)(μ-alkoxide) dicopper(II) complex as a model for tyrosinase, **281**, 111
chain-like crystal structure and magnetic properties of [catena-bis(1,2-diaminoethane)copper(II)-μ-dicyano-argentate]-dicyanoargentate, **281**, 134
preparation and characterization of twin-cubane-type clusters with molybdenum–copper–sulfur cores: X-ray structure determination of $[\{(NH_3)_6(H_2O)(OH)Mo_3CuS_4(O)\}_2]Cl_4 \cdot 8H_2O$, **282**, 50
synthesis, structure and characterization of zinc(II), copper(II), zinc(II)barium(II) and copper(II)barium(II) complexes of macrocyclic heteronucleating ligands based on isothiosemicarbazide, **282**, 61
crystal structures and magnetic properties of uniform and alternating azido-bridged (2,2′-bipyridyl)copper(II) chains, **282**, 82
phosphinoamino complexes of copper(I) and silver(I): a combined NMR and IR spectroscopy, and electrospray ionization and fast atom bombardment mass spectrometry analysis, **282**, 119

new cyclam-type copper(II) complexes with amide molecular padlock: synthesis, properties and crystal structure, **282**, 142
synthetic models for biological trinuclear copper clusters. Trinuclear and binuclear complexes derived from an octadentate tetraamine-tetrabenzimidazole ligand, **282**, 180
binuclear nickel(II) and copper(II) complexes of amide-containing two-armed and macrocyclic ligands, **282**, 222
synthesis, crystal structure and magnetic properties of the first single azido-bridged copper(II) chain $[Cu(bpym)(N_3)_2]_n$ (bpym = 2,2′-bipyrimidine), **282**, 252
how non-bonding amino acid side-chains may enormously increase the stability of a Cu(II)–peptide complex, **283**, 1
hydrolysis of phosphate esters catalyzed by copper(II)-triamine complexes. The effect of triamine ligands on the reactivity of the copper(II) catalysts, **283**, 51
mixed ligand copper(II) coordination polymers constructed by Cu–bpm–Cu dimer unit (bpm = 2,2′-bipyrimidine) as a building block. Crystal structures and magnetic properties of $\{[Cu(bpm)(SO_4)](H_2O)\}_n$, $[Cu_2(bpm)(suc)_{0.5}(ClO_4)_2(OH)(H_2O)_2]_n$ and $\{[Cu(bpm)_{1.5}(suc)_{0.5}](ClO_4)(H_2O)_2\}_n$ (suc = succinate), **283**, 80
an infinite chiral single-helical structure formed in Cu(II)-L-/D-glutamic acid system, **283**, 105
ternary complexes in solution. Intramolecular stacking interactions in mixed ligand complexes formed by copper(II), 2,2′-bipyridyl or 1,10-phenanthroline and a pyrimidine-nucleoside 5′-diphosphate (CDP^{3-}, UDP^{3-}, $dTDP^{3-}$), **283**, 193
metal ion–FTS nonapeptide interactions. II. Copper(II) complex equilibria and structures under physiological conditions, **283**, 211

Copper(I) complexes
stability constants of copper(I) and silver(I) complexes with aza-crown ethers and cryptands in dimethylsulfoxide, **279**, 186

Copper(II) complexes
copper(II) complexes with tridentate O,N,O-chelate dianions and additive monodentate donor ligands. I. Crystal structures and EPR results of urea adducts of *N*-salicylidene-α-aminoacidato-copper(II) compared with *N*-salicylideneglycinato-thiourea-copper(II), **268**, 117
copper(II) complexes of pyrazol-3-yl amines. Synthesis, crystal and solution structure, **269**, 117
copper(II) complexes of (*R,S*)-α-hydroxymethylornithine and its N^{δ}-benzoyl derivative, **269**, 279
axial–equatorial interactions in chelates of β-diones. Crystal and molecular structure of bis(3-nitro-2,4-pentanedionato)copper(II), **269**, 332
synthesis of new copper(II) complexes of polydentate ligands containing pyridyl pendant arms. Crystal structure of [*N,N,N′,N′*-tetrakis(2-pyrid-2-yl-ethyl)1,2-ethylenediaminecopper(II)]-bis(hexafluorophosphate) · 1.5 acetone, **278**, 237
spectroscopic characterization of coaxially and π–π stacked binuclear copper(II) complexes of ω,ω′-bis(pyridine-2-carboxamido)-alkanes in acidic solution, **283**, 61

Copper(II) ion trimer
synthesis and single crystal X-ray structure of a novel trinuclear copper(II) methoxide complex, **269**, 241

Copper ligands
electron tunneling in Ru-modified His46Asp azurin. Coupling through the Cu ligands, **275–276**, 470

Copper proteins
characterization of the active site of galactose oxidase and its active site mutational variants Y495F/H/K and W290H by circular dichroism spectroscopy, **275–276**, 175
spectroscopic distinction between two Co(II) ions substituted for types 1 and 2 Cu in nitrite reductase, **275–276**, 289

Cosalen complexes
intramolecularly alkylated Cosalen complexes: thermolysis and photolysis of coenzyme B_{12} models with a cobalt-to-ligand carbon bridge, **269**, 203

Croconate complexes
synthesis, crystal structures and magnetic properties of $[Fe_2(bpym)(C_5O_5)_2(H_2O)_4] \cdot 2H_2O$ and two polymorphs of $[Fe_2(bpym)(C_4O_4)_2(H_2O)_6] \cdot 2H_2O$ (bpym = 2,2′-bipyrimidine), **279**, 127

Cross-coupling
enantioselective catalysis. XVI: regio- and enantioselectivity in nickel-catalysed cross-coupling reactions of allylic substrates with Grignard reagents, **269**, 34
synthesis of porphyrin–spacer–quinone compounds via metal-mediated cross-coupling: new systems for probing the relative magnitudes of axial and equatorial electronic coupling at the porphyrin macrocycle in thermal and photoactivated electron transfer reactions, **275–276**, 427
structures, dynamic behaviour in solution and cross-coupling reactions of the α-palladiated (η^6-alkylarene)tricarbonylchromium complexes containing a palladium–chromium bond, **280**, 71

Crown ether complexes
synthesis and crystal structure of rubidium hydrogen oximate complexes with 18-crown-6: is it possible to reach a perfect fit of the rubidium atom inside the crown ether cavity?, **268**, 93
synthesis, characterization and crystal structures of complexes of sodium hexafluorophosphate with dibenzo-18-crown-6 and dibenzo-24-crown-8 macrocycles, **282**, 76
the effect of second-sphere coordination. 7. Isolation of 18-crown-6 ether adducts of ruthenium–ammine complexes, **282**, 247

Crown thioether complexes
cobalt(III) complexes of crown thioethers: the crystal structure of (1,4,7,11,14,17-hexathiacycloeicosane)cobalt(III) tetrafluoroborate dihydrate, **274**, 192

Cryptand complexes
thermodynamic study of complexation of alkali and alkaline-earth metal ions with benzene-containing cryptands, **278**, 6

Cryptands
a large cavity reinforced cryptand for the binding of metal cations and anions, **273**, 326
stability constants of copper(I) and silver(I) complexes with aza-crown ethers and cryptands in dimethylsulfoxide, **279**, 186

Crystallographic data
dynamic properties of bovine Cu,Zn superoxide dismutase from crystallographic data, **275–276**, 349

Crystal packing
investigations into the engineering of inorganic/organic solids: hydrothermal synthesis and crystal structures of aromatic ammonium sodium hydrogen phosphomolybdates(VI), **282**, 10

Crystal structures
synthesis and characterization of new hydrotri(3,5-dimethyl-1-pyrazolyl)borate vanadium halfsandwich complexes, **267**, 19
antitumour copper(II) salicylaldehyde benzoylhydrazone (H_2sb) complexes: physicochemical properties and the single-crystal X-ray structures of $[\{Cu(H_2sb)(CCl_3CO_2)_2\}_2]$ and $[\{Cu(Hsb)(ClO_4)(C_2H_5OH)\}_2]$ and the related salicylaldehyde acetylhydrazone (H_2sa) complex, $[Cu(Hsa)Cl(H_2O)] \cdot H_2O$, **267**, 27
syntheses, structures and properties of hydrosulfido-bridged diiridium and dirhodium complexes, $[Cp^*MCl(\mu_2\text{-}SH)_2MCp^*Cl]$ and $[Cp^*M(\mu_2\text{-}SH)_3MCp^*]^+$ (M = Ir, Rh; $Cp^* = \eta^5\text{-}C_5Me_5$), **267**, 73
a bis(9-methyladeninium) complex of Hg(II) with a highly irregular coordination geometry: $[Hg(9\text{-MeAH-}N7)_2(H_2O)(NO_3)_3]ClO_4$, **267**, 87
synthesis and characterization of molybdenum(II) and molybdenum(V) monothiocarbamate complexes. Crystal and molecular structures of tetrakis(isopropylmonothiocarbamato)dimolyb-

denum(II) and oxobis[di(oxoisopropylmonothiocarbamato)-molybdenum(V)], **267**, 101
synthesis, X-ray and spectroscopic characterization of $[SnI_2(mbit)_2]$-$(I_3)_2 \cdot 2/3I_2$ obtained through the one-step reaction of mbit $\cdot 2I_2$ with tin metal powder (mbit = 1,1′-bis(3-methyl-4-imidazoline-2-thione)methane), **267**, 115
the synthesis, characterization and X-ray crystal structures of the *fac-cis* and *mer-trans* isomers of $ReOCl_3(PMe_3)_2$, **267**, 123
synthesis and crystal structure of the *trans*-dichloro(*N,N′*-bis-[2(*S*)-pyrrolidylmethyl]-ethane-1,2-diamine)cobalt(III) tetrachlorocobaltate(II) complex, **267**, 127
the structural characterization of the novel ribbon sheet, $[Cu_2Cl_2(pyz)]_n$ (pyz = pyrazine), **267**, 143
planar or tetrahedral coordination in copper(II) complexes induced by bromo substitution, **267**, 151
the synthesis and characterization of $[Tc(SPh)_3(N{=}NC_5H_4N)$-$(HN{=}NC_5H_4N)]_2$, a paramagnetic dimer with bridging phenylthiolate ligands. The X-ray structure of $[Tc(SPh)_2$-$(N{=}NC_5H_4N)(HN{=}NC_5H_4N)]_2$, **267**, 165
an infrequent geometry for an element of the first transition series: structural characterization of a distorted eight-coordinate compound $TiCl_4[C_5H_4(PMe_2)_2]_2$, **267**, 173
crystal and molecular structure of protonated (*N*-propyl)-aminomethyl ferrocene, a proton-sensitive redox-responsive fragment, **267**, 177
three dinuclear copper(II) carboxylates with the paddle-wheel cage structure as intermediates in copper(II) catalyzed oxidations of carboxylic acids. X-ray crystal structures of [tetrakis(diphenyl acetato-μ-*O,O′*)bis(acetonitrile-*N*)dicopper(II)] tetrakis-(acetonitrile), [tetrakis(diphenylacetato-μ-*O,O′*)bis(acetone-*O*)dicopper(II)] and [tetrakis(1-phenyl-1-carboxylato-μ-*O,O′-cyclo*-pentane)-bis(ethanol-*O*)dicopper(II)] bis(ethanol), **267**, 183
coordination compounds of the rigid rod ligand 1-(2-pyridyl)-2-(2-thienyl)ethyne (pythio). Single crystal X-ray structures of $[Cu(pythio)_2(NO_3)_2]$ and $[Zn(pythio)_2Cl_2]$, **267**, 193
the reactivity of five-coordinate Ru(II)(1,4-bis(diphenylphosphino)-butane) complexes with the N-donor ligands: ammonia, pyridine, 4-substituted pyridines, 2,2′-bipyridine, bis(*o*-pyridyl)amine, 1,10-phenanthroline, 4,7-diphenylphenanthroline and ethylenediamine, **267**, 209
site-selective PMe_3 addition to the donor–acceptor complex $Ru_2(CO)_6[(Z)$-$Ph_2PCH{=}CHPPh_2]$. X-ray diffraction structure of $Ru_2(CO)_5(PMe_3)[(Z)$-$Ph_2PCH{=}CHPPh_2]$, **267**, 223
the preparation and synthetic potential of chlorovanadium(V and IV) complexes supported by enamines and bis(enamines), **267**, 229
complexes of the terdentate N-donor ligand 6-(2-aminophenyl)-2,2′-bipyridine (L): crystal structures of mononuclear $[ZnL_2][PF_6]_2$ and tetranuclear $[\{CuL(MeCN)\}_4(\mu^4\text{-}PO_4)][PF_6]_5$ containing an unusual μ^4-bridging phosphate ion, **267**, 239
effect of substitution on packing in *trans*-di(Z-chlorobenzoato-*O*)-di(1,3-diaminopropane-*N,N′*)nickel(II) (*Z* = 2, 3 or 4) complexes, **267**, 249
partial hydrolysis of Ti(III) and Ti(IV) chlorides in the presence of $[PPh_4]Cl$, **267**, 271
the preparation and X-ray structure of $[Ho(oxam)_3(H_2O)_3]_4 \cdot 2.75H_2O$, **267**, 299
the crystal structure of trifluoromethylcobalamin, a coenzyme B_{12} analog, **267**, 305
phosphide complexes of divalent europium: synthesis and structural characterization of $Eu[PPh_2]_2(N\text{-}MeIm)_4$, **267**, 309
a tetranuclear silver complex with dppm and i-mnt ligands. Synthesis and structural characterization of $[Ag_4(\mu\text{-dppm})_4(\mu_4\text{-i-mnt})_2] \cdot$ 0.5 DMF (i-mnt = 2,2-dicyano-1,1-ethylenedithiolate; dppm = bis(diphenylphosphino)methane), **267**, 313
the synthesis and characterization of a technetium nitrosyl complex with *cis*-{2-pyridyl,diphenylphosphine} coligands. The X-ray crystal structure of $[TcCl_2(NO)(pyPPh_2\text{-}P,N)(pyPPh_2\text{-}P)]$, **267**, 319
complexes of thallium(I) and lead(II) with the potentially tetradentate ligand bis[3-(2-pyridyl)-pyrazolyl]dihydroborate, **267**, 323
preparation and molecular structure of (*N*-benzyl-1,2-didehydroglycinato) bis(1,10-phenanthroline)ruthenium(II) chloride: $[Ru(N\text{-}Bn\text{-}gly\text{-}H_2)(phen)_2]Cl$, **267**, 329
effect of hydrogen bonding on coordination in *trans*-di(Z-aminobenzoato-*O*)di(1,3-diaminopropane-*N,N′*)copper(II) (*Z* = 2, 3 or 4) complex units, **268**, 21
synthesis, X-ray structure, spectroscopic and magnetic properties of [bis(4-amino-3,5-bis(pyridin-2-yl)-1,2,4-triazole-*N⁴,N1*)-(aqua)copper(II)]bis(hydrogensulfate), **268**, 37
synthesis and reactivity of Ru(II) complexes containing the phosphino-amine $Ph_2PCH_2CH_2NMe_2$, **268**, 69
synthesis and structural studies on three-dimensionally hydrogen-bonded metal perchlorate complexes with 1,3-dimethyllumazine and 1,3,6,7-tetra-methyllumazine. Crystal structure of diaqua-bis-(1,3,6,7-tetramethyl-pteridine-2,4(1*H*,3*H*)-dione-O^4,N^5)-copper(II) perchlorate dihydrate, **268**, 77
synthesis and crystal structure of rubidium hydrogen oximate complexes with 18-crown-6: is it possible to reach a perfect fit of the rubidium atom inside the crown ether cavity?, **268**, 93
synthesis, electrical properties of $[Ni(dmit)_2]^-$ complexes and X-ray crystal structure of $[EDA][Ni(dmit)_2]$ (dmit = 1,3-dithiol-2-thione-4,5-dithiolate), **268**, 103
ligand exchange reactions of $ReCl_4(PPh_3)_2$ with tridentate diacidic ligands with the donor set O–N–O(S). Molecular structures of the resulting oxo-rhenium(V) complexes, **268**, 109
copper(II) complexes with tridentate O,N,O-chelate dianions and additive monodentate donor ligands I. Crystal structures and EPR results of urea adducts of *N*-salicylidene-α-aminoacidato-copper(II) compared with *N*-salicylideneglycinato-thiourea-copper(II), **268**, 117
X-ray structures of $Cu(poph)Cl_2$, $[Cu(pxo)Cl_2]_2$ and $[Cu(pxo)(pxo\text{–}H)Cl]_2$ (poph = 2-pyridinecarboxaldehyde 1-oxide 2′-pyridinylhydrazone; pxo = 2-acetylpyridine 1-oxide oxime). One monomeric structure and two *N*-oxide-bridged dimeric structures, involving multidentate pyridine *N*-oxide ligands, **268**, 135
the synthesis and characterization of *trans*-$ReCl_2(dppe)_2$ and α-$Re_2Cl_4(dppe)_2$, **268**, 151
syntheses and structure of dinuclear cobalt(II) complexes of the polypyridine ligand 6,6′-bis[bis(2-pyridylmethyl)aminomethyl]-2,2′-bipyridine, **268**, 159
synthesis and characterization of uranyl thiolate complexes. Crystal and molecular structures of $[Me_4N][UO_2(NO_3)_2(C_5H_4NS)]$ and $[C_{16}H_{25}N_2S_2Si_2][UO_2(NO_3)_2(C_8H_{12}NSSi)]$, **268**, 163
synthesis and structural characterization of the [2,9-dimethyl-4,7-diaza-2,9-decanedithiolato]oxorhenium(V) complex, **268**, 169
synthesis and crystal structures of complexes containing an Fe→Cd donor–acceptor bond formed by reaction of *trans*-$(CO)_3Fe$-$(Ph_2Ppym)_2$ (Ph_2Ppym = 2-(diphenylphospino)pyrimidine) with CdX_2 (X = Cl, Br, I, SCN, ClO_4), **268**, 177
reactivity of molybdenum complexes containing both NHNPhR and NNPhR ligands (R = Ph, Me) toward hydridotris(pyrazolyl)-borate, Tp and hydridotris(3,5-dimethylpyrazolyl)borate, Tp*, ligands. X-ray crystal structure of $[TpMo(NNMePh)_2Cl]$, **268**, 199
investigation on coordination ability of *N*-chloro-acetylglycine towards Cu(II) in solid and solution state, **268**, 205
square-pyramidal copper(II) complexes of linear tetradentate edda-type ligands forming six-membered rings. Molecular structures of $[Cu(1,3\text{-pdda})(H_2O)]$ and $[Cu(eddp)(H_2O)] \cdot 3.5H_2O$, **268**, 221
crystal structure, spectroscopic, magnetochemical, thermoanalytical and electrochemical properties of binuclear copper(II) complexes of Suprofen, **268**, 239

hydrothermal synthesis and structural characterization of a three-dimensional organically-templated vanadium phosphate $[H_3NCH_2CH_2NH_3][(VO)_3(H_2O)_2(PO_4)_2(HPO_4)]$, **268**, 257
structure and magnetic properties of a linear oximato-bridged tetranuclear copper(II) complex, **268**, 263
the preparation and crystal and molecular structure of the complex *cis*-$PtCl_2$(razoxane), **268**, 313
assembly of a mixed-valence Cu(I/II) system coupled by multiple hydrogen bonding through tetrahydroxybenzoquinone, **268**, 317
nitrosamine complexes of $SbCl_5$ and $CuCl_2$, **269**, 7
reactivity and chemical exchange processes of bis(ether-phosphine)iridium(III) complexes, **269**, 13
'activated cyanide', **269**, 23
clusters $N_2B_4R_6$ from the fusion of rings NB_2R_3: mechanism, **269**, 43
taranakite – the mineral with the longest crystallographic axis, **269**, 47
LMU-3: a new cobalt aluminophosphate with exclusively five-coordinated aluminium and octahedrally coordinated cobalt, **269**, 73
synthesis and coordination properties of amphiphilic 3-oxodithiocarboxylic esters, **269**, 83
syntheses and structure of dimeric yttrium tris[bis(trimethylsilyl)phosphanide]; X-ray structures of $[(thf)LiCl\cdot(thf)_2LiP^tBu_2]_2$ and $Y[CH(SiMe_3)_2]_3$, **269**, 91
some oxidative addition reactions with a diphosphadibora[1.1.0]bicyclobutane: formation of cage compounds and of an unusual trinuclear coordination compound of palladium, **269**, 101
coordination chemistry of perhalogenated cyclopentadienes. XXII. Reaction of tetrabromodiazocyclopentadiene with some halide bridged Rh(I), Ru(II) and Ru(III) complexes. Molecular structure of $[C_5Br_5]Rh(COD)$, a complex with an η^1,η^4-coordinated cyclopentadienyl ligand, **269**, 111
copper(II) complexes of pyrazol-3-yl amines. Synthesis, crystal and solution structure, **269**, 117
tripod ligands of constitutional C_1 symmetry containing a deliberate set of mixed P/S donor groups: synthesis and coordination chemistry, **269**, 143
substitution reactions of $CpRu(PPh_3)_2Cl$ with PPh_2H and dppa, **269**, 157
structures of $HRh(PMe_2Ph)_4$ and $HRh(PPh_2Me)_4$, **269**, 162
synthesis and structures of tin(II) and lead(II) 1-aza-allyls; the $[N(SiMe_3)C(Ph)C(SiMe_3)_2]^-$ ligand, **269**, 181
lanthanide complexes with 1-hydroxyethane-1,1-diphosphonic acid: solvent organization and coordination geometry in crystalline and amorphous solids, **269**, 211
electronic spectra and structures of Cu^{2+} and Ni^{2+} complexes with dioxotetraazacycloalkanediacetates and their diester derivatives, **269**, 224
the first X-ray structures of chromium(III) sulfoxide complexes, **269**, 229
the coordination mode of 3-hydroxypicolinic acid in oxorhenium(V) complexes. Crystal and molecular structures of $(n\text{-}Bu_4N)[ReOCl_3(HO\text{–}C_5H_3N\text{–}CO_2)]$ and $[ReOCl(HO\text{–}C_5H_3N\text{–}CO_2)_2]$, **269**, 235
synthesis and single crystal X-ray structure of a novel trinuclear copper(II) methoxide complex, **269**, 241
synthesis, structure, and properties of {*N,N'*-bis[2-(1-pyrazolyl)-ethyl]pyridine-2,6-dicarboxamido}copper(II), **269**, 269
synthesis and X-ray structure of the thioethereal mercury(II) complex $[N(CH_2CH_2SCHMe_2)_3HgCl]_2(Hg_2Cl_6)$, **269**, 274
synthesis, spectral and structural characterization of the zinc carboxylate $[Zn(2\text{-quinolinecarboxylato})_2(1\text{-methylimidazole})_2]$, **269**, 283
new efficient synthesis of the triangular cluster $[W_3Se_4(H_2O)_9]^{4+}$ and X-ray structure of the first mixed-metal seleno-bridged cubane-type tungsten–tin cluster $(Me_2NH_2)_6[(SCN)_9W_3Se_4SnCl_3]\cdot 0.5H_2O$, **269**, 292
synthetic, spectroscopic and X-ray crystallographic studies on copper(II) complexes with pyruvic acid and pyridoxal thiosemicarbazones, **269**, 297
syntheses and structural characterization of chromium(III) complexes containing hydrotris(1-pyrazolyl)borate, **269**, 302
{*catena*-($\mu_{N,S}$-NCS)[Ni(Medien)(NCS)]}: a new 1D nickel(II) complex with bridging thiocyanato ligands in *cis* position. Crystal structure and magnetic properties, **269**, 313
synthesis and crystal structure of $[Cd\{(c\text{-}C_6H_{11})_2P(Se)C(S)NPh\}_2]$, **269**, 317
X-ray diffraction and EPR studies of binuclear complex $[CuCl(C_6H_9N_3O_3)_2\cdot H_2O]_2Cl_2$ crystals, **269**, 326
axial–equatorial interactions in chelates of β-diones. Crystal and molecular structure of bis(3-nitro-2,4-pentanedionato)copper(II), **269**, 332
synthesis of trichlorooxo-bis(2,6-dimethylphenylisocyanide)rhenium(V) and crystal structure of μ-oxo-bis[dichlorooxo-bis(2,6-dimethylphenylisocyanide)dirhenium(V)], **269**, 347
salicylaldehyde-2-mercaptoanilato(2−)-2-hydroxyphenyl-benzthiazolato-nitrosyl-molybdenum(II) — a chelate which simultaneously contains an azomethine ligand and its cyclic oxidation product, **269**, 350
ligand connected metal clusters. The molecular structures and solid state packing of $\{Ru_3(CO)_{11}\}_2$(bis(diphenylphosphino)ethane) and $\{Ru_3(CO)_{11}\}_2$(1,4-bis(diphenylphosphino)benzene), **270**, 28
rhodium aryloxide complexes containing terdentate nitrogen ligands, C–O bond formation, hydrogen bonding with phenol and oxidative addition reactions. Molecular structure of an Rh(III)–acetyl complex, **270**, 34
synthesis and structures of the cuboidal iron–sulfur–nitrosyl–phosphine clusters $[Fe_4S_3(NO)_4(PR_3)_3]^{0,1+}$ (R = Et, Pr^i, C_6H_{11}), **270**, 46
thermal and photochemical substitution reactions of $CpRe(PPh_3)_2H_2$ and $CpRe(PPh_3)H_4$. Catalytic insertion of ethylene into the C–H bond of benzene, **270**, 77
palladium-catalyzed seven-membered N,O-heterocycle ring formation, **270**, 123
triply-bridged diruthenium(II) 1,4-bis(diphenylphosphino)butane (dppb) and (*R*)-2,2′-bis(diphenylphosphino)-1,1′-binaphthyl (binap) complexes, including structural characterisation of $[(dppb)ClRu(\mu\text{-}D_2O)(\mu\text{-}Cl)_2RuCl(dppb)]$, $[(\eta^2\text{-}H_2)(dppb)Ru(\mu\text{-}Cl)_3RuCl(dppb)]$ and the $[(dppb)ClRu(\mu\text{-}Cl)_3RuCl(dppb)]^-$ anion, **270**, 130
alkylcobalt chelates with Schiff bases derived from a β-diketone bearing both alkyl and aryl groups, **270**, 169
eight-coordinate chelate complexes of zirconium(IV) and niobium(IV): X-ray diffractometric and EPR investigations, **270**, 177
interaction of mono- and triphosphate anions with a protonated polyaza macrocycle and its Cu(II) complexes, **270**, 207
misdirected π-donor ligands: $(\eta^5\text{-}C_5H_5)_2Zr(Cl)(SR)$ with sterically more demanding R groups have lower rotational barriers, **270**, 227
solution and crystal structure of the dihydrogen complex $[Ru(H_2)(H)(PMe_2Ph)_4]PF_6$, an active alkyne hydrogenation catalyst, **270**, 238
silyl reagents (Me_3Si–X) efficiently transfer X to $Ir(H)_2F(P^tBu_2Ph)_2$, **270**, 261
transition metal complexes. IX. Nickel complexes with a chiral azaphosphole ligand, **270**, 273
titanium–carbon functionalities on an oxo surface defined by a calix[4]arene moiety and its redox chemistry, **270**, 298
chemistry of the azine phosphine ligand *Z,E*-$PPh_2CH_2C(Bu^t)$=N–N=$CMe(C_6H_4NO_2$-4): crystal structure of $[Mo(CO)_4\{PPh_2CH_2C(Bu^t)$=N–N=$CMe(C_6H_4NO_2$-4)$\}]$, **270**, 312
the synthesis and structures of tantalum complexes that contain a triamido or a diamidoamine ligand, **270**, 353

preparation of trimethylsilyl alkyne complexes of bis(pentamethylcyclopentadienyl)zirconium, $(\eta^5\text{-}C_5Me_5)_2Zr(RC{\equiv}CSiMe_3)$, and their regioselective reactions with nitrous oxide, **270**, 399

the propensity of alkoxide and aryloxide derivatives of tungsten carbonyls to aggregate in solution. Synthesis and X-ray structures of dinuclear, trinuclear and tetranuclear complexes derived from the $MeOW(CO)_5^-$ anion, **270**, 405

the synthesis of substituted bis[(diarylphosphinomethyl)cyclopentadienyl]zirconocene dichloride complexes for the preparation of heterodimetallic complexes containing early/late transition metal combinations, **270**, 446

unique complexation by the tridentate ligands, 6-diphenylphosphino-2-pyridonate (pyphos) and 6-diphenylphosphino-2-pyridone (pyphosH), on ruthenium(II): crystal structures of (pyphosH)-$RuCl(\mu\text{-}Cl)Mo_2Cl(pyphos)_3$, $RuCl_2(CO)(pyphosH)_2$, $[Ru(pyphos)_3](\mu\text{-}Na)_2[Ru(pyphos)_3]$, **270**, 459

the synthesis of the tripodal phosphine $CH_3C\{CH_2P(m\text{-}CF_3C_6H_4)_2\}_3$ and its rhodium coordination chemistry, **270**, 499

nitrogen atom transfer and redox chemistry of terpyridyl phosphoraniminato complexes of osmium (IV), **270**, 511

synthesis, characterization, and reactivity of organometallic Zr(IV) carboxylate complexes, **270**, 527

ferrocenyl-derived face-capping ligands: syntheses and molecular structures of $[(FcTt)Cu]_4$ and $[FcTt]_2M$ (M = Fe, Co, Ni; $FcTt^-$ = ferrocenyltris((methylthio)methyl)borate), **270**, 543

synthesis of rhodium porphyrin aryls via intermolecular arene carbon–hydrogen bond activation, **270**, 551

a ferrocene–perfluoroarene molecular complex, **270**, 559

complexes of copper(I), silver(I), nickel(II) and cadmium(II) with 6-tert-butyldimethylsilylpyridine-2-thionate ($6\text{-}Bu^tSidmepyS^-$): crystal structure of $[Cu_2(6\text{-}Bu^tSidmepyS)_2dppm_2]$, **271**, 1

syntheses and properties of bivalent metal complexes with *N*-(2-hydroxyethyl)ethylenediamine-*N*,*N*′,*N*′-triacetic acid and the crystal structure of *N*-(2-hydroxyethyl)ethylenediamine-*N*,*N*′,*N*′-triacetato-copper(II) monohydrate, **271**, 13

transition metal compounds of chlorophenols: synthesis, characterization and single-crystal X-ray structures of (*N*,*N*,*N*′,*N*′-tetramethylethylenediamine)bis(2,4,6-trichlorophenolato)cobalt(II) and bis(*N*-methylimidazole)bis(2,6-dichlorophenolato)copper(II), **271**, 19

3-ferrocene-substituted 3-mercaptopropenales and their Schiff bases as ligands for transition metal complexes, **271**, 40

the synthesis and characterization of $\alpha\text{-}Re_2Cl_4(depe)_2$ and $\alpha\text{-}W_2Cl_4(depe)_2$, **271**, 49

[cyclo{tetrakis-$\mu_3(\eta^2\text{-}S,\eta^1\text{-}N)$-(1,3-thiazolidine-2-thionato)-bis-$(\eta^1\text{-}S)$-(1,3-thiazolidine-2-thione)tetracopper(I)}]: electrochemical synthesis, characterisation and crystal structure at 160 K, **271**, 57

synthesis, characterization and study of the redox properties of rhenium(V) diolates, **271**, 65

triphenyltellurium(IV) dichalcogenoimidodiphosphinates containing novel six-membered TeXYP$_2$N chelate rings. X-ray structures of $Ph_3Te[(SPPh_2)(OPPh_2)N]$ and $Ph_3Te[(OPPh_2)_2N]$, **271**, 75

synthesis, spectroscopic and electrochemical study of nickel(II) complexes with tetradentate asymmetric Schiff bases derived from salicylaldehyde and methyl-2-amino-1-cyclopentenedithiocarboxylate, **271**, 83

syntheses and spectroscopic characterizations of the compounds containing univalent coinage metal and tetraselenomolybdate. Crystal structure of $[MoSe_4(AuPPh_3)_2]$, **271**, 93

synthesis, structure and magnetic properties of one-dimensional azide-bridged manganese(III) uniform chain complex Mn(salpn)N_3, **271**, 99

synthesis, characterization and reactivity of bis(diphenylphosphino)-amine nickel complexes, **271**, 112

synthesis and crystal structure of two ternary dicopper(I) complexes having the unsymmetrical coordination arrangement bridged by 1,8-naphthyridine (napy). $[Cu_2(napy)_2(Me_2CO)](PF_6)_2 \cdot 2Me_2CO$ and $[Cu_2(napy)_2(dppm)(CH_3CN)](PF_6)_2$, **271**, 129

synthesis and structure of a novel bis($\mu\text{-}\eta^2$-thianthrene)disilver(I) bis(perchlorate), **271**, 145

dinuclear complexes of first transition series metals with 4,4′-dipyrazolylmethane: characterisation, DNA binding and anticancer properties, **271**, 151

trimethylphosphine halide complexes of titanium, zirconium and hafnium in oxidation states IV, III and II. Crystal structure of $MI_4(PMe_3)_n$ (M = Zr, n = 3; M = Hf, n = 2.5), $M_2X_6(PMe_3)_4$ (M = Hf, Ti, X = I; M = Zr, X = Br) and $(\mu\text{-}\eta^6,\eta^6\text{-}C_6H_6)\text{-}Hf_2I_4(PMe_3)_4 \cdot C_6H_6$, **271**, 180

metal–metal multiply-bonded complexes of technetium. 7. Oxidative decomposition of tetrachlorotetrakis(dimethylphenylphosphino)-ditechnetium(II) by an aminium hexachloroantimonate(V), **271**, 187

the synthesis and characterization of a technetium(III) isodiazene complex. The X-ray crystal structure of $[TcCl_3(N{=}NPh_2)\text{-}(PPh_3)_2]$, **271**, 191

structure of an indium(III) complex with a 15-membered macrocycle, dioxopentaazacyclopentadecanetriacetate, **271**, 207

ligand behaviour of a bis-pyrazinecarboxamide copper(II) complex towards a copper(II) β-diketonate. X-ray crystal structures of $[Cu(bpzpn)(H_2O)] \cdot 3H_2O$ and $[Cu(bpzpn)Cu(hfacac)_2]_2$ (H_2bpzpn = *N*,*N*′-bis(2′-pyrazinecarboxamide)-1,3-propane, Hhfacac = hexafluoroacetylacetone), **271**, 210

synthesis and physical studies of copper(I) complexes with the ambidentate ligand (1-piperidyl-2-diphenylphosphino)ethane. X-ray structural characterization of a copper(I) dimer, **271**, 217

synthesis of a new series of Nb(IV) β-diketonates with the $Nb_2S_4^{4+}$ core: crystal structure of $Nb_2S_4(CF_3COCHCOCH_3)_4$, **271**, 222

synthesis of the first sulfido-bridged octahedral rhenium(III) aqua ion $[Re_6S_8(H_2O)_6]$, **271**, 228

crystal structure and ^{63}Cu NQR of bis[1-aza-2-(1-pyrrolidinyl)]-copper(I) chloride and bromide, **271**, 231

an unusual cyanide-bridged extended A-frame complex: the molecular structure of $[\{Pd_2Br_2(\mu\text{-}PPh_2)(\mu\text{-}dppm)\}_2(\mu\text{-}CN)_2]$, **271**, 236

synthesis and characterization of copper(I) complexes with triphenylphosphine and heterocyclic thione ligands: the crystal structure of (thiazolidine-2-thione)(bis-triphenylphosphine)copper(I) chloride, **271**, 243

Cu^{II}–herbicide complexes: structure and bioactivity, **272**, 24

metal complexes of Imazapyr, a herbicide provided with efficient metal-chelating ability: crystal structure of the cobalt(III) and manganese(II) complexes, **272**, 68

synthesis and structure of borylated organocobaloximes containing neutral nitrile ligands, **272**, 74

comparison of the neutral benzyldiphenylphosphine complexes $C_6H_6Cr(CO)_2PPh_2Bz$ and $C_5H_5Mn(CO)_2PPh_2Bz$ with the isoelectronic manganiobenzyldiphenylphosphonium salts $[C_5R_5Mn(CO)(NO)PPh_2Bz]BF_4$, **272**, 101

lanthanide metal complexes of *N*,*N*′-ethylenebis(pyridin-2-one); from chelates to metallacyclic frameworks, **272**, 131

rhenium(VII) and (V) imido and oxo complexes containing 1,4,7-triazacyclononane: synthesis, electrochemistry and crystal structures, **272**, 176

mono- and bis-alkynyl ruthenium(II) complexes containing the ferrocenyl moiety; crystal structure of *trans*-$[Ru(C{\equiv}CC_5H_4FeC_5H_5)_2(Ph_2PCH_2CH_2PPh_2)_2]$ and electrochemical studies, **272**, 188

transition metal complexes with sulfur ligands. Part CXXV. Synthesis and characterization of hydrido and chloro complexes with rhodium sulfur cores, **272**, 211

functionalized acyclic Schiff bases and related complexes with d- and f-metal ions, **272**, 235

technetium and rhenium carbonyl complexes with 1,4-benzodiazepine derivatives: X-ray crystal structure, **272**, 267

Ni(II) and Cu(II) Schiff base complexes with an extended H-bond network, **272**, 283
chelate ring-opening ruthenium complexes: X-ray crystal structure and solution studies of *cis,trans*-bis(2-dimethyl-aminoethyl)diphenylphosphino(dichloro)ruthenium(II), **273**, 1
a new class of biomimetically relevant 'Scorpionate' ligands. III. The bis(pyrazolyl)methane(phen-2′-ol)s: synthesis and structural characterization of mono and dinuclear copper(II) complexes, **273**, 14
oligonuclear ferrocenolato and 1,1′-ferrocenediolato derivatives of phosphorus: synthesis, structure and electrochemical behaviour, **273**, 24
synthesis, crystal structure and magnetic properties of bis[μ-bis(3-(pyridin-2-yl)-1,2,4-triazole-N^1,$N1$,$N2$)] bis[triaqua nickel(II)] tetranitrate, **273**, 54
rhodium(III) analogues of antitumour-active ruthenium(III) compounds: the crystal structure of [ImH][*trans*-$RhCl_4(Im)_2$] (Im = imidazole), **273**, 62
oxidative addition of 2,3,4-benzotrithiepin 1-oxide to platinum(0) compounds: a convenient synthetic route to platinum(II) thiosulfinato complexes, **273**, 106
crystalline dihydrogen complexes. Intramolecular and intermolecular interactions and dynamic behavior, **273**, 116
synthesis and coordinating properties of the functionalized macrocyclic ligand 1,4-bis(1-methylimidazol-2-ylmethyl)-7-carboxymethyl-1,4,7-triazacyclononane. Crystal structures of cobalt(III), nickel(II) and zinc(II) complexes, **273**, 151
X-ray structures of apo and tungstate derivatives of vanadium chloroperoxidase from the fungus *Curvularia inaequalis*, **273**, 160
novel oxidation and reduction products of the neutral nickel-dithiolene [$Ni(Pr^i_2timdt)_2$] (Pr^i_2timdt is the monoanion of 1,3-diisopropylimidazolidine-2,4,5-trithione), **273**, 175
a reversible Cu^{II}/Cu^{I} electron transfer centre based on a flexible di-imidazole ligand, **273**, 225
syntheses of ruthenium hydridotris(1-pyrazoryl)borate complexes having sulfur-donor coligands, **273**, 238
synthesis and characterisation of rhodium(I) complexes containing the dihydrobis(pyrazolyl)borate ligands $[H_2B(pz')_2]^-$. The X-ray structures of $Bp^*Rh(C_2H_4)_2$ and $BpRh(PPh_3)_2$ ($Bp = H_2B(pz)_2$; $Bp^* = H_2B(3,5\text{-}Me_2\text{-}pz)_2$, **273**, 244
solvothermal synthesis and crystal structure of [La(ethylenediamine)$_4$Cl]In_2Te_4: A 1-D indium telluride, **273**, 255
synthesis, characterization and crystal structure of *trans*-aquatrichlorobis(5,7-dimethyl[1,2,4]triazolo[1,5-*a*]pyrimidine-N^3)-ruthenium(III) monohydrate, **273**, 259
synthesis of chloro and methyl imido cyclopentadienyl molybdenum and tungsten complexes. X-ray molecular structures of [WCp^*Cl_3(N^tBu)], [$MoCp^*ClMe_2$(N^tBu)] and [WCp^*ClMe_2(N^tBu)], **273**, 270
synthesis, structure characterization and magnetic properties of tellurostannates [$M(en)_3]_2(Sn_2Te_6)$ (M = Mn, Zn), **273**, 310
metal complexes of $P(Ph_2)CH_2(Ph_2)PSe$. Crystal structure of [$Ni\{P(Ph_2)CH_2(Ph_2)PSe\}_2]Cl_2 \cdot 2EtOH$, **273**, 320
copper complexes modelling the interaction between benzolamide and Cu-substituted carbonic anhydrase. Crystal structure of Cu(bz)-$(NH_3)_4$ complex, **273**, 334
macrocyclic ligand design: structure–function relationships involving the interaction of cobalt(II), nickel(II) and copper(II) with mixed donor macrocyclic ligands, **273**, 372
unexpected assembly of a (μ-oxo)(μ-formato)diiron(III) complex from an aerobic methanolic solution of Fe(III) and the TPA ligand, **273**, 393
spin crossover in six-coordinate [$Fe(L)_2(NCX)_2$] compounds with L = DPQ = 2,3-bis-(2′-pyridyl)-quinoxaline, ABPT = 4-amino-3,5-bis(pyridin-2-yl)-1,2,4-triazole and X = S, Se: synthesis, magnetic properties and single crystal studies, **274**, 1
quadruply bonded complexes containing η^2- or η^3-tridentate phosphine ligands; syntheses, structures and $^{31}P\{^1H\}$ NMR spectra of $Mo_2Cl_4(\eta^3\text{-etp})(L)$ and $Mo_2Cl_4(\eta^2\text{-etp})_2$ (L = CH_3OH, PMe_3 and PEt_3; etp = $Ph_2PCH_2CH_2P(Ph)CH_2CH_2PPh_2$), **274**, 7
oxidative addition of methyl iodide and iodine to new binuclear rhodium(I) and iridium(I) compounds containing diaminoanthraquinonate-bridging ligands. Crystal structure of [Rh_2(μ-1,4-DA)-$(CO)_2(PPh_3)_2$] (1,4-H_2DA = 1,4-diaminoanthraquinone), **274**, 15
synthesis and properties of a new tetraaza macrocyclic ligand containing one N-acetic acid pendant arm and its copper(II) and nickel(II) complexes, **274**, 24
structural and electronic studies of the coordination of 6-(thiazol-2-yl)-2,2′-bipyridine and related systems to Fe(II), Co(II) and Ni(II), **274**, 32
ternary copper(II) complexes with the versatile 4,7-dihydro-5-methyl-7-oxo-[1,2,4]triazolo[1,5-*a*]pyrimidine ligand, **274**, 53
tris(4,4′-dimethoxy-2,2′-bipyridine)osmium(II), amperometric properties and crystal structure, **274**, 64
structural characterization of an eight-coordinate Nb(IV)–edta complex, [$Nb(edta)(H_2O)_2] \cdot 2H_2O$, **274**, 103
exchange coupling across multiple-atom bridges: crystal structure and magnetic properties of a chloranilate bridged dicopper(II) complex, **274**, 111
crystal structures of aryloxide complexes of zinc embracing sodium aryl group interactions: Na[$Zn(2,6\text{-diphenylphenoxide})_3(H_2O)$] and Na[$Zn_2(2,6\text{-diisopropylphenoxide})_4Cl] \cdot 3THF$, **274**, 115
different picrate coordination modes along the lanthanide series: synthesis and molecular structure of two isomorphous compounds [$Ln_2(pic)_4(TDTD)_3(H_2O)_2](pic)_2(H_2O)_2$ (Ln = Tm, Lu; pic = picrate; TDTD = *trans*-1,4-dithiane-*S*,*S*′-dioxide), **274**, 122
electrochemical synthesis and characterization of indium pyrimidine-2-thiolate complexes. Molecular structure of tris(pyrimidine-2-thiolato)indium(III) and tris[(5-ethyl-4,6-dimethyl)pyrimidine-2-thiolato]indium(III), **274**, 131
the synthesis, NMR and X-ray analysis of tris(1-ethyl-2-isopropylimidazol-5-yl)phosphine gold(I)-2,3,4,6-tetra-O-acetyl-1-thio-β-D-glucopyranoside, a bifunctional auranofin analogue, **274**, 137
nickel(II), palladium(II), platinum(II) and platinum(IV) complexes of *cis*-1,2-bis(diphenylphosphino)ethene and nitrogen-containing heterocycles or sulfur ligands, **274**, 143
binuclear oxo-bridged iron(III) complexes of chiral tetradentate ligands, **274**, 155
synthesis, structure and properties of lanthanide dithionates and their triphenylphosphine oxide complexes, **274**, 177
photosubstitution reactivity, crystal structures, and electrochemistry of ruthenium(II)(III) complexes containing tetradentate (O_2N_2, S_2N_2, and P_2N_2) Schiff base ligands, **274**, 184
cobalt(III) complexes of crown thioethers: the crystal structure of (1,4,7,11,14,17-hexathiacycloeicosane)cobalt(III) tetrafluoroborate dihydrate, **274**, 192
preparation and X-ray crystal structures of $Li_4[Hg_2(edta)_2] \cdot 8H_2O$ and $Ba_2[Hg_2(teta)_2] \cdot 9H_2O$ (teta = *N*,*N*,*N*′,*N*′-diaminobutanetetraacetate), **274**, 219
synthesis and crystal structure of a seven-coordinate dinuclear iron(III) complex containing only bidentate ligands, **274**, 224
polynuclear cobalt cluster compounds formed via sulfur–carbon bond cleavage of 2,2-dicyano-1,1-ethylenedithiolate, **274**, 229
trimethylsilyl iodide as a halide exchange reagent in inorganic chemistry: synthesis and structure of $Cp^*_2ThI_2$ ($Cp^* = \eta^5\text{-}C_5Me_5$), **274**, 232
synthesis and crystal structure of a new polynuclear lanthanum picrate complex with *trans*-1,4-dithiane-1,4-dioxide, **274**, 236
synthesis and molecular structure of the molybdate-bridged cyclic imido complex [$Mo(S_2CNEt_2)_2(NR)(\mu\text{-}MoO_4)]_2$ (R = 2,6-$^iPr_2C_6H_3$), **274**, 243
novel trigonal-bipyramidal copper(II) complexes containing one discrete amino acid chelate: crystal structures of [Cu(L-val)-

(pmdt)]ClO_3 and [Cu(aiba)(pmdt)]$ClO_4 \cdot H_2O$ (L-val = L-valinate ion, aiba = α-aminoisobutyrate ion, and pmdt = *N,N,N′,N″,N″*-pentamethyldiethylenetriamine), **274**, 247

complexes of niobium and tantalum with chloride, sulfide and oxide ligands: synthesis and crystal structure of the (tetrachloro)bis-(η^2-disulfido)(μ-oxo)(μ-η^2:η^2-disulfido) bismetallates $[Cl_2(S_2)Nb(\mu\text{-}O)(\mu\text{-}S_2)Nb(S_2)Cl_2]^{2-}$ and $[Cl_2(S_2)Ta(\mu\text{-}O)(\mu\text{-}S_2)Ta(S_2)Cl_2]^{2-}$ as their tetraphenylphosphonium salts, **274**, 251

crystal structure, EPR and magnetic susceptibility studies of tetrakis[μ-(β-alanine)-*O,O′*]dichlorodicopper(II) dichloromonohydrate, **275–276**, 28

the structure of the azide coordinated superoxide dismutase of *Propionibacterium shermanii* investigated by X-ray structure analysis, extended X-ray absorption fine structure, Mössbauer and electron paramagnetic resonance spectroscopy, **275–276**, 65

hydrothermal synthesis and structural characterization of the three-dimensional framework solids $(NH_4)[V_2(OH)(PO_4)_2(H_2O)] \cdot H_2O$ and $Rb_6[(Mo_9V_3O_6)(PO_4)_{10}(H_2PO_4)_3(OH)_9] \cdot 8.5H_2O$, **275–276**, 122

binding properties and crystal structures of azamacrocycles containing nitrophenol side arms, **275–276**, 168

nucleophilic attack within Ge, Sn and Pb complexes containing $Me_2N(CH_2)_3$– as a potential intramolecular donor ligand, **275–276**, 203

X-ray crystallographic and absorption spectroscopic analyses of structures of catecholato(pyridine)iron chloride complexes in relevance to functional model complexes for catechol 1,2-dioxygenases, **275–276**, 222

structural preferences of η^4-dienecyclopentadienyl complexes: molecular mechanics, molecular orbital and crystallographic studies, **275–276**, 263

a ternary complex of carbonic anhydrase: X-ray crystallographic structure of the adduct of human carbonic anhydrase II with the activator phenylalanine and the inhibitor azide, **275–276**, 295

crystal and molecular structure of $[Ru(bpy)_2(2,3\text{-}dpp)]Cl_2 \cdot 3H_2O \cdot CH_3CN$ and ^1H and ^{99}Ru NMR spectra of $[Ru(bpy)_2(2,n\text{-}dpp)][PF_6]_2$ (bpy = 2,2′-bipyridine, dpp = bis(2-pyridyl)-pyrazine, *n* = 3 or 5), **275–276**, 320

synthesis, X-ray structure and bonding of tris(2,2-6,6-tetramethylheptane-3,5-dionato) bismuth(III), **275–276**, 340

4′-(4-fluorophenyl)-2,2′:6′,2″-terpyridine – an aryl-2,2′:6′,2″-terpyridine with an electron-releasing aryl substituent, **275–276**, 359

copper(I) and copper(II) complexes of polydentate phosphinoamines, **275–276**, 401

boat versus chair conformation in *N*-methyl- and *N,N′*-dimethylpiperazine platinum(II) complexes studied by X-ray analysis. A rare example of metal chelate piperazine: *cis*-$[PtCl_2(Me_2ppz)]$. Part I, **275–276**, 410

synthesis and molecular structure of *N*-methylpiperazinium(+1) and *N,N′*-dimethylpiperazinium(+2) platinum(II) and platinum(IV) complexes. Part II, **275–276**, 419

syntheses and structures of some Ru_2^{5+} complexes containing weakly coordinating anions as axial ligands, **275–276**, 447

five-coordinate platinum(II) alkyne complexes: synthesis, ab initio calculations and crystal and molecular structure of $[PtI_2(Me_2phen)(\eta^2\text{-}PhC{\equiv}CPh)] \cdot CHCl_3$, **275–276**, 500

three reactions of Ru_2^{5+} compounds of the paddlewheel type that lead to cleavage of the Ru–Ru bond, **275–276**, 541

synthesis and X-ray crystal structure of an oxorhenium(V) complex with a double Schiff base, **275–276**, 551

syntheses, properties and structural characteristics of several new (polypyrazolylborato)samarium(III) complexes: agostic interaction of the BH_2 proton with samarium, **277**, 8

complexes of heterocyclic thiones and group twelve metals. Part 1. Preparation and characterisation of 1:1 complexes of mercury(II) halides with 1-methylimidazoline-2(3H)-thione: the crystal structure of [(μ-dibromo)bis(*trans*{(bromo)(1-methyl-imidazoline-2(3H)-thione)}mercury(II))] at 160 K, **277**, 14

crystal structure and properties of a new five-coordinate manganese superoxide dismutase mimic, **277**, 21

structural study and electron exchange kinetics of the bis(2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline)copper(II)/(I) couple in acetonitrile, **277**, 31

synthetic, spectroscopic, and X-ray crystallographic studies on the uranyl(VI) complexes of the β-diketophenol ligands 1-(2-hydroxyphenyl)-1,3-butanedione and 1-(2-hydroxyphenyl)-3-phenyl-1,3-propanedione, **277**, 37

hydrothermal synthesis and characterization of mixed-valence hexatungstates: crystal structures of $[(C_3H_5)_4N]_3[W^{V}W^{VI}_5O_{19}] \cdot 0.5H_2O$ and $[H_3N(CH_2)_2NH_3]_2[W^{V}W^{VI}_5O_{19}] \cdot [H_2N(CH_2)_2NH_2]Cl \cdot 8H_2O$, **277**, 69

manifestation of redox duality of 2-propanone oxime: Pt(II)-assisted reduction versus Pt(IV)-mediated oxidation of $Me_2C{=}NOH$ species, **277**, 83

electrophilic substitution of metal-coordinated pyrazole: nitration, sulfonation and bromination of $[Co(NH_3)_5(pyzH)]^{3+}$ (pyzH = pyrazole), **277**, 89

synthesis and structural studies on metal nitrate complexes with 1,3-dimethyllumazine and 1,3,6,7-tetramethyllumazine: crystal structure of two new cobalt(II) and copper(II) three-dimensionally hydrogen-bonded complexes and a cadmium(II) complex with unusual geometry, **277**, 103

2-(diphenylphosphino)pyridine (dppy) as monodentate or bridging ligand in homoleptic silver(I) complexes. Crystal structures of $[Ag(\eta^1\text{-}dppy)_4][ClO_4]$ and $[Ag_2(\eta^1\text{-}dppy)(\mu\text{-}dppy)_2][ClO_4]_2$, **277**, 111

palladium(II) and platinum(II) complexes containing new phosphinoxy chiral P,N-ligands, **277**, 119

synthesis and characterization of a new unsymmetrical μ-oxo-bridged iron(III) complex with triphenylarsine oxide ligands, **277**, 127

synthesis and characterization of the tungsten–gallium heterometallic cluster $Na_2[W_3GaO_4(O_2CEt)_8]_2$, **277**, 130

crystal structure and characterization of binuclear copper(II) complex with *N*-(2-hydroxyphenyl)-3-methoxycarbonyl-4,6-dimethylsalicylamide, **277**, 134

Anderson–Evans type molybdotellurates with diazolium cations: preparation, structure and properties of $(H_2imz)_6[TeMo_6O_{24}] \cdot 4H_2O$ and $(H_2pyz)_6[TeMo_6O_{24}] \cdot Te(OH)_6$, **277**, 139

five-coordinate Salen(*t*Bu) complexes of zinc, **277**, 157

variable coordination modes for 1,1′-bis(diphenylphosphino)-ferrocene (dppf) mono- and di-chalcogenides or -oxides (dppfE and dppfE$_2$; E = O, S or Se) in copper(I) complexes. Crystal structure of $[\{Cu(dppf)\}_2(\mu\text{-}dppfS_2)](BF_4)_2$ and $[Cu(dppfSe)_2]BF_4$, **277**, 163

polynuclear silver compound formed from aggregation of Se^{2-} and Ag(I)-thiolate complex. Synthesis, structure and spectroscopic characterization of $Ag_{11}(\mu_5\text{-}Se)(\mu_4\text{-}Et_2NCS_2)_3(\mu_3\text{-}Et_2NCS_2)_6$, **277**, 171

bis(2-aminomethylpyridine) complexes of Cr(III) and Co(III), **277**, 177

reaction of the dinuclear cobalt complex $[Co_2\{\mu\text{-}PPh_2CHCHC(O)CHCHPPh_2\}(CO)_4]$ with two-electron donor ligands; the stabilisation of cobalt–alkene bonds by incorporation of alkenes into a bridging diphosphine ligand, **277**, 186

ligand exchange reactions of $ReCl_4(PPh_3)_2$, $ReOCl_3(PPh_3)_2$ and $NiCl_2(PPh_3)_2$ with acetylacetone thiobenzoylhydrazone. An unexpected template reaction at the ReO^{3+} core. Molecular structures of the resulting oxorhenium(V) and nickel complexes, **277**, 219

pentamethylphenylcopper(I): a square-planar tetranuclear cluster, **277**, 232

trans influence of triphenylstibine. Crystal and molecular structures of *cis*-$[PtCl_2(SbPh_3)_2]$ and *trans*-$[PtI_2(SbPh_3)_2]$, **277**, 236

crystal structure and molecular stereochemistry of the 1:2 adduct of anionic complex, dicyano-(5,10,15,20-tetramesityleneporphinato)-iron(II) with bis(triphenylphosphine)nitrogen (1+)·I_2, $CDCl_3$ solvate {[$(CN)_2Fe^{II}$(TMP)][PPN]$_2$}·I_2·2$CDCl_3$, **277**, 242

crystal structure and fluxional behavior of novel Pd(II) allyl complexes containing a potentially tridentate S,N,S ligand: an example of selective *syn–anti* isomerism, **277**, 247

synthesis of a dinuclear, phosphido-bridged complex of rhenium, X-ray crystal structure of $Re_2(\mu\text{-H})[\mu\text{-(t-Bu)}PC_6H_4P(H)\text{t-Bu}](CO)_7$, **277**, 253

synthesis and catalytic activity of manganese(II) complexes of pentanedioic acid; X-ray crystal structure of $[Mn(phen)_2(H_2O)_2][Mn(O_2C(CH_2)_3CO_2)(phen)_2H_2O](O_2C(CH_2)_3CO_2)\cdot 12H_2O$ (phen = 1,10-phenanthroline), **277**, 257

a new form of decavanadate: hydrothermal synthesis and structural characterization of $Ba_{5.33}(H_3O)_{0.33}[V_{10}O_{30}]$, **277**, 263

a series of $[Cr_2(\mu\text{-OH})_n(\mu\text{-RCO}_2)_{3-n}(bispma)_2]$ type chromium(III) dinuclear complexes ($n = 1, 2$) with tridentate bis(2-pyridylmethyl)amine (bispma): synthesis, crystal structure, and magnetic properties, **278**, 15

kinetic studies using Co(III) and Cr(III) complexes of 1,4,7,11-tetraazaundecane (2,2,3-tet) and the crystal structure of *trans*-$[CrCl_2(2,2,3\text{-tetH})(OH_2)]Cl\cdot ClO_4$, **278**, 24

synthesis, structural characterization and magnetic behaviour of two new $(\mu_{1,3}\text{-}N_3)_2$ dinuclear nickel compounds: study of the magnetic transition at 235 K for the *cis*-$(\mu_{1,3}\text{-}N_3)_2[Ni(dl\text{-cth})]_2(ClO_4)_2$ compound, **278**, 43

some hexadentate Ni(II)-edta-type complexes containing five-membered diamine rings. The molecular and crystal structure of the *trans*(O_5) isomer of barium(ethylenediamine-*N*,*N*′-diacetato-*N*,*N*′-di-3-propionato)nickelate(II) hexahydrate, *trans*(O_5)-Ba[Ni(eddadp)]·6H_2O, and strain analysis of edta-type chelates in relation to their octahedral distortion, **278**, 66

the reactivity of the $[Re{\equiv}O]^{3+}$ core toward aromatic 1,2-diamines: the formation of rhenium(V)-imido complexes, **278**, 96

synthesis and crystal structure of a new salt of the complex anion $[(TCNQ)_3]^{2-}$ $[Mo_2(\mu_2\text{-}S_2)_2(S_2CNMe_2)_4](TCNQ)_3$, **278**, 108

bis(1,2-diselenooxalato)nickelate(II) and -cuprate(II) — the first diselenooxalate complexes, **278**, 118

spectroscopic and magnetic properties and structure of a five-coordinate, O_2-binding cobalt(II) Schiff base complex and of the copper(II) analogue, **278**, 127

the chemistry of dinitrato-2,2′-bipyridinecopper(II): preparation and characterization of binuclear complexes having a $Cu(\mu\text{-OH})_2Cu$ and $Cu(\mu\text{-}N_3)_2Cu$ core, **278**, 143

spectroscopic and magnetic properties of the pyridine-2-carbaldehyde thiosemicarbazonateiron(III) complexes: $[Fe(C_7H_7N_4S)_2]X\cdot nH_2O$ (X = Cl, ClO_4, NO_3, PF_6). Crystal structure of the hexafluorophosphate compound, **278**, 150

complexes of a new bidentate chelating pyridyl/sulfonamide ligand with copper(II), cobalt(II) and palladium(II): crystal structures and spectroscopic properties, **278**, 178

synthesis and characterization of a neutral, low spin iron(III) complex of a hexadentate tripodal ligand containing three imidazolate arms. Use as a dinucleating agent, **278**, 197

solvent-induced reaction of rhodium(III) halide with N-heterocyclic ligands. Crystal structure of *trans*-dichlorotetrakis(*N*-methylimidazole)rhodium(III) chloride dihydrate, **278**, 209

the 1,8-bis(2′-pyridyl)-3,6-dithiaoctane complex of nickel(II): X-ray crystal structure and borohydride adduct formation, **278**, 217

silver(I)–silver(I) interactions in the tetrameric silver thiolate phosphine complex $[Ag_4(SPh)_4(PPh_3)_4]$, **278**, 229

reactions of $Ti(O\text{–i-Pr})Cl_3$ with 2-aminophenol and the crystal structure of the zwitterionic complex $[Ti(O\text{–i-Pr})Cl_3(THF)(^-OC_6H_4NH_3^+)]\cdot THF$, **278**, 232

synthesis of new copper(II) complexes of polydentate ligands containing pyridyl pendant arms. Crystal structure of [*N*,*N*,*N*′,*N*′-tetrakis(2-pyrid-2-yl-ethyl)1,2-ethylenediaminecopper(II)]-bis(hexafluorophosphate)·1.5 acetone, **278**, 237

the synthesis, characterization and X-ray crystal structures of the *fac-cis* and *mer-trans* isomers of $Re(NEt)Cl_3(PMe_3)_2$, **278**, 245

$(C_3H_7N_2O)Cu_3Br_4$: template synthesis of a two-dimensional copper(I) bromide lattice, **279**, 1

protonation of Fischer-type alkylidyne carbonyltungsten complexes. Structural comparison of alkylidyne and alkylidene metal complexes, including a neutron diffraction study of $[W(CHCH_3)Cl_2(CO)(PMe_3)_2]$, **279**, 7

syntheses and X-ray crystal structures of *cis*-$[Mn(bipy)_2Cl_2]\cdot 2H_2O\cdot EtOH$ and *cis*-$[Mn(phen)_2Cl_2]$ (bipy = 2,2′-bipyridine; phen = 1,10-phenanthroline); catalysts for the disproportionation of hydrogen peroxide, **279**, 24

synthesis of tetracarboxylatediruthenium(II) compounds: reduction and disproportionation reactions of diruthenium(II,III) units. Crystal structure of $Ru_2(\mu\text{-}O_2CCOPh)_4(thf)_2$, **279**, 30

structural information on *trans*-1,2-diaminocyclohexane-*N*,*N*′-tetraacetateferrate(III) in the solid and aqueous phase, **279**, 37

Lewis base adducts of halogenorhenium(VII) oxides: ^{17}O NMR spectroscopy, structural aspects and catalysis, **279**, 44

intercalator-linked cisplatin: synthesis and antitumor activity of *cis*-dichloroplatinum(II) complexes connected to acridine and phenylquinolines by one methylene chain, **279**, 51

exchange interactions through hydrogen-bond bridges. Crystal structure, spectroscopic characterisation and magnetic properties of the complex $[\{Cu(en)\}_2(\mu\text{-egta})]\cdot 4H_2O$ (H_4egta = 3,12-bis(carboxymethyl)-6,9-dioxa-3,12-diazatetradecanedioic acid), **279**, 58

the synthesis, molecular structure and fluxional properties of the 1-phenylpropyne complex $[WI_2(CO)\{Ph_2P(CH_2)PPh_2\}(\eta^2\text{-}MeC_2Ph)]$, **279**, 65

transition metal complexes with nitronyl nitroxides. Part 1. Crystal structure and properties of square-planar *trans* radical–copper(II)–radical complexes, **279**, 69

transition metal complexes with nitronyl nitroxides. Part 2. Crystal structure and magnetism of manganese(II) compounds including radicals and their reduced derivatives, **279**, 76

some chlorocarbonylruthenium(II) complexes of P,N-donor ligands: crystal structures of $[RuCl(CO)\{PPh_2CH_2C(Bu^t){=}NNH_2\}_2]Cl$ and *fac*,*cis*-$[RuCl_2(CO)\{PPh_2CH_2C(Bu^t){=}N\text{–}N{=}C(Bu^t)\text{-}CH_2PPh_2\}]$, **279**, 95

a phosphane analog of nedaplatin: preparation and structural characterization of *cis*-bis(trimethylphosphane)(glycolato-*O*,*O*′)-platinum(II), **279**, 105

synthesis and structural characterization of metal complexes of an unsymmetric N_3O tetradentate Schiff base ligand, **279**, 111

synthesis and structural properties of (diamine)platinum(II) complexes of itaconic acid, **279**, 116

synthesis, crystal structures and magnetic properties of $[Fe_2(bpym)(C_5O_5)_2(H_2O)_4]\cdot 2H_2O$ and two polymorphs of $[Fe_2(bpym)(C_4O_4)_2(H_2O)_6]\cdot 2H_2O$ (bpym = 2,2′-bipyrimidine), **279**, 127

synthesis through an in situ esterification process and characterization of oxo isopropoxo titanium clusters, **279**, 144

tris- and tetrakis(1-methylcytosine) complexes of Pt(II): syntheses and X-ray structures of $[Pt(1\text{-MeC-}N3)_3Cl]^+$ and $[Pt(1\text{-MeC-}N3)_4]^{2+}$ compounds, **279**, 152

synthesis and crystal structure of a new isomer of the μ-oxo-bis[dichlorooxobis(pyridine)rhenium(V)] complex $\{ORecl_2py_2\}_2O$, **279**, 159

preparation and properties of $C_8H_4S_8$-platinum(II) complexes and electrical conductivities of their oxidized species and X-ray crystal structure of $C_8H_4S_8(CH_2CH_2CN)_2$ as a pro-ligand compound, **279**, 165

solid-state synthesis of cluster compound $[MoCu_3OS_3(PPh_3)_3(CH_3COO)]$ with its structure, reactivity and optical non-linearities, **279**, 172

synthesis and characterisation of mononuclear and binuclear iron(II) complexes of pentadentate and bis(pentadentate) ligands derived from 1,4,7-triazacyclononane, **279**, 192

the reaction chemistry of a technetium(I) nitrosyl complex with potentially chelating organohydrazines: the X-ray crystal structure of $[TcCl_2(NO)(HN{=}NC_5H_4N)(PPh_3)]$, **279**, 206

synthesis and characterization of the new complex tris[(2-oxazolinyl)phenolato] manganese(III) and its use as a catalyst in oxidation reactions, **279**, 217

synthesis and metal complexes of a chelating ligand containing thiomorpholine and pyridine: 2,6-bis(thiomorpholinomethyl)pyridine, **279**, 221

structure and magnetic properties of a novel chloro-bridged polymeric cadmium(II) complex with pyridyl-substituted nitronyl nitroxide, **279**, 233

synthesis and characterization of nickel(II) complexes of di-*N*-alkylated 14-membered tetraaza macrocycles containing cyclohexane rings via regioselective alkylation reaction, **279**, 238

preparation and properties of ferrocenyl bimetallic compounds for non-linear optics, **279**, 243

molecular structure of a five-coordinate binuclear zinc complex, **279**, 249

cis- and *trans*-titanium complexes with doubly silyl-bridged dicyclopentadienyl ligands: molecular structure of $[(TiCl)_2(\mu\text{-}O)\{(SiMe_2)_2(\eta^5\text{-}C_5H_3)_2\}]_2(\mu\text{-}O)_2$, **280**, 1

transition metal complexes with sulfur ligands. Part CXXIX. Retention and reactivity of the $[Fe{\backsim}NH{\backsim}NH{\backsim}Fe]$ chromophore in the iron sulfur diazene complex $[\mu\text{-}N_2H_2\{Fe(PPr_3)('S_4')\}_2]$ in exchange and oxidation processes. ($'S_4'^{2-}$ = 1,2-bis(2-mercaptophenylthio)ethane(2−)), **280**, 39

the *trans* influence of F, Cl, Br and I ligands in a series of square-planar Pd(II) complexes: relative affinities of halide anions for the metal centre in *trans*-$[(Ph_3P)_2Pd(Ph)X]^+$, **280**, 87

tetravalent niobium alkyls and enolate stabilized by anionic organic amides, **280**, 143

synthesis and characterization of homologous nickel(II) and palladium(II) complexes with biphosphine monoxide ligands $Ph_2P(CH_2)_nP(O)Ph_2$ (n = 1–3) and $pTol_2P(CH_2)P(O)pTol_2$, **280**, 150

reactivities of the coordinated organonitriles in molybdenum(0) and tungsten(0) phosphine complexes: protonation of the nitrile carbon and cleavage of the C≡N triple bond, **280**, 163

preparation, structure and properties of triphenylphosphine rhodium aryloxides, **280**, 172

preparation of doubly-silylene-bridged zirconocene alkyl complexes, $(Me_2Si)_2\{\eta^5\text{-}C_5H_2\text{-}4\text{-}CHMe_2\}\{\eta^5\text{-}C_5H\text{-}3,5\text{-}(CHMe_2)_2\}ZrR_2$ ($R = CH_3$, CH_2Ph) and investigations of their activity in 1-pentene polymerization. Molecular structure of $(Me_2Si)_2\{\eta^5\text{-}C_5H_2\text{-}4\text{-}CHMe_2\}\{\eta^5\text{-}C_5H\text{-}3,5\text{-}(CHMe_2)_2\}Zr(CH_2Ph)_2$, **280**, 226

synthesis, solution behavior and molecular structure of 16-electron *closo*-2-(Ph_3P)-1-N,2-$[\mu\text{-}(\eta^2\text{-}CH_2CH{=}CH_2)]$-1-N-$(\sigma\text{-}CH_2CH{=}CH_2)$-2,1-$RhCB_{10}H_{10}$, the first η^2-olefinic monocarbon metallacarborane, **280**, 233

binuclear and tetranuclear complexes prepared by reactions of dodecacarbonyltriruthenium with the N-substituted cinnamic acid amides. Crystal and molecular structures of $[Ru_2(\mu\text{-}H)(\mu\text{-}\eta^3\text{-}PhC{=}CHC(O)N(CH_2)_4)(CO)_6]$ and $[Ru_4(\mu_3\text{-}H)_2(\mu\text{-}\eta^3\text{-}PhC{=}CHC(O)N(CH_2)_4)_2(CO)_{10}]$, **280**, 243

synthesis and properties of the dichloro[tetramethyl(trifluoromethyl)cyclopentadienyl]ruthenium dimer: X-ray structure of $[M_2(\eta^5\text{-}C_5Me_3CF_3)_2Cl_2(\mu\text{-}Cl)_2]$ (M = Ru, Rh), **280**, 249

on the reduction behaviour of 2-methoxyethyl and 2-methylthioethyl substituted zirconocene dichlorides: crystal structure of $(\eta^5,\sigma\text{-}C_5Me_4CH_2CH_2S)_2Zr$, **280**, 257

chiroptical properties of heterotrimetallic 'wing-tip' butterflies; synthesis and crystal structure of $(\mu\text{-}H)Ru_2Fe_2PdC(CO)_{12}(\eta^3\text{-}\beta\text{-}C_{10}H_{15})$, **280**, 275

bimetallic cyclooligosiloxanolate complexes of copper and nickel, **280**, 282

molecular structure and reactions of azobenzene complexes with iron–lithium compounds, **280**, 288

syntheses and properties of organocyanamide, cyanoguanidine and dinitrogen complexes of rhenium. Crystal structure of *mer*-$[ReCl_2(NCNEt_2)(PMePh_2)_3]$, **280**, 308

synthesis and X-ray crystal determination of four novel lanthanide(III) oxydiacetate complexes, **281**, 18

structural and spectroscopic models of the manganese catalase active site. Isolation and structures of the asymmetric $[(H_2O)Mn^{III}(\mu\text{-}O)(\mu\text{-}O_2CR)_2Mn^{III}(L)]$ ($L = Cr_2O_7^{2-}$, CH_3OH) cores: analogs of a substrate-bound catalase active site intermediate, **281**, 25

cation-π interactions between potassium ions and aromatic rings. Crystal structures of three potassium complexes of calix[6]arene, **281**, 36

synthesis, structure, spectroelectrochemical and theoretical investigations of $[(\eta^5\text{-}Me_5Cp)\overline{Co(\eta^3\text{-}MeC{=}CMe\text{–}CMe{=}CMe)Sn}\{CH(SiMe_3)_2\}_2]$. Formation of a cobalt complexed stannacycle via a mixed alkyne–stannylene cycloaddition reaction, **281**, 53

binuclear diorganophosphinothioformamido complexes of gold(I): synthesis and crystal structure of $[Au(Ph_2PC(S)NPh)]_2$ and related systems, **281**, 64

electrochemical synthesis and characterisation of gallium, indium and thallium compounds with 1-hydroxopyridine-2-thione. The crystal structures of tris(1-oxopyridine-2-thionato)indium(III) and tris(1-oxopyridine-2-thionato)thallium(III), **281**, 70

syntheses and crystal structures of geometrical isomeric pairs: *trans*- and *cis*-$(PPh_4)[Co(acac)_2(CN)_2]$ and *trans*- and *cis*-$[Co(acac)_2(PMe_3 \text{ or } PEt_3)_2]PF_6$ (acac = pentane-2,4-dionate), **281**, 77

synthesis of polynuclear copper(II) complexes by controlling the conformation (*cis* or *trans*) of *N,N'*-bis(3-aminopropyl)oxamidate and the coordination mode of 2,2′-bipyrimidine, **281**, 95

crystal structure, spectral and magnetic properties of a new (μ-acetate)(μ-alkoxide) dicopper(II) complex as a model for tyrosinase, **281**, 111

synthesis and crystal structure of dinuclear palladium(I) dppm complex with 4-mercaptopyridine in the thione mode. $[Pd_2(\mu\text{-}dppm)_2(HpyS)_2](ClO_4)_2 \cdot CH_3OH$, **281**, 116

silver(I) complexes with heterocyclic thiones and tertiary phosphines as ligands. Part 3. Solution behaviour of mononuclear complexes of silver(I) nitrate. The crystal structure of bis(triphenylphosphino)bis(benzoxazoline-2-thione)silver(I) nitrate, **281**, 121

chain-like crystal structure and magnetic properties of [catena-bis(1,2-diaminoethane)copper(II)-μ-dicyano-argentate]-dicyanoargentate, **281**, 134

reactivity of $[NBu_4]_2[\{(C_6X_5)_2M(\mu\text{-}OH)\}_2]$ (X = F or Cl; M = Pd or Pt) towards heterocyclic thiones: crystal structure of $[NBu_4]_2[\{(C_6F_5)_2Pd(\mu\text{-}\eta^2\text{-}LL')\}_2]$ (LL′ = methylimidazole-2-thiolate), **281**, 165

complexes of the bis(trimethylsilyl)-benzamidinato ligand '*siam*': synthesis and X-ray structures of $(siam)_2M$, $(siam)(siamH)MX$ (M = Ni, Pd), $(siam)_2MnI$ and $(siam)ReO_3$, and their reactivity towards CO_2, **281**, 181

mixed valence, tetradisordered structure of iodine-doped bismuth(III) diphthalocyanine, **281**, 195

nephelauxetic series of in-plane ligands for square-pyramidal nitridochromium(V) complexes as studied by means of ESR spectroscopy, **281**, 207

synthesis of manganese tricarbonyl cationic complexes of ferrocenyl substituted arenes via a manganese tricarbonyl cation transfer reaction, **281**, 229

coordination chemistry of the sterically hindered N_3-donor hydrotris(3,5-diphenylpyrazol-1-yl)borate toward silver(I)triorganophosphino compounds. Synthesis, structural and spectroscopic characterization, **282**, 1

investigations into the engineering of inorganic/organic solids: hydrothermal synthesis and crystal structures of aromatic ammonium sodium hydrogen phosphomolybdates(VI), **282**, 10

syntheses, spectroscopic characterization and X-ray crystal structures of $[HP\text{-}i\text{-}Pr_3]_2[IrCl_6]$ (**1**), $[HP\text{-}i\text{-}Pr_3]_2[Ir_2Cl_8(P\text{-}i\text{-}Pr_3)]$ (**2**) and $[HP\text{-}i\text{-}Pr_3][Ir_2Cl_7(P\text{-}i\text{-}Pr_3)_2]$ (**3**): three possible precursors of the complex $Ir(H)(Cl)_2(P\text{-}i\text{-}Pr_3)_2$ (**4**), **282**, 17

the reaction of cobalt powder with tetraiodo{1,2-bis(dibenzylphosphino)ethane} to form 1,2-bis(dibenzylphosphino)-ethane cobalt diiodide, $Co(dBzP_2)I_2$; and the X-ray crystal structure of the diphosphinodioxide complex, $Co\{dBzP_2(O)_2\}I_2$, **282**, 25

crystal and molecular structures of some tri-aryltin isothiocyanates (Ar_3SnNCS; Ar = C_6H_5, *p*-$CH_3C_6H_4$, *o*-$CH_3C_6H_4$, *o*-$CH_3OC_6H_4$) and of the adduct hexamethylphosphoramide(*N*-thiocyanato)-tris(*o*-tolyl)tin, **282**, 30

lanthanide(III) complexes of two oxaazadiamine macrocyclic ligands derived from 2,6-diformylpyridine: the crystal structures of a reduced macrocyclic ligand and the corresponding diprotonated macrocycle, **282**, 42

preparation and characterization of twin-cubane-type clusters with molybdenum–copper–sulfur cores: X-ray structure determination of $[\{(NH_3)_6(H_2O)(OH)Mo_3CuS_4(O)\}_2]Cl_4\cdot 8H_2O$, **282**, 50

synthesis, structure and characterization of zinc(II), copper(II), zinc(II)barium(II) and copper(II)barium(II) complexes of macrocyclic heteronucleating ligands based on isothiosemicarbazide, **282**, 61

the effect of aromatic–aromatic interactions on the crystallization of helical nickel(II) complexes, **282**, 71

synthesis, characterization and crystal structures of complexes of sodium hexafluorophosphate with dibenzo-18-crown-6 and dibenzo-24-crown-8 macrocycles, **282**, 76

crystal structures and magnetic properties of uniform and alternating azido-bridged (2,2′-bipyridyl)copper(II) chains, **282**, 82

electrocrystallization and characterization of phosphonium-based radical anion salts $[MePh_3P][Pd(dmit)_2]_3$ and $[Ph_4P][Pd(dmit)_2]_3$ (dmit = 1,3-dithiole-2-thione-4,5-dithiolate). Comparison with related Pd and Ni isologues, **282**, 90

the phenomenon of conglomerate crystallization. Part 51. The crystallization behavior of nickel dinitro amine complexes. *Trans*-$[Ni(3,2,3\text{-tet})(NO_2)_2]\cdot H_2O$ (**I**), *mer*-[Ni(N5-Me-1,5,9-triazanonane)$(NO_2)_2$ (**II**), *cis*-$[Ni(tren)(NO_2)_2]\cdot H_2O$ (**III**) and *cis*-$[Ni(edda)(H_2O)_2]\cdot H_2O$ (**IV**) and control of the bonding mode of the nitrite ligand by the choice of the amine, **282**, 96

ferrocenyl-substituted imine ligands and their reactivity towards $Fe_2(CO)_9$; crystal structure of Fc–C(H)=C(H)–C(H)=N–C_6H_{11} and (η^3-R^1–C(H)=C(H)–C(H)=N–R^2)$Fe(CO)_3$ (R^1 = C_6H_5, R^2 = Fc; R^1 = Fc, R^2 = C_6H_{11}, R^1 = R^2 = Fc; Fc = (η^5-C_5H_5)Fe(η^5-C_5H_4)), **282**, 111

a bimetallic main group oxide cluster of the oxovanadium borate system: $(H_3NCH_2CH_2NH_3)_4[(VO)_{12}O_4\{B_8O_{17}(OH)_4\}_2\text{-}\{Mn(H_2O)_2\}_2]\cdot H_2O$, **282**, 123

new cyclam-type copper(II) complexes with amide molecular padlock: synthesis, properties and crystal structure, **282**, 142

synthesis and characterization of two novel monooxovanadium(V) complexes with bidentate benzohydroxamate ligand, **282**, 149

the synthesis and characterization of Ni, Pd and Pt maleonitriledithiolate complexes: X-ray crystal structures of the isomorphous Ni, Pd and Pt ($Ph_2PCH_2CH_2PPh_2$)M(maleonitriledithiolate) congeners, **282**, 155

pentamethylcyclopentadienyl ruthenium(II) complexes containing chiral diphosphines: synthesis, characterisation and electrochemical behaviour. X-ray structure of (η^5-C_5Me_5)Ru{(*S*,*S*)-$Ph_2PCH(CH_3)CH(CH_3)PPh_2$}Cl, **282**, 163

electrochemical, spectrochemical characterization and crystal structure of (1,4,8,11-tetrazacyclotetradecane)(acetylacetonate)cobalt(III) complexes, **282**, 173

zinc thiolate complexes with chelating nitrogen ligands, **282**, 193

the synthesis and characterization of platinum monodithiolene complexes containing 1,3-dithiole-2-oxo-4,5-dithiolate ($dmid^{2-}$) and 1,3-dithiole-2-thione-4,5-dithiolate ($dmit^{2-}$), **282**, 200

mononuclear, dinuclear, and polynuclear dioxomolybdenum(VI)–tridentate Schiff base complexes: crystal structures and properties, **282**, 209

pentacoordinated cationic rhodium(I) and iridium(I) organocomplexes with α-diimine ligands derived from glyoxal: crystal structure of [Rh(norbornadiene)(glyoxalbis(4-dimethylamino-anil))-$(PPh_3)]BF_4\cdot 1/2CH_2Cl_2$, **282**, 230

synthesis and structure of (1,3-dimethyluracil-5-yl)mercury(II) complexes with aromatic nitrogen donor ligands, **282**, 237

synthesis, crystal structure and magnetic properties of the first single azido-bridged copper(II) chain $[Cu(bpym)(N_3)_2]_n$ (bpym = 2,2′-bipyrimidine), **282**, 252

synthesis, crystal structure and magnetic properties of the one-dimensional chain compound *catena*-poly[aquabis(4,4′-bis-1,2,4-triazole-*N*1)(thiocyanato-*N*)-manganese-μ-[4,4′-bis-1,2,4-triazole-*N*1,*N*1′]thiocyanate], **282**, 257

spectroscopic characterization of coaxially and π–π stacked binuclear copper(II) complexes of ω,ω′-bis(pyridine-2-carboxamido)-alkanes in acidic solution, **283**, 61

dinuclear $Cu^{II}M^{II}$ complexes of a phenol-based macrocycle with $N(amine)_2O_2$ and $N(imine)_2O_2$ metal-binding sites: the effect of chloride ligation upon the site selectivity of metal ions, **283**, 72

mixed ligand copper(II) coordination polymers constructed by Cu–bpm–Cu dimer unit (bpm = 2,2′-bipyrimidine) as a building block. Crystal structures and magnetic properties of $\{[Cu(bpm)(SO_4)]\text{-}(H_2O)\}_n$, $[Cu_2(bpm)(suc)_{0.5}(ClO_4)_2(OH)(H_2O)_2]_n$ and $\{[Cu(bpm)_{1.5}(suc)_{0.5}](ClO_4)(H_2O)_2\}_n$ (suc = succinate), **283**, 80

molecular structure of the methoxy-iron(III) derivative of 5,15-(*o*,*o*′(2-methyl-2′-hydroxy-3,3′-diamidobiphenyl)-diphenylporphyrin and CO binding properties of iron(II)–pyridine complexes of 2,2′-substituted biphenyl strapped porphyrins, **283**, 98

an infinite chiral single-helical structure formed in Cu(II)-L-/D-glutamic acid system, **283**, 105

synthesis, structure and characterization of dinuclear nickel(II) complexes containing the derivatives of *N*,*N*,*N*′,*N*′-tetrakis[(2-benzimidazolyl)methyl]-2-hydroxy-1,3-diaminopropane and an acetate or a propionate ion, **283**, 118

synthesis and characterization of mixed valence μ-alkoxo-diiron(II,III) complexes with an unsymmetric dinucleating ligand, **283**, 124

structures and electrochemical properties of the Co(III) ternary complexes containing NO_3-type tripodal tetradentate ligands and amino acids: effect of the outer coordination sphere on the electrochemical properties, **283**, 160

interactions between $[AuX_4]^-$ (X = Cl, CN) and cytosine and guanine model nucleobases: salt formation with (hemi-) protonated bases, coordination, and oxidative degradation of guanine, **283**, 223

synthesis and crystal structure of a dinuclear platinum(II) bis(*N*,*N*-diethylamino)carbeniumdithiocarboxylate complex $Pt_2\{\mu\text{-}(Et_2N)_2C_2S_2\}_2Cl_4\cdot 0.5CH_2Cl_2$: structural study of the coordinated inner salt, **283**, 256

CT photochemistry

thermal and photochemical reactions of bis(8-quinolinolato)-copper(II), bis(8-quinolinethiolato)copper(II) and (8-quinolinolato,8-quinolinethiolato)copper(II) with chloroalkanes, **273**, 403

Cubane-type complexes

preparation and characterization of twin-cubane-type clusters with molybdenum–copper–sulfur cores: X-ray structure determination of $[\{(NH_3)_6(H_2O)(OH)Mo_3CuS_4(O)\}_2]Cl_4 \cdot 8H_2O$, **282**, 50

synthesis of double incomplete-cubane clusters $[PPh_4]_2[(\eta^5\text{-}C_5Me_5)MS_3Cu_3X_3]_2$ (M = Mo, W; X = Cl, Br) and reactions with PPh_3, dppm (1,2-bis(diphenylphosphino)methane), and $AsPh_3$, **283**, 136

Cuboidal cluster complexes

synthesis and structures of the cuboidal iron–sulfur–nitrosyl–phosphine clusters $[Fe_4S_3(NO)_4(PR_3)_3]^{0,1+}$ (R = Et, Pr^i, C_6H_{11}), **270**, 46

Cyanamide complexes

syntheses and properties of organocyanamide, cyanoguanidine and dinitrogen complexes of rhenium. Crystal structure of *mer*-$[ReCl_2(NCNEt_2)(PMePh_2)_3]$, **280**, 308

Cyanide bridged complexes

the cyanide ligand as an efficient bridge in mixed-valence complexes, **269**, 246

an unusual cyanide-bridged extended A-frame complex: the molecular structure of $[\{Pd_2Br_2(\mu\text{-}PPh_2)(\mu\text{-}dppm)\}_2(\mu\text{-}CN)_2]$, **271**, 236

Cyanide complexes

cyanide binding to phthalocyaninatoiron(II) in dimethyl sulfoxide in the presence of carbon monoxide: kinetic and equilibrium study, **267**, 109

syntheses and crystal structures of geometrical isomeric pairs: *trans*- and *cis*-$(PPh_4)[Co(acac)_2(CN)_2]$ and *trans*- and *cis*-$[Co(acac)_2(PMe_3 \text{ or } PEt_3)_2]PF_6$ (acac = pentane-2,4-dionate), **281**, 77

Cyanide inhibition

proton ENDOR spectroscopy of cyanide-inhibited copper–zinc superoxide dismutase in randomly oriented samples, **273**, 131

Cyano complexes

kinetics and mechanism for reduction of *trans*-dichloro-tetracyanoplatinate(IV) by tetraammineplatinum(II) and bis(ethylenediamine)platinum(II), **282**, 55

Cyanoguanidine complexes

syntheses and properties of organocyanamide, cyanoguanidine and dinitrogen complexes of rhenium. Crystal structure of *mer*-$[ReCl_2(NCNEt_2)(PMePh_2)_3]$, **280**, 308

Cyclam complexes

electrochemical, spectrochemical characterization and crystal structure of (1,4,8,11-tetrazacyclotetradecane)(acetylacetonate)cobalt(III) complexes, **282**, 173

Cyclic phosphite complexes

NMR investigations of di-rhodium-tetraacetate complexes with cyclic phosphites, **267**, 293

Cyclic voltammetry

an electrospray mass spectrometric and voltammetric study of horse heart cytochrome *c* in the presence of metal ions, **267**, 281

nucleophilic attack at the five-coordinate anion $[Mn(CO)_3(3,5\text{-di-tert-butyl-catecholate})]^-$ controlled by electronic and steric effects, **268**, 1

thermodynamics of reduction of *Chromatium vinosum* high-potential iron–sulfur protein and its histidine depleted H42Q mutant, **275–276**, 230

Cyclization reactions

zirconium-catalyzed and zirconium-promoted cyclization reactions of non-conjugated dienes with alkylmagnesium halides to give cycloalkylmethylmagnesium derivatives, **280**, 8

Cycloalkanes

unprecedented facile reactions of propane and cycloalkanes with elemental sulfur leading to dialkylsulfides and dicycloalkylsulfides, **280**, 355

Cyclodextrin complexes

L- and D-cysteine derivatives of β-cyclodextrin: different molecular recognition properties of their copper(II) complexes for amino acids, **275–276**, 395

Cyclopalladate complexes

cyclopalladated complexes of Schiff bases of homoveratrylamine and tryptamine. Synthesis and CO insertion, **272**, 18

Cyclopentadienyl complexes

high pressure liquid chromatography resolution of racemic mixtures of chiral tricarbonyl arene chromium(0) and cyclopentadienyl-titanium(IV) complexes, **268**, 323

thermal and photochemical substitution reactions of $CpRe(PPh_3)_2H_2$ and $CpRe(PPh_3)H_4$. Catalytic insertion of ethylene into the C–H bond of benzene, **270**, 77

hydrogen site exchange in $CpIr(PR_3)H_3^+$ complexes, **272**, 95

synthesis of chloro and methyl imido cyclopentadienyl molybdenum and tungsten complexes. X-ray molecular structures of $[WCp^*Cl_3(N^tBu)]$, $[MoCp^*ClMe_2(N^tBu)]$ and $[WCp^*ClMe_2(N^tBu)]$, **273**, 270

Fourier transform IR study of the interaction of $[(\eta^5\text{-}C_5H_5)_2FeRu(CO)_4]$ with hydroxylated alumina surfaces, **277**, 61

synthesis and properties of the dichloro[tetramethyl(trifluoromethyl)cyclopentadienyl]ruthenium dimer: X-ray structure of $[M_2(\eta^5\text{-}C_5Me_3CF_3)_2Cl_2(\mu\text{-}Cl)_2]$ (M = Ru, Rh), **280**, 249

Cyclopentadienyl ligands

organometallic Rh(I) and Rh(III) complexes with [2-(dimethylamino)ethyl]cyclopentadiene, **283**, 24

Cyclophosphazenes

an amazing example of yin–yang aminolysis in cyclophosphazenes, **268**, 307

Cycloplatination

synthesis and properties of 3,2′-polymethylene-6-(2″-pyridyl)-2-phenylpyridines and their cycloplatinated complexes, **267**, 265

Cyclopropene complexes

protonation of cyclopropene complexes of platinum(0) and the reduction of cyclopropene by nitrogenases, **280**, 193

Cyclosiloxanolate complexes

bimetallic cyclooligosiloxanolate complexes of copper and nickel, **280**, 282

Cyclo-triphosphorus

the role of cyclo-triphosphorus (P_3) in-plane σ orbitals in coordinating transition metal fragments, **275–276**, 366

Cysteine complexes

solution behaviour and zinc complexation of tripeptides with cysteine and/or histidine at both termini, **272**, 115

Cysteine tetrapeptides

interaction of Hg(II) with tetrapeptides containing cysteinyl and histidinyl residues, **272**, 7

Cytochromes *c*

an electrospray mass spectrometric and voltammetric study of horse heart cytochrome *c* in the presence of metal ions, **267**, 281

effects of solvent on the redox properties of cytochrome *c*: cyclic voltammetry and ^1H NMR experiments in mixed water–dimethylsulfoxide solutions, **272**, 168

^1H and ^{11}B NMR evidence for specific binding of borate ion to cytochrome *c*, **273**, 388

Cytochrome c_2

isolation and characterization of cytochrome c_2 from *Rhodopseudomonas palustris*, **269**, 125

Cytochrome c_3

refinement of the three-dimensional structures of cytochrome c_3 from *Desulfovibrio vulgaris* Hildenborough at 1.67 Å resolution and from *Desulfovibrio desulfuricans* ATCC 27774 at 1.6 Å resolution, **273**, 213

desulfoviridin, the dissimilatory sulfite reductase from *Desulfovibrio desulfuricans* (Essex): new structural and functional aspects of the membranous enzyme, **275–276**, 52

Cytochrome c_4

photosensitized electron transfer reactions of cytochrome c_4 from *Pseudomonas stutzeri* with flavins and methyl viologen, **272**, 109

Cytochrome c_{551}

the effect of surface charge on the reduction potential and heme propionate ionization of a monoheme cytochrome: spectroscopic and potentiometric studies of trifluoroacetylated cytochrome c_{551} from *Pseudomonas aeruginosa*, **275–276**, 256
Cytochrome *c* peroxidase
spectroscopic study of the compound ES and the oxoferryl compound II states of cytochrome *c* peroxidase· comparison with the compound II of horseradish peroxidase, **275–276**, 250
Cytotoxicity
cytotoxic anionic tribromo platinum(II) complexes containing benzothiazole and benzoxazole donors: synthesis, characterization, and structure–activity correlation, **271**, 137
rhodium(III) analogues of antitumour-active ruthenium(III) compounds: the crystal structure of [ImH][*trans*-$RhCl_4(Im)_2$] (Im = imidazole), **273**, 62
gold(III) compounds as potential antitumor agents: cytotoxicity and DNA binding properties of some selected polyamine–gold(III) complexes, **281**, 90

Desulforedoxin
the use of ^{113}Cd NMR chemical shifts as a structural probe in tetrathiolate metalloproteins, **273**, 279
probing the iron environment in desulforedoxin. EXAFS of oxidized and reduced states, **273**, 409
Desulfoviridin
desulfoviridin, the dissimilatory sulfite reductase from *Desulfovibrio desulfuricans* (Essex): new structural and functional aspects of the membranous enzyme, **275–276**, 52
Deuterohaemin complexes
synthesis, characterization and stereoselective catalytic oxidations of chelated deuterohaemin-glycyl-L-histidine complexes, **273**, 339
DFT calculations
ricochet inter-ring haptotropic rearrangement of σ-methyl-(η^5-indenyl)chromium tricarbonyls. Experimental kinetic and theoretical DFT study, **280**, 328
Diamagnetic complexes
NMR conformational study of diamagnetic complexes of some triazatriacetate macrocycles, **273**, 288
Diamide complexes
spectroscopic characterization of coaxially and π–π stacked binuclear copper(II) complexes of ω,ω′-bis(pyridine-2-carboxamido)-alkanes in acidic solution, **283**, 61
Diamidoamine ligand complexes
the synthesis and structures of tantalum complexes that contain a triamido or a diamidoamine ligand, **270**, 353
Diamine complexes
intramolecular and intermolecular C–H activation at a cationic Pt^{II} center, **270**, 467
nickel(II), palladium(II), platinum(II) and platinum(IV) complexes of *cis*-1,2-bis(diphenylphosphino)ethene and nitrogen-containing heterocycles or sulfur ligands, **274**, 143
first solid state synthesis of the racemic compound and enantiomer of dinuclear Λ,Λ-di-μ-hydroxo-bis[bis((1*S*,2*S*)-1,2-*trans*-cyclohexanediamine)chromium(III)] and Δ,Δ-di-μ-hydroxo-bis-[bis((1*R*,2*R*)-1,2-*trans*-cyclohexanediamine)chromium(III)] bromides, **274**, 210
synthesis and structural properties of (diamine)platinum(II) complexes of itaconic acid, **279**, 116
Diaminedithiol complexes
synthesis and structural characterization of the [2,9-dimethyl-4,7-diaza-2,9-decanedithiolato]oxorhenium(V) complex, **268**, 169
Diamines
the stereochemistry of the oxidation of diamines and of the reduction of diimines in the coordination sphere of ruthenium(II), **272**, 153
Diaza-*nido*-hexaboranes
clusters $N_2B_3R_6$ from the fusion of rings NB_2R_3: mechanism, **269**, 43
Diazene complexes
transition metal complexes with sulfur ligands. Part CXXIX. Retention and reactivity of the [Fe⎓NH⎓NH⎓Fe] chromophore in the iron sulfur diazene complex [μ-N_2H_2{$Fe(PPr_3)('S_4')$}$_2$] in exchange and oxidation processes. ('S_4'$^{2-}$ = 1,2-bis(2-mercaptophenylthio)ethane(2−)), **280**, 39
Diazenido complexes
the synthesis and characterization of [$Tc(SPh)_2$(N=NC_5H_4N)-(HN=NC_5H_4N)]$_2$, a paramagnetic dimer with bridging phenylthiolate ligands. The X-ray structure of [$Tc(SPh)_2$-(N=NC_5H_4N)(HN=NC_5H_4N)]$_2$, **267**, 165
Diazolinium complexes
Anderson–Evans type molybdotellurates with diazolium cations: preparation, structure and properties of $(H_2imz)_6$[$TeMo_6O_{24}$]·$4H_2O$ and $(H_2pyz)_6$[$TeMo_6O_{24}$]·$Te(OH)_6$, **277**, 139
Dibenzylphosphine
the reaction of cobalt powder with tetraiodo{1,2-bis(dibenzylphosphino)ethane} to form 1,2-bis(dibenzylphosphino)-ethane cobalt diiodide, $Co(dBzP_2)I_2$; and the X-ray crystal structure of the diphosphinodioxide complex, $Co\{dBzP_2(O)_2\}I_2$, **282**, 25
Dicyanoargentate
chain-like crystal structure and magnetic properties of [catena-bis(1,2-diaminoethane)copper(II)-μ-dicyano-argentate]-dicyanoargentate, **281**, 134
Dicyano-porphyrin complexes
crystal structure and molecular stereochemistry of the 1:2 adduct of anionic complex, dicyano-(5,10,15,20-tetramesityleneporphinato)-iron(II) with bis(triphenylphosphine)nitrogen (1+)·I_2, $CDCl_3$ solvate {[$(CN)_2Fe^{II}$(TMP)][PPN]$_2$}·I_2·$2CDCl_3$, **277**, 242
Diene complexes
structural preferences of η^4-dienecyclopentadienyl complexes: molecular mechanics, molecular orbital and crystallographic studies, **275–276**, 263
Diene cyclomagnesation
cyclomagnesation of dienes catalyzed by a chiral ansa-zirconocene, **280**, 189
Dihydrogen activation
probing the nature of H_2 activation in catalytic asymmetric hydrogenation, **270**, 285
Dihydrogen complexes
solution and crystal structure of the dihydrogen complex [$Ru(H_2)(H)(PMe_2Ph)_4$]PF_6, an active alkyne hydrogenation catalyst, **270**, 238
crystalline dihydrogen complexes. Intramolecular and intermolecular interactions and dynamic behavior, **273**, 116
NMR evidence for formation of new alcohol rhenium complexes as intermediates in ionic hydrogenations of carbonyl groups with systems composed of $ReH_5(NO)(CO)(PR_3)_2$ (R = Pr^i, CH_3, OPr^i) and CF_3COOH, **278**, 122
Di-imidazole complexes
a reversible Cu^{II}/Cu^{I} electron transfer centre based on a flexible di-imidazole ligand, **273**, 225
Diimine complexes
the stereochemistry of the oxidation of diamines and of the reduction of diimines in the coordination sphere of ruthenium(II), **272**, 153
pentacoordinated cationic rhodium(I) and iridium(I) organocomplexes with α-diimine ligands derived from glyoxal: crystal structure of [Rh(norbornadiene)(glyoxalbis(4-dimethylamino-anil))-(PPh_3)]BF_4·$1/2CH_2Cl_2$, **282**, 230
Diiodotyrosine complexes
electrostatic ligand–ligand interactions in ternary copper(II) complexes with 3,5-diiodo-L-tyrosine and polar amino acids, **278**, 136
Diiron complexes
synthesis and characterization of mixed valence μ-alkoxo-diiron-(II,III) complexes with an unsymmetric dinucleating ligand, **283**, 124

Diketonate complexes
synthesis, X-ray structure and bonding of tris(2,2-6,6-tetramethylheptane-3,5-dionato) bismuth(III), **275–276**, 340
β-Diketonate complexes
synthesis of a new series of Nb(IV) β-diketonates with the $Nb_2S_4^{4+}$ core: crystal structure of $Nb_2S_4(CF_3COCHCOCH_3)_4$, **271**, 222
β-Diketone complexes
role of the intramolecular hydrogen bond and ligand rigidity in the complexation of trifluoroacetylcycloalkanones with lanthanides: novel strategy for the design of organic ligands of high selectivity, **267**, 201
the effect of end-group substitutions and use of a mixed solvent system on β-diketones and their iron complexes, **278**, 170
β-Diketonato complexes
eight-coordinate chelate complexes of zirconium(IV) and niobium(IV): X-ray diffractometric and EPR investigations, **270**, 177
β-Diketophenolato complexes
synthetic, spectroscopic, and X-ray crystallographic studies on the uranyl(VI) complexes of the β-diketophenol ligands 1-(2-hydroxyphenyl)-1,3-butanedione and 1-(2-hydroxyphenyl)-3-phenyl-1,3-propanedione, **277**, 37
Dimeric complexes
properties and kinetics of dihydroxy- and diaminoanthraquinone ruthenium bipyridyl dimers, **268**, 31
mechanistic details for metathetical exchange between XCO (X = O and RN) and the tin(II) dimer, $\{Sn[N(SiMe_3)_2](\mu\text{-}OBu^t)\}_2$, **270**, 118
Dimerization of olefins
zirconocene–alumoxane (1:1) – a catalyst for the selective dimerization of α-olefins, **270**, 20
Dimethyl sulfide complexes
thermodynamics of the formation of *cis*-bis(dialkylsulfide)dimethylplatinum(II) complexes from bis(μ-dialkylsulfide)bisdimethylplatinum(II) and free dialkylsulfide, **268**, 311
Dimethylsulfoxide complexes
synthesis of binuclear platinum(II) organometallic complexes as potential cross-linking reagents, **275–276**, 141
Dinitrogen complexes
syntheses and properties of organocyanamide, cyanoguanidine and dinitrogen complexes of rhenium. Crystal structure of *mer*-$[ReCl_2(NCNEt_2)(PMePh_2)_3]$, **280**, 308
Dinuclear complexes
effects of ligand topology on the properties of dinuclear ruthenium complexes of bis-semiquinone bridging ligands, **267**, 1
syntheses, structures and properties of hydrosulfido-bridged diiridium and dirhodium complexes, $[Cp^*MCl(\mu_2\text{-}SH)_2MCp^*Cl]$ and $[Cp^*M(\mu_2\text{-}SH)_3MCp^*]^+$ (M = Ir, Rh; $Cp^* = \eta^5\text{-}C_5Me_5$), **267**, 73
three dinuclear copper(II) carboxylates with the paddle-wheel cage structure as intermediates in copper(II) catalyzed oxidations of carboxylic acids. X-ray crystal structures of [tetrakis(diphenyl acetato-μ-*O*,*O*′)bis(acetonitrile-*N*)dicopper(II)] tetrakis(acetonitrile), [tetrakis(diphenylacetato-μ-*O*,*O*′)bis(acetone-*O*)dicopper(II)] and [tetrakis(1-phenyl-1-carboxylato-μ-*O*,*O*′-*cyclo*-pentane)-bis(ethanol-*O*)dicopper(II)] bis(ethanol), **267**, 183
site-selective PMe_3 addition to the donor–acceptor complex $Ru_2(CO)_6[(Z)\text{-}Ph_2PCH{=}CHPPh_2]$. X-ray diffraction structure of $Ru_2(CO)_5(PMe_3)[(Z)\text{-}Ph_2PCH{=}CHPPh_2]$, **267**, 223
NMR investigations of di-rhodium-tetraacetate complexes with cyclic phosphites, **267**, 293
NMR spectra of dinuclear manganese and iron compounds containing 1,4,7-triazacyclononane and 1,4,7-trimethyl-1,4,7-triazacyclononane, **268**, 43
copper(II) complexes with tridentate O,N,O-chelate dianions and additive monodentate donor ligands. I. Crystal structures and EPR results of urea adducts of *N*-salicylidene-α-aminoacidato-copper(II) compared with *N*-salicylideneglycinato-thiourea-copper(II), **268**, 117
homodinuclear lanthanide complexes: Ln_2L_3 (H_2L = tetradentate Schiff bases). Magnetic properties (solid state) and spectroscopic studies (solution), **268**, 125
syntheses and structure of dinuclear cobalt(II) complexes of the polypyridine ligand 6,6′-bis[bis(2-pyridylmethyl)aminomethyl]-2,2′-bipyridine, **268**, 159
an asymmetrically-coordinated 2,3-dimethylpyrazine bridging ligand in the (II,II) binuclear ion, $\{[Ru^{II}(hedta)]_2(2,3\text{-}Me_2pz)\}^{2-}$ ($hedta^{3-}$ = *N*-hydroxyethylethylenediaminetriacetate), **268**, 279
copper(II) complexes of pyrazol-3-yl amines. Synthesis, crystal and solution structure, **269**, 117
the synthesis and characterization of α-$Re_2Cl_4(depe)_2$ and α-$W_2Cl_4(depe)_2$, **271**, 49
synthesis, characterization and reactivity of bis(diphenylphosphino)-amine nickel complexes, **271**, 112
synthesis and crystal structure of two ternary dicopper(I) complexes having the unsymmetrical coordination arrangement bridged by 1,8-naphthyridine (napy). $[Cu_2(napy)_2(Me_2CO)](PF_6)_2 \cdot 2Me_2CO$ and $[Cu_2(napy)_2(dppm)(CH_3CN)](PF_6)_2$, **271**, 129
synthesis and physical studies of copper(I) complexes with the ambidentate ligand (1-piperidyl-2-diphenylphosphino)ethane. X-ray structural characterization of a copper(I) dimer, **271**, 217
an unusual cyanide-bridged extended A-frame complex: the molecular structure of $[\{Pd_2Br_2(\mu\text{-}PPh_2)(\mu\text{-}dppm)\}_2(\mu\text{-}CN)_2]$, **271**, 236
synthesis, crystal structure and magnetic properties of bis[μ-bis(3-(pyridin-2-yl)-1,2,4-triazole-*N*′,*N*1,*N*2)] bis[triaqua nickel(II)] tetranitrate, **273**, 54
unexpected assembly of a (μ-oxo)(μ-formato)diiron(III) complex from an aerobic methanolic solution of Fe(III) and the TPA ligand, **273**, 393
exchange coupling across multiple-atom bridges: crystal structure and magnetic properties of a chloranilate bridged dicopper(II) complex, **274**, 111
synthesis and crystal structure of a seven-coordinate dinuclear iron(III) complex containing only bidentate ligands, **274**, 224
syntheses and structures of some Ru_2^{5+} complexes containing weakly coordinating anions as axial ligands, **275–276**, 447
liquid secondary ion mass spectrometric study of diruthenium(II,III) complexes, **277**, 76
reaction of the dinuclear cobalt complex $[Co_2\{\mu\text{-}PPh_2CHCHC(O)\text{-}CHCHPPh_2\}(CO)_4]$ with two-electron donor ligands; the stabilisation of cobalt–alkene bonds by incorporation of alkenes into a bridging diphosphine ligand, **277**, 186
synthesis of a dinuclear, phosphido-bridged complex of rhenium, X-ray crystal structure of $Re_2(\mu\text{-}H)[\mu\text{-}(t\text{-}Bu)PC_6H_4P(H)t\text{-}Bu](CO)_7$, **277**, 253
a series of $[Cr_2(\mu\text{-}OH)_n(\mu\text{-}RCO_2)_{3-n}(bispma)_2]$ type chromium(III) dinuclear complexes (n = 1, 2) with tridentate bis(2-pyridylmethyl)amine (bispma): synthesis, crystal structure, and magnetic properties, **278**, 15
synthesis, structural characterization and magnetic behaviour of two new $(\mu_{1,3}\text{-}N_3)_2$ dinuclear nickel compounds: study of the magnetic transition at 235 K for the *cis*-$(\mu_{1,3}\text{-}N_3)_2[Ni(dl\text{-}cth)]_2(ClO_4)_2$ compound, **278**, 43
synthesis, characterization and kinetics of the metallation of the intermediate compound $[Rh_2(\mu\text{-}O_2CCH_3)_3(O_2CCH_3)\{\eta^2\text{-}\underline{P}(2\text{-}(CH_3\underline{O})C_6H_4)Ph_2\}] \cdot H_2O$ to produce compound $[Rh_2(\mu\text{-}O_2CCH_3)_3\text{-}\{\mu\text{-}(\underline{C}\underline{C}_5H_4)\underline{P}(2\text{-}(CH_3O)C_6H_4)Ph\}] \cdot (H_2O)$ from Rh(II) tetraacetate, **278**, 61
synthesis and crystal structure of a new salt of the complex anion $[(TCNQ)_3]^{2-}$ $[Mo_2(\mu_2\text{-}S_2)_2(S_2CNMe_2)_4](TCNQ)_3$, **278**, 108
the chemistry of dinitrato-2,2′-bipyridinecopper(II): preparation and characterization of binuclear complexes having a $Cu(\mu\text{-}OH)_2Cu$ and $Cu(\mu\text{-}N_3)_2Cu$ core, **278**, 143

synthesis of tetracarboxylatediruthenium(II) compounds: reduction and disproportionation reactions of diruthenium(II,III) units. Crystal structure of $Ru_2(\mu\text{-}O_2CCOPh)_4(thf)_2$, **279**, 30

synthesis and crystal structure of a new isomer of the μ-oxo-bis[dichlorooxobis(pyridine)rhenium(V)] complex $\{OReCl_2py_2\}_2O$, **279**, 159

dialkyl-μ-ethylidene-μ-methylene-bis(pentamethylcyclopentadienyl)-dirhodium complexes: synthesis, spectra and decomposition reactions, **280**, 62

crystal structure, spectral and magnetic properties of a new (μ-acetate)(μ-alkoxide) dicopper(II) complex as a model for tyrosinase, **281**, 111

syntheses, spectroscopic characterization and X-ray crystal structures of $[HP\text{-}i\text{-}Pr_3]_2[IrCl_6]$ (**1**), $[HP\text{-}i\text{-}Pr_3]_2[Ir_2Cl_8(P\text{-}i\text{-}Pr_3)]$ (**2**) and $[HP\text{-}i\text{-}Pr_3][Ir_2Cl_7(P\text{-}i\text{-}Pr_3)_2]$ (**3**): three possible precursors of the complex $Ir(H)(Cl)_2(P\text{-}i\text{-}Pr_3)_2$ (**4**), **282**, 17

synthesis and characterization of mixed valence μ-alkoxo-diiron-(II,III) complexes with an unsymmetric dinucleating ligand, **283**, 124

preparation and properties of a series of (μ-oxo)di(μ-acetato) diruthenium(III) complexes, $[Ru_2(\mu\text{-}O)(\mu\text{-}CH_3COO)_2(pyridine)_4(L)_2]^{2+}$ and $[Ru_2(\mu\text{-}O)(\mu\text{-}CH_3COO)_2(AA)_2(L)_2]^{2+}$, where AA is 2,2′-bipyridine or 1,10-phenanthroline, and L is a monodentate ligand, **283**, 151

synthesis and crystal structure of a dinuclear platinum(II) bis(*N*,*N*-diethylamino)carbeniumdithiocarboxylate complex $Pt_2\{\mu\text{-}(Et_2N)_2C_2S_2\}_2Cl_4\cdot 0.5CH_2Cl_2$: structural study of the coordinated inner salt, **283**, 256

Dinucleating ligand complexes

synthesis and characterization of mixed valence μ-alkoxo-diiron-(II,III) complexes with an unsymmetric dinucleating ligand, **283**, 124

Dinucleating macrocycles

dinuclear $Cu^{II}M^{II}$ complexes of a phenol-based macrocycle with $N(amine)_2O_2$ and $N(imine)_2O_2$ metal-binding sites: the effect of chloride ligation upon the site selectivity of metal ions, **283**, 72

Diolate complexes

synthesis, characterization and study of the redox properties of rhenium(V) diolates, **271**, 65

β-Dione complexes

axial–equatorial interactions in chelates of β-diones. Crystal and molecular structure of bis(3-nitro-2,4-pentanedionato)copper(II), **269**, 332

Diorganotin(IV) complexes

unusual coordination behavior of D-fructose towards dimethyltin(IV): metal-promoted deprotonation of alcoholic OH groups in aqueous solutions of low pH, **274**, 167

Dioxygen complexes

electrochemistry of 2-thio- and 2-oxo-pyrimidines in dimethyl formamide in the presence of dioxygen, **270**, 145

spectroscopic and magnetic properties and structure of a five-coordinate, O_2-binding cobalt(II) Schiff base complex and of the copper(II) analogue, **278**, 127

electrochemistry of dioxygen activation in the presence of adenine, **280**, 322

Diphosphadiboracyclobutane complexes

some oxidative addition reactions with a diphosphadibora[1.1.0]bicyclobutane: formation of cage compounds and of an unusual trinuclear coordination compound of palladium, **269**, 101

Diphosphene complexes

comparative DFT and HF-MP2 study on structure, relative stabilities and rotational barriers of end-on (η^1) and side-on (η^2) $(PH_3)_2Pd$ HPPH model complexes, **273**, 72

Diphosphine-bridged complexes

synthesis and crystal structure of dinuclear palladium(I) dppm complex with 4-mercaptopyridine in the thione mode. $[Pd_2(\mu\text{-}dppm)_2(HpyS)_2](ClO_4)_2\cdot CH_3OH$, **281**, 116

Diphosphine complexes

site-selective PMe_3 addition to the donor–acceptor complex $Ru_2(CO)_6[(Z)\text{-}Ph_2PCH{=}CHPPh_2]$. X-ray diffraction structure of $Ru_2(CO)_5(PMe_3)[(Z)\text{-}Ph_2PCH{=}CHPPh_2]$, **267**, 223

the synthesis and characterization of *trans*-$ReCl_2(dppe)_2$ and α-$Re_2Cl_4(dppe)_2$, **268**, 151

ligand connected metal clusters. The molecular structures and solid state packing of $\{Ru_3(CO)_{11}\}_2$(bis(diphenylphosphino)ethane) and $\{Ru_3(CO)_{11}\}_2$(1,4-bis(diphenylphosphino)benzene), **270**, 28

triply-bridged diruthenium(II) 1,4-bis(diphenylphosphino)butane (dppb) and (*R*)-2,2′-bis(diphenylphosphino)-1,1′-binaphthyl (binap) complexes, including structural characterisation of $[(dppb)ClRu(\mu\text{-}D_2O)(\mu\text{-}Cl)_2RuCl(dppb)]$, $[(\eta^2\text{-}H_2)(dppb)Ru(\mu\text{-}Cl)_3RuCl(dppb)]$ and the $[(dppb)ClRu(\mu\text{-}Cl)_3RuCl(dppb)]^-$ anion, **270**, 130

probing the nature of H_2 activation in catalytic asymmetric hydrogenation, **270**, 285

hydride complexes of platinum(II) containing unsymmetrical di(phosphine) ligands: synthesis, characterization and evidence for pairwise addition of hydrogen based on parahydrogen induced polarization, **270**, 363

synthesis and crystal structure of two ternary dicopper(I) complexes having the unsymmetrical coordination arrangement bridged by 1,8-naphthyridine (napy). $[Cu_2(napy)_2(Me_2CO)](PF_6)_2\cdot 2Me_2CO$ and $[Cu_2(napy)_2(dppm)(CH_3CN)](PF_6)_2$, **271**, 129

nickel(II), palladium(II), platinum(II) and platinum(IV) complexes of *cis*-1,2-bis(diphenylphosphino)ethene and nitrogen-containing heterocycles or sulfur ligands, **274**, 143

reaction of the dinuclear cobalt complex $[Co_2\{\mu\text{-}PPh_2CHCHC(O)CHCHPPh_2\}(CO)_4]$ with two-electron donor ligands; the stabilisation of cobalt–alkene bonds by incorporation of alkenes into a bridging diphosphine ligand, **277**, 186

a simple and convenient synthesis of *cis*/*trans*-$RuH_2(Ph_2PCH_2PPh_2)_2$ and of *trans*-$RuHCl(Ph_2PCH_2PPh_2)_2$, **278**, 226

Diphosphine selenide complexes

metal complexes of $P(Ph_2)CH_2(Ph_2)PSe$. Crystal structure of $[Ni\{P(Ph_2)CH_2(Ph_2)PSe\}_2]Cl_2\cdot 2EtOH$, **273**, 320

Diphosphinodioxide complexes

the reaction of cobalt powder with tetraiodo{1,2-bis(dibenzylphosphino)ethane} to form 1,2-bis(dibenzylphosphino)-ethane cobalt diiodide, $Co(dBzP_2)I_2$; and the X-ray crystal structure of the diphosphinodioxide complex, $Co\{dBzP_2(O)_2\}I_2$, **282**, 25

Diphosphino-pyridone complexes

unique complexation by the tridentate ligands, 6-diphenylphosphino-2-pyridonate (pyphos) and 6-diphenylphosphino-2-pyridone (pyphosH), on ruthenium(II): crystal structures of (pyphosH)-$RuCl(\mu\text{-}Cl)Mo_2Cl(pyphos)_3$, $RuCl_2(CO)(pyphosH)_2$, $[Ru(pyphos)_3](\mu\text{-}Na)_2[Ru(pyphos)_3]$, **270**, 459

Diphosphorane complexes

the reaction of cobalt powder with tetraiodo{1,2-bis(dibenzylphosphino)ethane} to form 1,2-bis(dibenzylphosphino)-ethane cobalt diiodide, $Co(dBzP_2)I_2$; and the X-ray crystal structure of the diphosphinodioxide complex, $Co\{dBzP_2(O)_2\}I_2$, **282**, 25

Dipyrazolylmethane complexes

dinuclear complexes of first transition series metals with 4,4′-dipyrazolylmethane: characterisation, DNA binding and anticancer properties, **271**, 151

Distance geometry calculations

Boc-Glu-Thr-Ile-His-OMe/Zn^{2+} in terms of the zinc-binding sites of proteases, **283**, 243

Disulfide complexes

first- and second-order mechanisms for oxidative addition for bound methyl disulfide in the complex $W(CO)_3(phen)(MeSSMe)$, **270**, 488

Disulfide compounds

X-ray absorption fine structure study of the atomic and electronic structure of molybdenum disulfide intercalation compounds with transition metals, **280**, 211

Dithiocarbamate complexes
synthesis and characterization of molybdenum(II) and molybdenum(V) monothiocarbamate complexes. Crystal and molecular structures of tetrakis(isopropylmonothiocarbamato)dimolybdenum(II) and oxobis[di(oxoisopropylmonothiocarbamato)-molybdenum(V)], **267**, 101
substituent effects of *cis*-dioxobis(dithiocarbamato)molybdenum(VI) on redox properties: redox potentials for one-electron reduction and second-order rate constants for oxygen atom transfer, **269**, 260
syntheses of ruthenium hydridotris(1-pyrazoryl)borate complexes having sulfur-donor coligands, **273**, 238
synthesis and molecular structure of the molybdate-bridged cyclic imido complex $[Mo(S_2CNEt_2)_2(NR)(\mu\text{-}MoO_4)]_2$ (R = 2,6-$^iPr_2C_6H_3$), **274**, 243
polynuclear silver compound formed from aggregation of Se^{2-} and Ag(I)-thiolate complex. Synthesis, structure and spectroscopic characterization of $Ag_{11}(\mu_5\text{-}Se)(\mu_4\text{-}Et_2NCS_2)_3(\mu_3\text{-}Et_2NCS_2)_6$, **277**, 171
insulin-mimetic vanadyl–dithiocarbamate complexes, **283**, 175

Dithiocarbamato complexes
synthesis and crystal structure of a new salt of the complex anion $[(TCNQ)_3]^{2-}$ $[Mo_2(\mu\text{-}S_2)_2(S_2CNMe_2)_4](TCNQ)_3$, **278**, 108

Dithiodipyridine complexes
synthesis and properties of the unsymmetrical binuclear complexes: $Na[(CN)_5Fe(dtdp)Ru(NH_3)_5]6H_2O$ and $K[(CN)_5Ru(dtdp)Ru(NH_3)_5]6H_2O$, dtdp = 4,4′-dithiodipyridine, **267**, 93

Dithiolate complexes
a tetranuclear silver complex with dppm and i-mnt ligands. Synthesis and structural characterization of $[Ag_4(\mu\text{-dppm})_4(\mu_4\text{-i-mnt})_2]\cdot 0.5$ DMF (i-mnt = 2,2-dicyano-1,1-ethylenedithiolate; dppm = bis(diphenylphosphino)methane), **267**, 313
synthesis, electrical properties of $[Ni(dmit)_2]^-$ complexes and X-ray crystal structure of [EDA][Ni(dmit)$_2$] (dmit = 1,3-dithiol-2-thione-4,5-dithiolate), **268**, 103
nickel(II) 1,1-dithiolates and related complexes. Solid-state and solution structural evidence for mixed-ligand complexes with substituted ethylenediamines, **271**, 195
polynuclear cobalt cluster compounds formed via sulfur–carbon bond cleavage of 2,2-dicyano-1,1-ethylenedithiolate, **274**, 229
preparation and properties of $C_8H_4S_8$-platinum(II) complexes and electrical conductivities of their oxidized species and X-ray crystal structure of $C_8H_4S_8(CH_2CH_2CN)_2$ as a pro-ligand compound, **279**, 165
electrocrystallization and characterization of phosphonium-based radical anion salts $[MePh_3P][Pd(dmit)_2]_3$ and $[Ph_4P][Pd(dmit)_2]_3$ (dmit = 1,3-dithiole-2-thione-4,5-dithiolate). Comparison with related Pd and Ni isologues, **282**, 90

Dithiolene complexes
synthesis of nitridoosmium(VI) dithiolene complexes, **267**, 335
novel oxidation and reduction products of the neutral nickel-dithiolene [Ni(Pr$^i_2$timdt)$_2$] (Pr$^i_2$timdt is the monoanion of 1,3-diisopropylimidazolidine-2,4,5-trithione), **273**, 175
a single-crystal electron paramagnetic resonance, ^{13}C and ^1H electron nuclear double resonance study of tetra-*n*-butylammonium-bis(1,2-dicyanoethylene-1,2-dithiolato)aurate(II), $[(n\text{-}C_4H_9)_4N]_2$-$[Au^{II}(mnt)_2]$, **281**, 160
the synthesis and characterization of platinum monodithiolene complexes containing 1,3-dithiole-2-oxo-4,5-dithiolate ($dmid^{2-}$) and 1,3-dithiole-2-thione-4,5-dithiolate ($dmit^{2-}$), **282**, 200

Dithionate complexes
synthesis, structure and properties of lanthanide dithionates and their triphenylphosphine oxide complexes, **274**, 177

Dithionite complexes
reduction of *Coprinus macrorhizus* peroxidase by dithionite, **274**, 215

Dithiophosphate complexes
stopped-flow spectrophotometric study on the reactions between copper(II) ions and some dithiophosphates, **272**, 291

DMF complexes
synthesis and properties of $[M_6X_{12}(C_3H_7NO)_6]X_2$ (M = Nb, Ta; X = Cl, Br), **271**, 239

DMSO complexes
ligand photolysis and recombination of (dimethylsulfoxide)$_2$Fe(II)-porphyrin complexes, **271**, 119
comparison of the reactivity of *cis*-$[RuCl_2(DMSO)_4]$ and *trans*-$[RuCl_2(DMSO)_4]$ towards nucleosides, **281**, 10

DNA
intercalation into calf thymus DNA of 2-(4-arylphenyl)imidazo-[4,5-f]-[1,10]phenanthroline (aryl = -OMe, -NMe_2 or -NO_2) in its bipyridyl ruthenium(II) complex, **283**, 17

DNA binding
dinuclear complexes of first transition series metals with 4,4′-dipyrazolylmethane: characterisation, DNA binding and anticancer properties, **271**, 151
influence of *cis*- and *trans*-diamminedichloroplatinum(II) binding on the helix-coil transition of DNAs with different GC content, **275–276**, 510

DNA complexes
the role of $O_2^{\bullet-}$ radicals in DNA–metal complexes, **273**, 201

DNA interaction
zinc as a structural and folding element of proteins which interact with DNA, **275–276**, 481

DNA nicking complexes
DNA nicking by a trinuclear chromium complex, **268**, 211

DNA oxidation
metallodrug complexes that mediate DNA and lipid damage via sulfite autoxidation: copper(II) famotidine and iron(III) bis(salicylglycine), **275–276**, 314

Drug metabolism
sulfonated and acetamidosulfonylated tetraarylporphyrins as biomimetic oxidation catalysts under aqueous conditions, **272**, 228

D–s mixing
the effects of d–s mixing in low symmetry transition metal complexes, **268**, 55

DTPA complexes
multinuclear NMR study of the interactions between the La(III) complex of DTPA-bis(glucamide) and Zn(II) or borate, **268**, 249

EDA complexes
structures and electronic states of the electron donor–acceptor (EDA) complexes: trimethylamine–halogen complex $(CH_3)_3N\cdots F_2$, **281**, 85

Edta complexes
structural characterization of an eight-coordinate Nb(IV)–edta complex, $[Nb(edta)(H_2O)_2]\cdot 2H_2O$, **274**, 103
preparation and X-ray crystal structures of $Li_4[Hg_2(edta)_2]\cdot 8H_2O$ and $Ba_2[Hg_2(teta)_2]\cdot 9H_2O$ (teta = *N,N,N′,N′*-diaminobutanetetraacetate), **274**, 219
some hexadentate Ni(II)-edta-type complexes containing five-membered diamine rings. The molecular and crystal structure of the *trans*(O_5) isomer of barium(ethylenediamine-*N,N′*-diacetato-*N,N′*-di-3-propionato)nickelate(II) hexahydrate, *trans*(O_5)-$Ba[Ni(eddadp)]\cdot 6H_2O$, and strain analysis of edta-type chelates in relation to their octahedral distortion, **278**, 66
substitution reaction of aqua(ethylenediaminetetraacetato) ruthenium(III) with 2-mercaptobenzoic acid, **282**, 38

EDTMP complexes
potentiometric and computer studies of yttrium–EDTMP, **275–276**, 37

Eight-coordinate complexes
an infrequent geometry for an element of the first transition series: structural characterization of a distorted eight-coordinate compound $TiCl_4[C_2H_4(PMe_2)_2]_2$, **267**, 173

eight-coordinate chelate complexes of zirconium(IV) and niobium(IV): X-ray diffractometric and EPR investigations, **270**, 177
structural characterization of an eight-coordinate Nb(IV)–edta complex, $[Nb(edta)(H_2O)_2]\cdot 2H_2O$, **274**, 103

Electrical conductivity
synthesis, electrical properties of $[Ni(dmit)_2]^-$ complexes and X-ray crystal structure of $[EDA][Ni(dmit)_2]$ (dmit = 1,3-dithiol-2-thione-4,5-dithiolate), **268**, 103

Electric properties
synthesis and characterization of electric and magnetic properties of the new intermetallic compounds $M_3(GaTe_3)_2$ (M = Cr, Fe, Co), **268**, 271

Electrocatalysis
electrochemistry of 2-thio- and 2-oxo-pyrimidines in dimethyl formamide in the presence of dioxygen, **270**, 145
application of the electrocatalytic reduction of nitric oxide mediated by ferrioxamine B to the determination of nitric oxide concentrations in solution, **273**, 316

Electrochemical kinetics
infrared spectroelectrochemical and electrochemical kinetics studies of the reaction of nickel cluster radicals $[Ni_3(\mu_2\text{-dppm})_3(\mu_3\text{-L})(\mu_3\text{-I})]^+$(L = CNR, R = CH_3, i-C_3H_7, C_6H_{11}, $CH_2C_6H_5$, t-C_4H_9, 2,6-$Me_2C_6H_3$; L = CO) with carbon dioxide, **270**, 111

Electrochemical synthesis
electrochemical synthesis and characterization of indium pyrimidine-2-thiolate complexes. Molecular structure of tris(pyrimidine-2-thiolato)indium(III) and tris[(5-ethyl-4,6-dimethyl)pyrimidine-2-thiolato]indium(III), **274**, 131

Electrochemistry
4- and 3-hydroxypyridine complexes of pentacyanoferrate(II, III), **267**, 81
substituent effects of *cis*-dioxobis(dithiocarbamato)molybdenum(VI) on redox properties: redox potentials for one-electron reduction and second-order rate constants for oxygen atom transfer, **269**, 260
redox behavior of a binuclear ruthenium complex having a di-μ-nitrosyl ligand, $[\{Ru(acac)_2\}_2(\mu\text{-NO})_2]$ (acac = acetylacetonato), **269**, 342
cooperative bimetallic reactivity, **270**, 13
nitrogen atom transfer and redox chemistry of terpyridyl phosphoraniminato complexes of osmium (IV), **270**, 511
complexes of copper(I), silver(I), nickel(II) and cadmium(II) with 6-tert-butyldimethylsilylpyridine-2-thionate (6-Bu^{t}SidmepyS$^-$): crystal structure of $[Cu_2(6\text{-Bu}^t\text{SidmepyS})_2\text{dppm}_2]$, **271**, 1
synthesis, spectroscopic and electrochemical study of nickel(II) complexes with tetradentate asymmetric Schiff bases derived from salicylaldehyde and methyl-2-amino-1-cyclopentenedithiocarboxylate, **271**, 83
a new manganese-β-heptanitro-porphyrin with extreme redox potentials: spectral, electrochemical and catalytic properties, **272**, 4
effects of solvent on the redox properties of cytochrome *c*: cyclic voltammetry and ^{1}H NMR experiments in mixed water–dimethylsulfoxide solutions, **272**, 168
rhenium(VII) and (V) imido and oxo complexes containing 1,4,7-triazacyclononane: synthesis, electrochemistry and crystal structures, **272**, 176
mono- and bis-alkynyl ruthenium(II) complexes containing the ferrocenyl moiety; crystal structure of *trans*-$[Ru(C{\equiv}CC_5H_4FeC_5H_5)_2(Ph_2PCH_2CH_2PPh_2)_2]$ and electrochemical studies, **272**, 188
oligonuclear ferrocenolato and 1,1′-ferrocenediolato derivatives of phosphorus: synthesis, structure and electrochemical behaviour, **273**, 24
tris(4,4′-dimethoxy-2,2′-bipyridine)osmium(II), amperometric properties and crystal structure, **274**, 64
photosubstitution reactivity, crystal structures, and electrochemistry of ruthenium(II)(III) complexes containing tetradentate (O_2N_2, S_2N_2, and P_2N_2) Schiff base ligands, **274**, 184
electrochemical and photophysical properties of two-component coordination compounds containing a metallocyclam and an $Re^I(bipy)(CO)_3Cl$ subunit, **275–276**, 117
synthesis, UV–visible and electrochemical studies of lipophilic and hydrophilic lanthanide(III) bis(porphyrinates), **275–276**, 182
study of the relative π-acceptor ability of unsaturated ligands $:C(=C)_nPh_2$ (n = 0, 1, 2 and 4) based on cyclic voltammetry of complexes *trans*-$[RhCl(L)(P^iPr_3)_2]$ (L = $:C(=C)_nPh_2$, C_2H_4 and CO), **275–276**, 435
migration of $[Ru^{II}(hedta)]^-$ among three binding sites on 6-azauridine, **277**, 46
manifestation of redox duality of 2-propanone oxime: Pt(II)-assisted reduction versus Pt(IV)-mediated oxidation of $Me_2C{=}NOH$ species, **277**, 83
preparation and electrochemistry of six-coordinate monooxo molybdenum(VI) complexes containing bidentate catecholate and tridentate NOS-donor Schiff base ligands, **278**, 1
preparation and some properties of perfluoroalkoxy-substituted phthalocyanine complexes of iron(III), nickel(II) and zinc(II), **278**, 253
structural information on *trans*-1,2-diaminocyclohexane-*N*,*N*′-tetraacetateferrate(III) in the solid and aqueous phase, **279**, 37
molecular structure of a five-coordinate binuclear zinc complex, **279**, 249
electrochemistry of dioxygen activation in the presence of adenine, **280**, 322
synthesis, structure, spectroelectrochemical and theoretical investigations of $[(\eta^5\text{-}Me_5Cp)\overline{Co(\eta^4\text{-}MeC{=}CMe{-}CMe{=}CMe)Sn}\{CH(SiMe_3)_2\}_2]$. Formation of a cobalt complexed stannacycle via a mixed alkyne–stannylene cycloaddition reaction, **281**, 53
electrochemical synthesis and characterisation of gallium, indium and thallium compounds with 1-hydroxopyridine-2-thione. The crystal structures of tris(1-oxopyridine-2-thionato)indium(III) and tris(1-oxopyridine-2-thionato)thallium(III), **281**, 70
formation and conversion of yellow heteropoly complexes in a Mo(VI)–Se(IV), Te(IV) system, **281**, 214
study of polyoxotungstates with the Keggin structure by cyclic voltammetry, **281**, 225
electron donating substituent effects on redox and spin state properties of iron(II) bis-terpyridyl complexes, **282**, 131
pentamethylcyclopentadienyl ruthenium(II) complexes containing chiral diphosphines: synthesis, characterisation and electrochemical behaviour. X-ray structure of $(\eta^5\text{-}C_5Me_5)Ru\{(S,S)\text{-}Ph_2PCH(CH_3)CH(CH_3)PPh_2\}Cl$, **282**, 163
electrochemical, spectrochemical characterization and crystal structure of (1,4,8,11-tetrazacyclotetradecane)(acetylacetonato)cobalt(III) complexes, **282**, 173
synthetic models for biological trinuclear copper clusters. Trinuclear and binuclear complexes derived from an octadentate tetraaminetetrabenzimidazole ligand, **282**, 180
structures and electrochemical properties of the Co(III) ternary complexes containing NO_3-type tripodal tetradentate ligands and amino acids: effect of the outer coordination sphere on the electrochemical properties, **283**, 160
potentiometric and spectroscopic studies of transition metal complexes of bis(imidazolyl) and bis(pyridyl) derivatives of amino acids, **283**, 233

Electrocrystallization
electrocrystallization and characterization of phosphonium-based radical anion salts $[MePh_3P][Pd(dmit)_2]_3$ and $[Ph_4P][Pd(dmit)_2]_3$ (dmit = 1,3-dithiole-2-thione-4,5-dithiolate). Comparison with related Pd and Ni isologues, **282**, 90

Electron density
alkyl exchange between aluminum trialkyls and zirconocene dichloride complexes — a measure of electron densities at the Zr center, **270**, 376

Electronic spectra

electronic spectra and structures of Cu^{2+} and Ni^{2+} complexes with dioxotetraazacycloalkanediacetates and their diester derivatives, **269**, 224

Electronic spectroscopy

electronic absorption and MCD spectra for the $Pt(PPh_3)(AuPPh_3)_6^{2+}$ and $Pt(H)(PPh_3)(AuPPh_3)_7^{2+}$ cluster complexes, **278**, 113

Electronic structures

an ab initio MO-LCAO investigation of the electronic structure of organic hydroperoxides and platinum(II) hydroperoxo complexes: a contribution to the knowledge of the mechanism of olefin epoxidation, **270**, 479

carbonyl bonding on ZnO(0001), CuCl(111) and Cu_2O(111) $3d^{10}$ metal ion surfaces: photoelectron spectroscopy of CO/Cu_2O(111) and electronic structure calculations, **275–276**, 327

simulation of the molecular and electronic structure of heterofullerenes $C_{55}Y_5$ (Y = Si, Ge, Sn, B, Al, N, P, SiH, GeH, SnH) and their η^5-π-complexes with Li by the MNDO method, **280**, 219

Electron paramagnetic resonance

an electron paramagnetic resonance study of tetra-*n*-butylammonium-tetrabromo-nitridorhenate(VI), $[(n\text{-}C_4H_9)_4N][ReNBr_4]$, **271**, 199

crystal structure, EPR and magnetic susceptibility studies of tetrakis[μ-(β-alanine)-*O*,*O*′]dichlorodicopper(II) dichloromonohydrate, **275–276**, 28

spectroscopic investigations on the water-soluble fragment of the Rieske [2Fe–2S] protein from *Paracoccus denitrificans*, **275–276**, 493

spectroscopic and magnetic properties of the pyridine-2-carbaldehyde thiosemicarbazonateiron(III) complexes: $[Fe(C_7H_7N_4S)_2]X \cdot nH_2O$ (X = Cl, ClO_4, NO_3, PF_6). Crystal structure of the hexafluorophosphate compound, **278**, 150

the 1,8-bis(2′-pyridyl)-3,6-dithiaoctane complex of nickel(II): X-ray crystal structure and borohydride adduct formation, **278**, 217

spectroscopic characterization of coaxially and π–π stacked binuclear copper(II) complexes of ω,ω′-bis(pyridine-2-carboxamido)-alkanes in acidic solution, **283**, 61

Electron spin echo

interspin distances determined by time domain EPR of spin-labeled high-spin methemoglobin, **273**, 354

Electron paramagnetic resonance–ENDOR study

a single-crystal electron paramagnetic resonance, ^{13}C and ^1H electron nuclear double resonance study of tetra-*n*-butylammonium-bis(1,2-dicyanoethylene-1,2-dithiolato)aurate(II), $[(n\text{-}C_4H_9)_4N]_2$-$[Au^{II}(mnt)_2]$, **281**, 160

Electron spin relaxation

interspin distances determined by time domain EPR of spin-labeled high-spin methemoglobin, **273**, 354

Electron transfer

preparation and characterization of water-soluble viologen-linked zinc porphyrin and bisviologen-linked zinc porphyrin, **267**, 257

isolation and characterization of cytochrome c_2 from *Rhodopseudomonas palustris*, **269**, 125

the cyanide ligand as an efficient bridge in mixed-valence complexes, **269**, 246

comparative time-resolved IR and UV spectroscopic study of monophosphine and diphosphine platinum(II) azido complexes, **271**, 160

roles of valine-98 and glutamic acid-72 of putidaredoxin in the electron-transfer complexes with NADH-putidaredoxin reductase and P450cam, **272**, 80

photosensitized electron transfer reactions of cytochrome c_4 from *Pseudomonas stutzeri* with flavins and methyl viologen, **272**, 109

stereoselectivity in reactions of metal complexes. Part XXI. Kinetics and mechanism of electron transfer between *rac*-$[Co(L)H_2O]^+$ (L = *N*,*N*′-[(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]) and optically active iron(II) complexes, **275–276**, 1

stereoselectivity in reactions of metal complexes. Part XXII. Kinetics and mechanism of electron-transfer reactions between *rac*-$[Co(L)X]^+$ (L = [(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]; X = pyridine, imidazole, 1-methylimidazole and pyrazine) and optically active iron(II) complexes, **275–276**, 9

synthesis of porphyrin–spacer–quinone compounds via metal-mediated cross-coupling: new systems for probing the relative magnitudes of axial and equatorial electronic coupling at the porphyrin macrocycle in thermal and photoactivated electron transfer reactions, **275–276**, 427

electron tunneling in Ru-modified His46Asp azurin. Coupling through the Cu ligands, **275–276**, 470

structural study and electron exchange kinetics of the bis(2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline)copper(II)/(I) couple in acetonitrile, **277**, 31

spectroscopic characterization and kinetic studies of a novel plastocyanin from the green alga *Ulva pertusa*, **283**, 184

Electron transfer metalloproteins

detection of two novel lowest lying singlet–triplet transitions of the C–S–C chromophore in L-methionine, **272**, 150

Electron tunneling

electron tunneling in Ru-modified His46Asp azurin. Coupling through the Cu ligands, **275–276**, 470

Electrophoresis

electrophoretic study of cationic cobalt(III) complexes with $[N_2O]$ and $[N_2O_2]$ Schiff base ligands, **280**, 295

Electroreduction

conductive polymeric films of *meso*-tetrakis[2,2]-*para*-cyclophanylmetalloporphyrins — spectroelectrochemistry and catalytic properties, **271**, 167

Electrospray mass spectrometry

an electrospray mass spectrometric and voltammetric study of horse heart cytochrome *c* in the presence of metal ions, **267**, 281

application of electrospray mass spectrometry to the characterisation of tertiary arsine ligands, **281**, 101

Electrospray study

an electrospray mass spectrometric investigation of gallium trihalide and indium trihalide solutions, **277**, 26

Electrostatics

the effect of surface charge on the reduction potential and heme propionate ionization of a monoheme cytochrome: spectroscopic and potentiometric studies of trifluoroacetylated cytochrome c_{551} from *Pseudomonas aeruginosa*, **275–276**, 256

Enamides

probing the nature of H_2 activation in catalytic asymmetric hydrogenation, **270**, 285

Enamine complexes

the preparation and synthetic potential of chlorovanadium(V and IV) complexes supported by enamines and bis(enamines), **267**, 229

Enantiomeric complexes

first solid state synthesis of the racemic compound and enantiomer of dinuclear Λ,Λ-di-μ-hydroxo-bis[bis((1*S*,2*S*)-1,2-*trans*-cyclohexanediamine)chromium(III)] and Δ,Δ-di-μ-hydroxo-bis-[bis((1*R*,2*R*)-1,2-*trans*-cyclohexanediamine)chromium(III)] bromides, **274**, 210

Enantioselective catalysis

enantioselective catalysis. XVI: regio- and enantioselectivity in nickel-catalysed cross-coupling reactions of allylic substrates with Grignard reagents, **269**, 34

the influence of the catalyst preparation protocol and silane structure on the rate and enantioselectivity of ansa-titanocene catalysed hydrosilation of prochiral ketones, **270**, 330

Enantioselectivity

enantioselective cleavage of activated amino acid esters promoted by chiral palladacycles, **280**, 57

ENDOR spectroscopy

proton ENDOR spectroscopy of cyanide-inhibited copper–zinc superoxide dismutase in randomly oriented samples, **273**, 131

Energy gap

electronic properties of some inorganic solids, **270**, 252
Enolate complexes
tetravalent niobium alkyls and enolate stabilized by anionic organic amides, **280**, 143
Enzyme activity
the immobilisation of proteins in carbon nanotubes, **272**, 261
Enzyme complexes
dynamic properties of bovine Cu,Zn superoxide dismutase from crystallographic data, **275–276**, 349
crystal structure, spectral and magnetic properties of a new (μ-acetate)(μ-alkoxide) dicopper(II) complex as a model for tyrosinase, **281**, 111
Enzyme reduction
reduction of *Coprinus macrorhizus* peroxidase by dithionite, **274**, 215
Epoxidation
synthesis and characterization of the new complex tris[(2-oxazolinyl)phenolato] manganese(III) and its use as a catalyst in oxidation reactions, **279**, 217
Equilibria
reinforced piperazine rings containing polyamines: metal complex equilibria and structural studies, **268**, 63
Equilibrium constants
thermodynamic study of complexation of alkali and alkaline-earth metal ions with benzene-containing cryptands, **278**, 6
thermodynamics of the binding of nitrogenous ligands to β-trifluoromethylcobinamide and β-cyanomethylcobinamide, **279**, 178
^{2}E relaxation
^{2}E relaxation of Cr(III) complexes in rigid and fluid media, **277**, 211
ESCA of humates
electron spectroscopy for chemical analysis investigation of the interaction of uranyl and calcium ions with humic acids, **273**, 234
Ester hydrolysis
kinetics and mechanism of ester hydrolysis by metal complexes of 2,6-diacetylpyridine dioxime, **273**, 167
Ether-phosphine complexes
reactivity and chemical exchange processes of bis(ether-phosphine) iridium(III) complexes, **269**, 13
Ethylene complexes
synthesis and reactivity of bis(ethylene)-phosphite complexes of molybdenum(0), **272**, 125
Ethylidene-bridged complexes
dialkyl-μ-ethylidene-μ-methylene-bis(pentamethylcyclopentadienyl)-dirhodium complexes: synthesis, spectra and decomposition reactions, **280**, 62
Europium complexes
phosphide complexes of divalent europium: synthesis and structural characterization of Eu[PPh_2]$_2$(*N*-MeIm)$_4$, **267**, 309
heteroleptic triple-decker (phthalocyaninato)(porphyrinato)-europium(III) complexes: synthesis and electrochemical study, **268**, 49
a new synthetic route to unsymmetrical bis(phthalocyaninato)-europium(III) complexes, **268**, 141
the dissociation kinetics of the ethylenediaminetetrakis(methylenephosphonato) complex of europium(III), **278**, 185
Europium(III) complexes
evaluation of solution structures of highly luminescent europium(III) chelates by using laser induced excitation of the $^7F_0 \rightarrow {}^5D_0$ transition, **267**, 63
EXAFS
EXAFS studies on Mn(III) tetraphenylporphyrin halides in Nafion, **268**, 231
probing the iron environment in desulforedoxin. EXAFS of oxidized and reduced states, **273**, 409
the structure of the azide coordinated superoxide dismutase of *Propionibacterium shermanii* investigated by X-ray structure analysis, extended X-ray absorption fine structure, Mössbauer and electron paramagnetic resonance spectroscopy, **275–276**, 65
X-ray absorption spectroscopic studies of the binding of ligands to FeMoco of nitrogenase from *Klebsiella pneumoniae*, **275–276**, 150
structural characterization of the zinc site in *Escherichia coli* L-threonine dehydrogenase using extended X-ray absorption fine structure spectroscopy, **275–276**, 215
Exchange interactions
exchange coupling across multiple-atom bridges: crystal structure and magnetic properties of a chloranilate bridged dicopper(II) complex, **274**, 111
Exchange rate constant
effect of pyridine on the alkaline isomerization of cytochrome *c*, **275–276**, 58

FAB mass spectrometry
chemistry of rare earth hypocoordinate oxocluster cations: formation of novel oxoacetate clusters, **281**, 201
Famotidine complexes
metallodrug complexes that mediate DNA and lipid damage via sulfite autoxidation: copper(II) famotidine and iron(III) bis(salicylglycine), **275–276**, 314
4f complexes
homodinuclear lanthanide complexes: Ln_2L_3 (H_2L = tetradentate Schiff bases). Magnetic properties (solid state) and spectroscopic studies (solution), **268**, 125
Fe–S clusters
spectroscopic investigations on the water-soluble fragment of the Rieske [2Fe–2S] protein from *Paracoccus denitrificans*, **275–276**, 493
FeMo cofactor complexes
catalytic reduction of acetylene in the presence of molybdenum and iron clusters, including FeMo cofactor of nitrogenase, **270**, 221
Ferrates
the kinetics and mechanism of the oxidation of 1,4-thioxane by potassium ferrate, **267**, 159
the kinetics and mechanism of oxidation of 3-mercaptopropionic acid, 2-mercaptoethanesulfonic acid and 2-mercaptobenzoic acid by potassium ferrate, **281**, 43
Ferrihydrite
reductive changes to polynuclear iron(III) clusters in iron-loaded human spleen tissue, **267**, 7
Ferrimagnetic properties
tuning up the T_c in Mn(II)Cu(II) bimetallic planes and design of molecular-based magnets, **278**, 159
Ferrioxamine B
application of the electrocatalytic reduction of nitric oxide mediated by ferrioxamine B to the determination of nitric oxide concentrations in solution, **273**, 316
Ferritin
reductive changes to polynuclear iron(III) clusters in iron-loaded human spleen tissue, **267**, 7
Ferrocene complexes
crystal and molecular structure of protonated (*N*-propyl)-aminomethyl ferrocene, a proton-sensitive redox-responsive fragment, **267**, 177
ferrocenyl-derived face-capping ligands: syntheses and molecular structures of [(FcTt)Cu]$_4$ and [FcTt]$_2$M (M = Fe, Co, Ni; FcTt$^-$ = ferrocenyltris((methylthio)methyl)borate), **270**, 543
a ferrocene–perfluoroarene molecular complex, **270**, 559
3-ferrocene-substituted 3-mercaptopropenales and their Schiff bases as ligands for transition metal complexes, **271**, 40
variable coordination modes for 1,1′-bis(diphenylphosphino)-ferrocene (dppf) mono- and di-chalcogenides or -oxides (dppfE and dppfE_2; E = O, S or Se) in copper(I) complexes. Crystal structure of [{Cu(dppf)}$_2$(μ-dppfS_2)](BF_4)$_2$ and [Cu(dppfSe_2)]BF_4, **277**, 163
synthesis of manganese tricarbonyl cationic complexes of ferrocenyl substituted arenes via a manganese tricarbonyl cation transfer reaction, **281**, 229

ferrocenyl-substituted imine ligands and their reactivity towards $Fe_2(CO)_9$; crystal structure of Fc–C(H)=C(H)–C(H)=N–C_6H_{11} and (η^4-R^1–C(H)=C(H)–C(H)=N–R^2)$Fe(CO)_3$ (R^1 = C_6H_5, R^2 = Fc; R^1 = Fc, R^2 = C_6H_{11}, R^1 = R^2 = Fc; Fc = (η^5-C_5H_5)Fe(η^5-C_5H_4)), **282**, 111
Ferrocenic polymers
new type of polymer containing 1,1′-ferrocenediyl unit in π-conjugated main chain. Preparation and redox behavior, **269**, 310
Ferrocenophanes
oligonuclear ferrocenolato and 1,1′-ferrocenediolato derivatives of phosphorus: synthesis, structure and electrochemical behaviour, **273**, 24
Ferrocenyl complexes
mono- and bis-alkynyl ruthenium(II) complexes containing the ferrocenyl moiety; crystal structure of *trans*-[Ru(C≡$CC_5H_4FeC_5H_5$)$_2$-($Ph_2PCH_2CH_2PPh_2$)$_2$] and electrochemical studies, **272**, 188
preparation and properties of ferrocenyl bimetallic compounds for non-linear optics, **279**, 243
Fischer–Tropsch mechanism
dialkyl-μ-ethylidene-μ-methylene-bis(pentamethylcyclopentadienyl)-dirhodium complexes: synthesis, spectra and decomposition reactions, **280**, 62
Five-coordinated complexes
crystal structure and properties of a new five-coordinate manganese superoxide dismutase mimic, **277**, 21
five-coordinate Salen(^{t}Bu) complexes of zinc, **277**, 157
molecular structure of a five-coordinate binuclear zinc complex, **279**, 249
organometallic polymers containing coordinatively saturated platinum(II) complexes as pendants, **281**, 141
Flavins
photosensitized electron transfer reactions of cytochrome c_4 from *Pseudomonas stutzeri* with flavins and methyl viologen, **272**, 109
Fluoride complexes
silyl reagents (Me_3Si–X) efficiently transfer X to $Ir(H)_2F(P^tBu_2Ph)_2$, **270**, 261
Fluorinated porphyrin complexes
spectroscopic characterization of fluorinated/hydrogenated mixed vesicles containing fluorinated Mn(III)-porphyrin, **272**, 274
Fluxional behaviour
crystal structure and fluxional behavior of novel Pd(II) allyl complexes containing a potentially tridentate S,N,S ligand: an example of selective *syn–anti* isomerism, **277**, 247
Fluxionality
stereochemically-rigid and fluxional pyrimidines within the bis-[Ru(hedta)(pym)$_2$]$^-$ complex, **268**, 287
the synthesis, molecular structure and fluxional properties of the 1-phenylpropyne complex [$WI_2(CO)${$Ph_2P(CH_2)PPh_2$}(η^2-MeC_2Ph)], **279**, 65
Force constants
Co–C force constants from resonance Raman spectra of alkylcobalamins: insensitivity to dimethylbenzylimidazole coordination, **270**, 392
Formation constants
potentiometric and computer studies of yttrium–EDTMP, **275–276**, 37
Formazan derivative complexes
iron and manganese complexes of benzothiazolylformazans, **280**, 30
D-Fructose complexes
unusual coordination behavior of D-fructose towards dimethyltin(IV): metal-promoted deprotonation of alcoholic OH groups in aqueous solutions of low pH, **274**, 167
FT-IR study
Fourier transform IR study of the interaction of [(η^5-C_5H_5)$_2$-$FeRu(CO)_4$] with hydroxylated alumina surfaces, **277**, 61
FT-Raman spectroscopy
the Co–CH_3 bond in Schiff base B_{12} models: influence of the *trans* and equatorial ligands as assessed by Fourier transform Raman spectroscopy, **275–276**, 90
FTS peptide complexes
metal ion–FTS nonapeptide interactions. II. Copper(II) complex equilibria and structures under physiological conditions, **283**, 211
Fullerenes
exohedral and endohedral complexation energies of complexes X@C_{70} between higher fullerene and rare-gas atoms (X = He, Ne, Ar, Kr, Xe), **272**, 33
simulation of the molecular and electronic structure of heterofullerenes $C_{55}Y_5$ (Y = Si, Ge, Sn, B, Al, N, P, SiH, GeH, SnH) and their η^5-π-complexes with Li by the MNDO method, **280**, 219
Fulvene complexes
synthesis and properties of the dichloro[tetramethyl(trifluoromethyl)-cyclopentadienyl]ruthenium dimer: X-ray structure of [M_2-(η^5-$C_5Me_4CF_3$)$_2Cl_2$(μ-Cl)$_2$] (M = Ru, Rh), **280**, 249
Functionalized rhodium complexes
organometallic Rh(I) and Rh(III) complexes with [2-(dimethylamino)ethyl]cyclopentadiene, **283**, 24

Gadolinium complexes
a ^{31}P and ^{1}H NMR relaxometric study of the interaction between adenosine triphosphate (ATP) and paramagnetic ions (Gd^{3+} and Mn^{2+}), **273**, 92
a tripod gadolinium(III) poly(amino carboxylate) relevant to magnetic resonance imaging: structural and dynamical ^{17}O NMR and ^{1}H NMRD studies, **275–276**, 15
Galactose oxidase
characterization of the active site of galactose oxidase and its active site mutational variants Y495F/H/K and W290H by circular dichroism spectroscopy, **275–276**, 175
Gallium complexes
complexation of gallium(III) by 2-hydroxynicotinic acid and 3-hydroxypicolinic acid in aqueous solution, **273**, 47
an electrospray mass spectrometric investigation of gallium trihalide and indium trihalide solutions, **277**, 26
synthesis and characterization of the tungsten–gallium heterometallic cluster $Na_2[W_3GaO_4(O_2CEt)_8]_2$, **277**, 130
electrochemical synthesis and characterisation of gallium, indium and thallium compounds with 1-hydroxopyridine-2-thione. The crystal structures of tris(1-oxopyridine-2-thionato)indium(III) and tris(1-oxopyridine-2-thionato)thallium(III), **281**, 70
Gas occlusion
synthesis and gas-occlusion properties of dinuclear molybdenum(II) dicarboxylates (fumarate, terephthalate, *trans-trans*-muconate, pyridine-2,5-dicarboxylate, and *trans*-1,4-cyclohexanedicarboxylate), **283**, 268
Gas-phase chemistry
gas-phase positive and negative ion chemistry of methyl hydroperoxide, **275–276**, 192
Gentamicin complexes
NMR studies of iron–gentamicin complexes and the implications for aminoglycoside toxicity, **273**, 85
Germane compounds
alkoxymesitylchlorogermanes and aryloxymesitylchlorogermanes, **277**, 98
Glass beads
a catalytic activity of glass beads, silica gels and anion-exchange resins modified with metal-porphines in oxidative reactions of ascorbic acid, **283**, 44
Glucamide complexes
multinuclear NMR study of the interactions between the La(III) complex of DTPA-bis(glucamide) and Zn(II) or borate, **268**, 249
Glucose oxidation
selective oxidation of glucose on bismuth-promoted Pd–Pt/C catalysts prepared from $NOct_4Cl$-stabilized Pd–Pt colloids, **270**, 95

Glutamic acid complexes
an infinite chiral single-helical structure formed in Cu(II)-L-/D-glutamic acid system, **283**, 105
Glutathione complexes
imidazole release from the antitumor-active ruthenium complex imidazolium *trans*-tetrachlorobis(imidazole)ruthenate(III) by biologically occurring nucleophiles, **267**, 137
interaction of [ReO(SNS)(S)] and [^{99m}TcO(SNS)(S)] mixed ligand complexes with glutathione: isolation and characterization of the product, **281**, 148
Glycolato complexes
a phosphane analog of nedaplatin: preparation and structural characterization of *cis*-bis(trimethylphosphane)(glycolato-*O*,*O'*)-platinum(II), **279**, 105
Glycylhistidyllysine complexes
Pd(gly-L-his-L-lys)Cl: solution structure and ternary complex formation with mono- and tetranucleotides, **273**, 31
Goethite
reductive changes to polynuclear iron(III) clusters in iron-loaded human spleen tissue, **267**, 7
Gold complexes
syntheses and spectroscopic characterizations of the compounds containing univalent coinage metal and tetraselenomolybdate. Crystal structure of $[MoSe_4(AuPPh_3)_2]$, **271**, 93
the synthesis, NMR and X-ray analysis of tris(1-ethyl-2-isopropylimidazol-5-yl)phosphine gold(I)-2,3,4,6-tetra-O-acetyl-1-thio-β-D-glucopyranoside, a bifunctional auranofin analogue, **274**, 137
toward a novel metal based chemotherapy against tropical diseases. 4. Synthesis and characterization of new metal-clotrimazole complexes and evaluation of their activity against *Trypanosoma cruzi*, **275–276**, 528
electronic absorption and MCD spectra for the $Pt(PPh_3)(AuPPh_3)_6^{2+}$ and $Pt(H)(PPh_3)(AuPPh_3)_7^{2+}$ cluster complexes, **278**, 113
binuclear diorganophosphinothioformamido complexes of gold(I): synthesis and crystal structure of $[Au(Ph_2PC(S)NPh)]_2$ and related systems, **281**, 64
gold(III) compounds as potential antitumor agents: cytotoxicity and DNA binding properties of some selected polyamine–gold(III) complexes, **281**, 90
a single-crystal electron paramagnetic resonance, ^{13}C and ^{1}H electron nuclear double resonance study of tetra-*n*-butylammonium-bis(1,2-dicyanoethylene-1,2-dithiolato)aurate(II), $[(n\text{-}C_4H_9)_4N]_2$-$[Au^{II}(mnt)_2]$, **281**, 160
interactions between $[AuX_4]^-$ (X = Cl, CN) and cytosine and guanine model nucleobases: salt formation with (hemi-) protonated bases, coordination, and oxidative degradation of guanine, **283**, 223
Gold compounds
metal cluster topology. 19. Beyond the auride ion: triangulated gold networks and ethane-like structural units in binary and ternary alkali metal gold intermetallics, **277**, 202
Group 10 metal complexes
synthesis, cyclic voltammetry and $^{31}P\{H\}$ NMR spectra of $[M_3E_2L_3]^{2+}$ clusters; M = Ni, Pd or Pt; E = S, Se or Te; L = bis(diphenylphosphino)methane, 1,2-bis(diphenylphosphino)-ethane, 1,3-bis(diphenylphosphino)propane, or 1,4-bis-(diphenylphosphino)butane, **281**, 174
Group theory
chemical applications of topology and group theory. 30. Distortions from idealized symmetrical coordination polyhedra by lifting molecular orbital degeneracies, **270**, 68
Guanine complexes
interactions between $[AuX_4]^-$ (X = Cl, CN) and cytosine and guanine model nucleobases: salt formation with (hemi-) protonated bases, coordination, and oxidative degradation of guanine, **283**, 223
Haem–peptide complexes
synthesis, characterization and stereoselective catalytic oxidations of chelated deuterohaemin-glycyl-L-histidine complexes, **273**, 339
Hafnium complexes
trimethylphosphine halide complexes of titanium, zirconium and hafnium in oxidation states IV, III and II. Crystal structure of $MI_4(PMe_3)_n$ (M = Zr, n = 3; M = Hf, n = 2.5), $M_2X_6(PMe_3)_4$ (M = Hf, Ti, X = I; M = Zr, X = Br) and $(\mu\text{-}\eta^6,\eta^6\text{-}C_6H_6)$-$Hf_2I_4(PMe_3)_4 \cdot C_6H_6$, **271**, 180
Halide complexes
an electrospray mass spectrometric investigation of gallium trihalide and indium trihalide solutions, **277**, 26
preparation of tetrabutylammonium octa-μ_3-bromohexa(trifluoromethanesulfonato)octahedro-hexamolybdate(2−), $(Bu_4N)_2$-$[Mo_6Br_8^i(CF_3SO_3)_6^a]$ and other derivatives containing the $Mo_6Br_8^{4+}$ core, **278**, 241
the *trans* influence of F, Cl, Br and I ligands in a series of square-planar Pd(II) complexes: relative affinities of halide anions for the metal centre in *trans*-$[(Ph_3P)_2Pd(Ph)X]^+$, **280**, 87
Halido complexes
transition metal complexes with sulfur ligands. Part CXXVII. Azido, halido and nitrido ruthenium complexes with sulfur-rich coordination spheres, **269**, 63
Haloarene complexes
bonding of halogenated arenes in photogenerated (arene)$M(CO)_5$ complexes (M = Cr, Mo, W), **278**, 202
Halogen complexes
structures and electronic states of the electron donor–acceptor (EDA) complexes: trimethylamine–halogen complex $(CH_3)_3N\cdots F_2$, **281**, 85
Haptotropic rearrangements
ricochet inter-ring haptotropic rearrangement of σ-methyl-(η^5-indenyl)chromium tricarbonyls. Experimental kinetic and theoretical DFT study, **280**, 328
Hard and soft
metal ion stabilities correlate with electron affinity rather than hardness or softness, **283**, 30
HDS modelling
synthesis and characterization of heterobimetallic complexes containing C–S cleaved thiophenes, **272**, 55
Heavy metal complexes
electronic absorption spectroscopy for the investigation of the solution binding of gold, palladium, mercury and silver salts to the lower rim substituted calix[4]arenes 25,26,27,28-(2-*N*,*N*-dimethyldithiocarbamoylethoxy)calix[4]arene and 25,26,27,28-(2-methylthioethoxy)calix[4]arene, **270**, 216
Helical structure
an infinite chiral single-helical structure formed in Cu(II)-L-/D-glutamic acid system, **283**, 105
Helicate complexes
dinuclear metal complexes of Cd(II), Zn(II) and Fe(II) with triple-helical structure and predetermined chirality, **271**, 36
Heme
reversible formation of high-valent-iron-oxo porphyrin intermediates in heme-based catalysis: revisiting the kinetic model for horseradish peroxidase, **275–276**, 98
structure and reactivity of reconstituted myoglobins: interaction between protein and polar side chain of chemically modified hemin, **275–276**, 159
Heme proteins
isolation and characterization of cytochrome c_2 from *Rhodopseudomonas palustris*, **269**, 125
photosensitized electron transfer reactions of cytochrome c_4 from *Pseudomonas stutzeri* with flavins and methyl viologen, **272**, 109
the orientation of the iron axial ligands in the low-potential cytochrome c_{549} from *Synechocystis sp.* PCC 6803 studied by NMR, **273**, 196

Hemoprotein models
hemoprotein models based on a covalent helix–heme–helix sandwich 4. Discrimination of paramagnetic Fe(III)–Mimochrome I Δ and Λ isomers by NMR spectroscopy, **278**, 76
Hemoproteins
miniaturized hemoproteins: design, synthesis and characterization of mimochrome II, **275–276**, 301
Hemosiderin
reductive changes to polynuclear iron(III) clusters in iron-loaded human spleen tissue, **267**, 7
Herbicide complexes
Cu^{II}–herbicide complexes: structure and bioactivity, **272**, 24
metal complexes of Imazapyr, a herbicide provided with efficient metal-chelating ability: crystal structure of the cobalt(III) and manganese(II) complexes, **272**, 68
Heterobimetallic complexes
synthesis and crystal structures of complexes containing an Fe→Cd donor–acceptor bond formed by reaction of *trans*-$(CO)_3Fe(Ph_2Ppym)_2$ (Ph_2Ppym = 2-(diphenylphospino)pyrimidine) with CdX_2 (X = Cl, Br, I, SCN, ClO_4), **268**, 177
synthesis and characterization of heterobimetallic complexes containing C–S cleaved thiophenes, **272**, 55
synthesis and X-ray crystal determination of four novel lanthanide(III) oxydiacetate complexes, **281**, 18
synthesis of manganese tricarbonyl cationic complexes of ferrocenyl substituted arenes via a manganese tricarbonyl cation transfer reaction, **281**, 229
Heterocyclic amine complexes
solvent-induced reaction of rhodium(III) halide with N-heterocyclic ligands. Crystal structure of *trans*-dichlorotetrakis(*N*-methylimidazole)rhodium(III) chloride dihydrate, **278**, 209
Heterocyclic thione complexes
synthesis and characterization of copper(I) complexes with triphenylphosphine and heterocyclic thione ligands: the crystal structure of (thiazolidine-2-thione)(bis-triphenylphosphine)copper(I) chloride, **271**, 243
electrochemical synthesis and characterization of indium pyrimidine-2-thiolate complexes. Molecular structure of tris(pyrimidine-2-thiolato)indium(III) and tris[(5-ethyl-4,6-dimethyl)pyrimidine-2-thiolato]indium(III), **274**, 131
complexes of heterocyclic thiones and group twelve metals. Part 1. Preparation and characterisation of 1:1 complexes of mercury(II) halides with 1-methylimidazoline-2(3H)-thione: the crystal structure of [(μ_2-dibromo) bis(*trans*{(bromo)(1-methyl-imidazoline-2(3H)-thione)}mercury(II))] at 160 K, **277**, 14
silver(I) complexes with heterocyclic thiones and tertiary phosphines as ligands. Part 3. Solution behaviour of mononuclear complexes of silver(I) nitrate. The crystal structure of bis(triphenylphosphino)bis(benzoxazoline-2-thione)silver(I) nitrate, **281**, 121
Heterodimetallic complexes
the synthesis of substituted bis[(diarylphosphinomethyl)cyclopentadienyl]zirconocene dichloride complexes for the preparation of heterodimetallic complexes containing early/late transition metal combinations, **270**, 446
Heterodinuclear complexes
dinuclear $Cu^{II}M^{II}$ complexes of a phenol-based macrocycle with $N(amine)_2O_2$ and $N(imine)_2O_2$ metal-binding sites: the effect of chloride ligation upon the site selectivity of metal ions, **283**, 72
Heterofullerenes
simulation of the molecular and electronic structure of heterofullerenes $C_{55}Y_5$ (Y = Si, Ge, Sn, B, Al, N, P, SiH, GeH, SnH) and their η^5-π-complexes with Li by the MNDO method, **280**, 219
Heteroleptic complexes
a new synthetic route to unsymmetrical bis(phthalocyaninato)-europium(III) complexes, **268**, 141
Heterometallic complexes
synthesis and characterization of the tungsten–gallium heterometallic cluster $Na_2[W_3GaO_4(O_2CEt)_8]_2$, **277**, 130
Heteronuclear complexes
structures, dynamic behaviour in solution and cross-coupling reactions of the α-palladiated (η^6-alkylarene)tricarbonylchromium complexes containing a palladium–chromium bond, **280**, 71
Heteropolyanions
preparation and characterization of a new heteropolyoxofluorotungstate anion $[InW_{17}O_{56}F_6NaH_4]$, **268**, 329
preparation and characterization of two novel heteropolyfluorooxotungstate anions $[NaCoW_{11}O_{43}FH_{12}]^{6-}$ and $[NaNiW_{11}O_{43}FH_{12}]^{6-}$, **282**, 136
Heteropolyfluorotungstate
preparation and characterization of a new heteropolyoxofluorotungstate anion $[InW_{17}O_{56}F_6NaH_4]$, **268**, 329
Heteropolymolybdates
formation and conversion of yellow heteropoly complexes in a Mo(VI)–Se(IV), Te(IV) system, **281**, 214
Heteropolyoxotungstates
preparation and characterization of the catalytically active heteropolyoxotungstate anion $[Co^{3+}ZnW_{11}O_{40}H_2]^{7-}$, **278**, 91
Hexatungstate clusters
hydrothermal synthesis and characterization of mixed-valence hexatungstates: crystal structures of $[(C_2H_5)_4N]_3[W^VW_5^{VI}O_{19}]\cdot 0.5H_2O$ and $[H_3N(CH_2)_2NH_3]_2[W^VW_5^{VI}O_{19}]\cdot[H_2N(CH_2)_2NH_2]Cl\cdot 8H_2O$, **277**, 69
High spin ions
angular overlap calculations of the spin Hamiltonian parameters of transition metal ions in low symmetry environments. High spin iron(II), iron(III) and manganese(III), **274**, 90
Histidine complexes
solution behaviour and zinc complexation of tripeptides with cysteine and/or histidine at both termini, **272**, 115
Histidine tetrapeptides
interaction of Hg(II) with tetrapeptides containing cysteinyl and histidinyl residues, **272**, 7
^1H NMR
the effect of net charge and π-backbonding contribution of ML_5 fragments containing Fe(II), Ru(II), Os(II), Co(III) and Rh(III) on the ^1H NMR shifts of coordinated N-heterocycles, **267**, 11
^1H and ^{11}B NMR evidence for specific binding of borate ion to cytochrome *c*, **273**, 388
Holmium complexes
the preparation and X-ray structure of $[Ho(oxam)_3(H_2O)_3]_4\cdot 2.75H_2O$, **267**, 299
Human carbonic anhydrase II complexes
a ternary complex of carbonic anhydrase: X-ray crystallographic structure of the adduct of human carbonic anhydrase II with the activator phenylalanine and the inhibitor azide, **275–276**, 295
Humic acid complexes
electron spectroscopy for chemical analysis investigation of the interaction of uranyl and calcium ions with humic acids, **273**, 234
Hybrid P,N ligand complexes
2-(diphenylphosphino)pyridine (dppy) as monodentate or bridging ligand in homoleptic silver(I) complexes. Crystal structures of $[Ag(\eta^1\text{-dppy})_4][ClO_4]$ and $[Ag_2(\eta^1\text{-dppy})(\mu\text{-dppy})_2][ClO_4]_2$, **277**, 111
Hydration
magnesium activation of nuclease enzymes — the importance of water, **275–276**, 24
Hydrazido complexes
reactivity of molybdenum complexes containing both NHNPhR and NNPhR ligands (R = Ph, Me) toward hydridotris(pyrazolyl)-borate, Tp and hydridotris(3,5-dimethylpyrazolyl)borate, Tp*, ligands. X-ray crystal structure of $[TpMo(NNMePh)_2Cl]$, **268**, 199
Hydrazine

the cations $[ICNI]^+$ and $[H_3N{-}NH_3]^{2+}$ — natural bond orbital analysis and some valence bond considerations, **269**, 1

Hydrazinium cations

the cations $[ICNI]^+$ and $[H_3N{-}NH_3]^{2+}$ — natural bond orbital analysis and some valence bond considerations, **269**, 1

Hydrazone complexes

antitumour copper(II) salicylaldehyde benzoylhydrazone (H_2sb) complexes: physicochemical properties and the single-crystal X-ray structures of $[\{Cu(H_2sb)(CCl_3CO_2)_2\}_2]$ and $[\{Cu(Hsb)(ClO_4)(C_2H_5OH)\}_2]$ and the related salicylaldehyde acetylhydrazone (H_2sa) complex, $[Cu(Hsa)Cl(H_2O)]\cdot H_2O$, **267**, 27

ligand exchange reactions of $ReCl_4(PPh_3)_2$, $ReOCl_3(PPh_3)_2$ and $NiCl_2(PPh_3)_2$ with acetylacetone thiobenzoylhydrazone. An unexpected template reaction at the ReO^{3+} core. Molecular structures of the resulting oxorhenium(V) and nickel complexes, **277**, 219

Hydride complexes

thermal and photochemical substitution reactions of $CpRe(PPh_3)_2H_2$ and $CpRe(PPh_3)H_4$. Catalytic insertion of ethylene into the C–H bond of benzene, **270**, 77

elimination of benzene from a benzyloxy ligand to form a carbonyl group on a tungsten-phosphine center, **270**, 202

silyl reagents ($Me_3Si{-}X$) efficiently transfer X to $Ir(H)_2F(P^tBu_2Ph)_2$, **270**, 261

hydride complexes of platinum(II) containing unsymmetrical di(phosphine) ligands: synthesis, characterization and evidence for pairwise addition of hydrogen based on parahydrogen induced polarization, **270**, 363

extracting absolute titanium–alkyl and –hydride bond enthalpies from relative D(TiR(H)) in $(silox)_2(^tBu_3SiNH)TiR$: electronegativity and *ECT* models, **270**, 414

a mechanistic study of thermal C–H reductive elimination from a six-coordinate d^6 iridium complex, **270**, 537

transition metal complexes with sulfur ligands. Part CXXV. Synthesis and characterization of hydrido and chloro complexes with rhodium sulfur cores, **272**, 211

NMR characterization of four new isomers of $H(\mu\text{-}H)Os_3(CO)_{10}$(phosphine), **275–276**, 521

NMR evidence for formation of new alcohol rhenium complexes as intermediates in ionic hydrogenations of carbonyl groups with systems composed of $ReH_2(NO)(CO)(PR_3)_2$ (R = Pr^i, CH_3, OPr^i) and CF_3COOH, **278**, 122

a simple and convenient synthesis of *cis*/*trans*-$RuH_2(Ph_2PCH_2PPh_2)_2$ and of *trans*-$RuHCl(Ph_2PCH_2PPh_2)_2$, **278**, 226

intermolecular Re–H···H–X hydrogen bonding (X = N, C) involving $ReH_5(PPh_3)_3$, **280**, 26

ammonia ruthenium complexes for the access to new hydrido, carbonyl and isocyanide ruthenium-alkynyl derivatives, **280**, 118

electrophilic abstractions on $Os(H)_2X(NO)L_2$: selective access to new unsaturated cationic Os(II) hydrides, **280**, 125

in situ IR and NMR study of the interactions between proton donors and the Re(I) hydride complex $[\{MeC(CH_2PPh_2)_3\}Re(CO)_2H]$. ReH···H bonding and proton-transfer pathways, **280**, 302

synthesis and characterization of arene, mono- and dihydrido-arene, monohydrido-cyclopentadienyl, and phosphite complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$, **281**, 153

β-Hydride elimination

reversible β-hydride elimination of amine ligands forming stable *cis*-η^2-iminium hydride Os^{II} complexes, **270**, 337

Hydrido complexes

structures of $HRh(PMe_2Ph)_4$ and $HRh(PPh_2Me)_4$, **269**, 162

intramolecular protonation and other mechanisms for substitution reactions of hydrido(thiolato)– and di(thiolato)–ruthenium(II) phosphine complexes, **280**, 75

Hydridotris(3,5-dimethylpyrazolyl)borate

reactivity of molybdenum complexes containing both NHNPhR and NNPhR ligands (R = Ph, Me) toward hydridotris(pyrazolyl)-borate, Tp and hydridotris(3,5-dimethylpyrazolyl)borate, Tp*, ligands. X-ray crystal structure of $[TpMo(NNMePh)_2Cl]$, **268**, 199

Hydridotris(pyrazolyl)borate

reactivity of molybdenum complexes containing both NHNPhR and NNPhR ligands (R = Ph, Me) toward hydridotris(pyrazolyl)-borate, Tp and hydridotris(3,5-dimethylpyrazolyl)borate, Tp*, ligands. X-ray crystal structure of $[TpMo(NNMePh)_2Cl]$, **268**, 199

Hydroformylation

the hydrogenation and hydroformylation of alkenes as catalyzed by polymer-anchored rhodium trichloride under water gas shift reaction conditions, **270**, 345

Hydrogen

reactivity studies on cationic non-hydridic triruthenium carbonyl clusters. Reactions of $[Ru_3(\mu_3\text{-ampy})(CO)_{10}][BF_4]$ with hydrogen, triethylsilane and triphenylstannane, **268**, 131

Hydrogenation

the hydrogenation and hydroformylation of alkenes as catalyzed by polymer-anchored rhodium trichloride under water gas shift reaction conditions, **270**, 345

Hydrogen bond complexes

polymeric and dimeric magnetic properties of square planar Cu(II) species controlled by hydrogen bond networks: $[Cu^{II}\{OCO\text{-}2,6\text{-}(CH_3CONH)_2C_6H_3\}_2(H_2O)_2]\cdot nH_2O$ (n = 1, 4), **275–276**, 43

Hydrogen bonding

effect of hydrogen bonding on coordination in *trans*-di(*Z*-aminobenzoato-*O*)di(1,3-diaminopropane-*N*,*N*′)copper(II) (*Z* = 2, 3 or 4) complex units, **268**, 21

intermolecular Re–H···H–X hydrogen bonding (X = N, C) involving $ReH_5(PPh_3)_3$, **280**, 26

in situ IR and NMR study of the interactions between proton donors and the Re(I) hydride complex $[\{MeC(CH_2PPh_2)_3\}Re(CO)_2H]$. ReH···H bonding and proton-transfer pathways, **280**, 302

structure and properties of tetraphenylporphinate iron(III) complexes with an intramolecular NH···S benzenethiolate or NH···O phenolate hydrogen bond, **283**, 91

Hydrogen peroxide

oxidations in the H_2O_2/V^V/AcOH system. Evidence of vanadium(V) complexes with superoxide anion and singlet dioxygen molecule as ligands, **275–276**, 453

Hydrogensulfate complexes

synthesis, X-ray structure, spectroscopic and magnetic properties of [bis(4-amino-3,5-bis(pyridin-2-yl)-1,2,4-triazole-*N*′,*N*1)-(aqua)copper(II)]bis(hydrogensulfate), **268**, 37

Hydrolysis

partial hydrolysis of Ti(III) and Ti(IV) chlorides in the presence of $[PPh_4]Cl$, **267**, 271

hydrolysis of nerve agent simulants by synergistic effects of tetraminecobalt(III) and microemulsions, **269**, 287

hydrolysis of phosphate esters catalyzed by copper(II)-triamine complexes. The effect of triamine ligands on the reactivity of the copper(II) catalysts, **283**, 51

Hydroperoxo complexes

an ab initio MO-LCAO investigation of the electronic structure of organic hydroperoxides and platinum(II) hydroperoxo complexes: a contribution to the knowledge of the mechanism of olefin epoxidation, **270**, 479

Hydroperoxyl radicals

the oxidation of nickel^{II}(1,4,8,11-tetraazacyclotetradecane) by hydroperoxyl radicals: a pulse radiolysis study, **273**, 266

Hydrophilic complexes

synthesis, UV–visible and electrochemical studies of lipophilic and hydrophilic lanthanide(III) bis(porphyrinates), **275–276**, 182

Hydrophobic interaction

spectroscopic characterization of coaxially and π–π stacked binuclear copper(II) complexes of ω,ω′-bis(pyridine-2-carboxamido)-alkanes in acidic solution, **283**, 61

Hydrophobic vitamin B_{12}

hydrophobic vitamin B_{12}. Part 15. Carbon-skeleton rearrangement reactions mediated by hydrophobic vitamin B_{12} covalently bound to an anionic lipid species in aqueous media, **273**, 299

Hydrosilation

the influence of the catalyst preparation protocol and silane structure on the rate and enantioselectivity of ansa-titanocene catalysed hydrosilation of prochiral ketones, **270**, 330

Hydrosulfido-bridged complexes

syntheses, structures and properties of hydrosulfido-bridged diiridium and dirhodium complexes, $[Cp^*MCl(\mu_2\text{-}SH)_2MCp^*Cl]$ and $[Cp^*M(\mu_2\text{-}SH)_3MCp^*]^+$ (M = Ir, Rh; $Cp^* = \eta^5\text{-}C_5Me_5$), **267**, 73

Hydrothermal synthesis

hydrothermal synthesis and characterization of mixed-valence hexatungstates: crystal structures of $[(C_2H_5)_4N]_3[W^VW^{VI}_5O_{19}]\cdot 0.5H_2O$ and $[H_3N(CH_2)_2NH_3]_2[W^VW^{VI}_5O_{19}]\cdot[H_2N\text{-}(CH_2)_2NH_2]Cl\cdot 8H_2O$, **277**, 69

Hydroxamate complexes

siderophore analogues. A new macrocyclic *bis*-(amine, amide, hydroxamate) ligand. Synthesis, solution chemistry, electrochemistry and molecular mechanics calculations for the iron complex, **278**, 51

Hydroxamic acid complexes

synthesis and characterization of two novel monooxovanadium(V) complexes with bidentate benzohydroxamate ligand, **282**, 149

Hydroxo-bridged complexes

first solid state synthesis of the racemic compound and enantiomer of dinuclear Λ,Λ-di-μ-hydroxo-bis[bis((1*S*,2*S*)-1,2-*trans*-cyclohexanediamine)chromium(III)] and Δ,Δ-di-μ-hydroxo-bis-[bis((1*R*,2*R*)-1,2-*trans*-cyclohexanediamine)chromium(III)] bromides, **274**, 210

a series of $[Cr_2(\mu\text{-}OH)_n(\mu\text{-}RCO_2)_{3-n}(bispma)_2]$ type chromium(III) dinuclear complexes ($n = 1, 2$) with tridentate bis(2-pyridylmethyl)amine (bispma): synthesis, crystal structure, and magnetic properties, **278**, 15

Hydroxybenzoquinone complexes

assembly of a mixed-valence Cu(I/II) system coupled by multiple hydrogen bonding through tetrahydroxybenzoquinone, **268**, 317

Hydroxy compounds

relative reactivities of hydroxy compounds with short-lived manganese(V), **281**, 235

Hydroxyferrocenes

oligonuclear ferrocenolato and 1,1′-ferrocenediolato derivatives of phosphorus: synthesis, structure and electrochemical behaviour, **273**, 24

Hydroxylated alumina complexes

Fourier transform IR study of the interaction of $[(\eta^5\text{-}C_5H_5)_2\text{-}FeRu(CO)_4]$ with hydroxylated alumina surfaces, **277**, 61

Hydroxymethylated amino acid complexes

copper(II) complexes of (*R*,*S*)-α-hydroxymethylornithine and its N^δ-benzoyl derivative, **269**, 279

Hydroxynicotinic acid complexes

complexation of gallium(III) by 2-hydroxynicotinic acid and 3-hydroxypicolinic acid in aqueous solution, **273**, 47

Hydroxypicolinic acid complexes

the coordination mode of 3-hydroxypicolinic acid in oxorhenium(V) complexes. Crystal and molecular structures of $(n\text{-}Bu_4N)\text{-}[ReOCl_3(HO\text{–}C_5H_3N\text{–}CO_2)]$ and $[ReOCl(HO\text{–}C_5H_3N\text{–}CO_2)_2]$, **269**, 235

complexation of gallium(III) by 2-hydroxynicotinic acid and 3-hydroxypicolinic acid in aqueous solution, **273**, 47

Hydroxypyridine complexes

4- and 3-hydroxypyridine complexes of pentacyanoferrate(II, III), **267**, 81

Hyperfine shift

isolation and characterization of cytochrome c_2 from *Rhodopseudomonas palustris*, **269**, 125

Imidazolate complexes

synthesis and characterization of a neutral, low spin iron(III) complex of a hexadentate tripodal ligand containing three imidazolate arms. Use as a dinucleating agent, **278**, 197

Imidazole bridge complexes

general synthesis of 'salicylaldehyde half-unit complexes': structural determination and use as synthon for the synthesis of dimetallic or trimetallic complexes and of 'self-assembling ligand complexes', **274**, 73

Imidazole-carboxylic acid complexes

oxovanadium(IV) complexes of imidazole-4-acetic, imidazole-4,5-dicarboxylic and pyrazole-3,5-dicarboxylic acids, **268**, 297

Imidazole complexes

synthesis, spectral and structural characterization of the zinc carboxylate [Zn(2-quinolinecarboxylato)$_2$(1-methylimidazole)$_2$], **269**, 283

Imidazole-type ligands

metal ion-coordinating properties of imidazole and derivatives in aqueous solution: interrelation between complex stability and ligand basicity, **280**, 50

Imidazol-pyridine complexes

spin crossover in bis(2,6-bis(benzimidazol-2-yl)pyridine) iron(II) tetraphenylborate, **278**, 190

Imidocarbyne bridged complexes

'activated cyanide', **269**, 23

Imido complexes

rhenium(VII) and (V) imido and oxo complexes containing 1,4,7-triazacyclononane: synthesis, electrochemistry and crystal structures, **272**, 176

synthesis of chloro and methyl imido cyclopentadienyl molybdenum and tungsten complexes. X-ray molecular structures of $[WCp^*Cl_3(N^tBu)]$, $[MoCp^*ClMe_2(N^tBu)]$ and $[WCp^*ClMe_2(N^tBu)]$, **273**, 270

synthesis and molecular structure of the molybdate-bridged cyclic imido complex $[Mo(S_2CNEt_2)_2(NR)(\mu\text{-}MoO_4)]_2$ (R = 2,6-$^iPr_2C_6H_3$), **274**, 243

the reactivity of the $[Re{\equiv}O]^{3+}$ core toward aromatic 1,2-diamines: the formation of rhenium(V)-imido complexes, **278**, 96

the synthesis, characterization and X-ray crystal structures of the *fac-cis* and *mer-trans* isomers of $Re(NEt)Cl_3(PMe_3)_2$, **278**, 245

Imine complexes

additions of carbon nucleophiles to acyclic imine complexes of the chiral rhenium Lewis acid $[(\eta^5\text{-}C_5H_5)Re(NO)(PPh_3)]^+$: controlling factors in 1,3-asymmetric induction and syntheses of non-racemic organic amines, **269**, 167

ferrocenyl-substituted imine ligands and their reactivity towards $Fe_2(CO)_9$; crystal structure of Fc–C(H)=C(H)–C(H)=N–C_6H_{11} and $(\eta^4\text{-}R^1\text{–}C(H)=C(H)\text{–}C(H)=N\text{–}R^2)Fe(CO)_3$ ($R^1 = C_6H_5$, R^2 = Fc; R^1 = Fc, $R^2 = C_6H_{11}$, $R^1 = R^2$ = Fc; Fc = $(\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4)$), **282**, 111

α-Imino-acidato complexes

preparation and molecular structure of (*N*-benzyl-1,2-didehydroglycinato) bis(1,10-phenanthroline)ruthenium(II) chloride: [Ru(*N*-Bn-gly-H_2)(phen)$_2$]Cl, **267**, 329

Iminoacyl complexes

synthesis, characterization, and reactivity of organometallic Zr(IV) carboxylate complexes, **270**, 527

Imino esters

oxidation of amino esters to imine complexes with chiral high-valent oxoruthenium porphyrins, **273**, 142

Iminophosphoranyl complexes

substitution reactions of $CpRu(PPh_3)_2Cl$ with PPh_2H and dppa, **269**, 157

Immobilisation
the immobilisation of proteins in carbon nanotubes, **272**, 261
Indene
ricochet inter-ring haptotropic rearrangement of σ-methyl-(η^5-indenyl)chromium tricarbonyls. Experimental kinetic and theoretical DFT study, **280**, 328
Indium complexes
structure of an indium(III) complex with a 15-membered macrocycle, dioxopentaazacyclopentadecanetriacetate, **271**, 207
electrochemical synthesis and characterization of indium pyrimidine-2-thiolate complexes. Molecular structure of tris(pyrimidine-2-thiolato)indium(III) and tris[(5-ethyl-4,6-dimethyl)pyrimidine-2-thiolato]indium(III), **274**, 131
an electrospray mass spectrometric investigation of gallium trihalide and indium trihalide solutions, **277**, 26
electrochemical synthesis and characterisation of gallium, indium and thallium compounds with 1-hydroxopyridine-2-thione. The crystal structures of tris(1-oxopyridine-2-thionato)indium(III) and tris(1-oxopyridine-2-thionato)thallium(III), **281**, 70
Induced circular dichroism
the synthesis and spectroscopic characterization of a chiral zinc(II) porphyrin dimer containing an amino acid bridge: model complexes for the induced circular dichroism of porphyrin protein supramolecules, **272**, 295
Inosine complexes
substitution of inosine for chloride in $[Pt_2(hdta)Cl_2]^{2-}$ ($hdta^{4-}$ = 1,6-hexanediamine-*N*,*N*,*N*′,*N*′-tetraacetate), **271**, 124
Insulin-mimetic complexes
insulin-mimetic vanadyl–dithiocarbamate complexes, **283**, 175
π–π Interaction
spectroscopic characterization of coaxially and π–π stacked binuclear copper(II) complexes of ω,ω′-bis(pyridine-2-carboxamido)-alkanes in acidic solution, **283**, 61
Intercalation
intercalation into calf thymus DNA of 2-(4-arylphenyl)imidazo-[4,5-f]-[1,10]phenanthroline (aryl = -OMe, -NMe_2 or -NO_2) in its bipyridyl ruthenium(II) complex, **283**, 17
Intercalation compounds
X-ray absorption fine structure study of the atomic and electronic structure of molybdenum disulfide intercalation compounds with transition metals, **280**, 211
Intermetallic compounds
synthesis and characterization of electric and magnetic properties of the new intermetallic compounds $M_3(GaTe_3)_2$ (M = Cr, Fe, Co), **268**, 271
metal cluster topology. 19. Beyond the auride ion: triangulated gold networks and ethane-like structural units in binary and ternary alkali metal gold intermetallics, **277**, 202
Intramolecular hydrogen bond
role of the intramolecular hydrogen bond and ligand rigidity in the complexation of trifluoroacetylcycloalkanones with lanthanides: novel strategy for the design of organic ligands of high selectivity, **267**, 201
Iodine complexes
the reaction of cobalt powder with tetraiodo{1,2-bis(dibenzylphosphino)ethane} to form 1,2-bis(dibenzylphosphino)-ethane cobalt diiodide, $Co(dBzP_2)I_2$; and the X-ray crystal structure of the diphosphinodioxide complex, $Co\{dBzP_2(O)_2\}I_2$, **282**, 25
Iodine-doped complexes
mixed valence, tetradisordered structure of iodine-doped bismuth(III) diphthalocyanine, **281**, 195
Iodotrimethylsilane
trimethylsilyl iodide as a halide exchange reagent in inorganic chemistry: synthesis and structure of $Cp^*_2ThI_2$ ($Cp^* = \eta^5$-C_5Me_5), **274**, 232
Ionic hydrogenation
NMR evidence for formation of new alcohol rhenium complexes as intermediates in ionic hydrogenations of carbonyl groups with systems composed of $ReH_2(NO)(CO)(PR_3)_2$ (R = Pr^i, CH_3, OPr^i) and CF_3COOH, **278**, 122
Ionspray mass spectrometry
characterisation of covalent copper and manganese organometallic complexes with Schiff bases by ionspray mass spectrometry, **275–276**, 462
Iridium complexes
syntheses, structures and properties of hydrosulfido-bridged diiridium and dirhodium complexes, $[Cp^*MCl(\mu_2\text{-}SH)_2MCp^*Cl]$ and $[Cp^*M(\mu_2\text{-}SH)_3MCp^*]^+$ (M = Ir, Rh; $Cp^* = \eta^5$-C_5Me_5), **267**, 73
reactivity and chemical exchange processes of bis(ether-phosphine) iridium(III) complexes, **269**, 13
silyl reagents (Me_3Si–X) efficiently transfer X to $Ir(H)_2F(P^tBu_2Ph)_2$, **270**, 261
probing the nature of H_2 activation in catalytic asymmetric hydrogenation, **270**, 285
model reactions of a carbonylation catalyst: phosphite induced migratory CO insertion in $[MeIr(CO)_2I_3]^-$, **270**, 382
a mechanistic study of thermal C–H reductive elimination from a six-coordinate d^6 iridium complex, **270**, 537
synthesis and characterization of heterobimetallic complexes containing C–S cleaved thiophenes, **272**, 55
hydrogen site exchange in $CpIr(PR_3)H_3^+$ complexes, **272**, 95
oxidative addition of methyl iodide and iodine to new binuclear rhodium(I) and iridium(I) compounds containing diaminoanthraquinonate-bridging ligands. Crystal structure of $[Rh_2(\mu\text{-}1,4\text{-}DA)(CO)_2(PPh_3)_2]$ (1,4-H_2DA = 1,4-diaminoanthraquinone), **274**, 15
syntheses, spectroscopic characterization and X-ray crystal structures of $[HP\text{-}i\text{-}Pr_3]_2[IrCl_6]$ (**1**), $[HP\text{-}i\text{-}Pr_3]_2[Ir_2Cl_8(P\text{-}i\text{-}Pr_3)]$ (**2**) and $[HP\text{-}i\text{-}Pr_3][Ir_2Cl_7(P\text{-}i\text{-}Pr_3)_2]$ (**3**): three possible precursors of the complex $Ir(H)(Cl)_2(P\text{-}i\text{-}Pr_3)_2$ (**4**), **282**, 17
pentacoordinated cationic rhodium(I) and iridium(I) organocomplexes with α-diimine ligands derived from glyoxal: crystal structure of $[Rh(norbornadiene)(glyoxalbis(4\text{-}dimethylamino\text{-}anil))(PPh_3)]BF_4 \cdot 1/2CH_2Cl_2$, **282**, 230
Iron–molybdenum cofactor
X-ray absorption spectroscopic studies of the binding of ligands to FeMoco of nitrogenase from *Klebsiella pneumoniae*, **275–276**, 150
Iron–sulfur proteins
thermodynamics of reduction of *Chromatium vinosum* high-potential iron–sulfur protein and its histidine depleted H42Q mutant, **275–276**, 230
Iron complexes
synthesis and properties of the unsymmetrical binuclear complexes: $Na[(CN)_5Fe(dtdp)Ru(NH_3)_5]6H_2O$ and $K[(CN)_5Ru(dtdp)Ru(NH_3)_5]6H_2O$, dtdp = 4,4′-dithiodipyridine, **267**, 93
cyanide binding to phthalocyaninatoiron(II) in dimethyl sulfoxide in the presence of carbon monoxide: kinetic and equilibrium study, **267**, 109
NMR spectra of dinuclear manganese and iron compounds containing 1,4,7-triazacyclononane and 1,4,7-trimethyl-1,4,7-triazacyclononane, **268**, 43
'activated cyanide', **269**, 23
the cyanide ligand as an efficient bridge in mixed-valence complexes, **269**, 246
synthesis and structures of the cuboidal iron–sulfur–nitrosyl–phosphine clusters $[Fe_4S_3(NO)_4(PR_3)_3]^{0,1+}$ (R = Et, Pr^i, C_6H_{11}), **270**, 46
catalytic reduction of acetylene in the presence of molybdenum and iron clusters, including FeMo cofactor of nitrogenase, **270**, 221
dinuclear metal complexes of Cd(II), Zn(II) and Fe(II) with triple-helical structure and predetermined chirality, **271**, 36
ligand photolysis and recombination of $(dimethylsulfoxide)_2Fe(II)$-porphyrin complexes, **271**, 119

integrated photocatalysts for hydrocarbon oxidation: polyoxotungstates/iron porphyrins systems in the reductive activation of molecular oxygen, **272**, 197
sulfonated and acetamidosulfonylated tetraarylporphyrins as biomimetic oxidation catalysts under aqueous conditions, **272**, 228
NMR studies of iron–gentamicin complexes and the implications for aminoglycoside toxicity, **273**, 85
generation of mixed iron chalcogenide clusters $MFeX_n^+$ (M = V, Fe; X = O, S; n = 2, 3) in the gas phase, **273**, 111
unexpected assembly of a (μ-oxo)(μ-formato)diiron(III) complex from an aerobic methanolic solution of Fe(III) and the TPA ligand, **273**, 393
spin crossover in six-coordinate $[Fe(L)_2(NCX)_2]$ compounds with L = DPQ = 2,3-bis-(2′-pyridyl)-quinoxaline, ABPT = 4-amino-3,5-bis(pyridin-2-yl)-1,2,4-triazole and X = S, Se: synthesis, magnetic properties and single crystal studies, **274**, 1
structural and electronic studies of the coordination of 6-(thiazol-2-yl)-2,2′-bipyridine and related systems to Fe(II), Co(II) and Ni(II), **274**, 32
tris(4,4′-dimethoxy-2,2′-bipyridine)osmium(II), amperometric properties and crystal structure, **274**, 64
homogeneous catalysis by tetra-t-butylphthalocyanine iron: intermediates in the oxidative dimerization of 2,6-di-t-butylphenol, **274**, 126
binuclear oxo-bridged iron(III) complexes of chiral tetradentate ligands, **274**, 155
reduction of *Coprinus macrorhizus* peroxidase by dithionite, **274**, 215
synthesis and crystal structure of a seven-coordinate dinuclear iron(III) complex containing only bidentate ligands, **274**, 224
stereoselectivity in reactions of metal complexes. Part XXI. Kinetics and mechanism of electron transfer between *rac*-$[Co(L)H_2O]^+$ (L = *N,N′*-[(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]) and optically active iron(II) complexes, **275–276**, 1
stereoselectivity in reactions of metal complexes. Part XXII. Kinetics and mechanism of electron-transfer reactions between *rac*-$[Co(L)X]^+$ (L = [(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]; X = pyridine, imidazole, 1-methylimidazole and pyrazine) and optically active iron(II) complexes, **275–276**, 9
X-ray crystallographic and absorption spectroscopic analyses of structures of catecholato(pyridine)iron chloride complexes in relevance to functional model complexes for catechol 1,2-dioxygenases, **275–276**, 222
sterically hindered iron(II) complex of a new tripodal polyimidazole ligand: structure and reactivity toward superoxide, **275–276**, 234
metallodrug complexes that mediate DNA and lipid damage via sulfite autoxidation: copper(II) famotidine and iron(III) bis(salicylglycine), **275–276**, 314
4′-(4-fluorophenyl)-2,2′:6′,2″-terpyridine – an aryl-2,2′:6′,2″-terpyridine with an electron-releasing aryl substituent, **275–276**, 359
Fourier transform IR study of the interaction of $[(\eta^5\text{-}C_5H_5)_2FeRu(CO)_4]$ with hydroxylated alumina surfaces, **277**, 61
synthesis and characterization of a new unsymmetrical μ-oxo-bridged iron(III) complex with triphenylarsine oxide ligands, **277**, 127
crystal structure and molecular stereochemistry of the 1:2 adduct of anionic complex, dicyano-(5,10,15,20-tetramesityleneporphinato)-iron(II) with bis(triphenylphosphine)nitrogen (1+) · I_2, $CDCl_3$ solvate $\{[(CN)_2Fe^{II}(TMP)][PPN]_2\} \cdot I_2 \cdot 2CDCl_3$, **277**, 242
siderophore analogues. A new macrocyclic *bis*-(amine, amide, hydroxamate) ligand. Synthesis, solution chemistry, electrochemistry and molecular mechanics calculations for the iron complex, **278**, 51
hemoprotein models based on a covalent helix–heme–helix sandwich 4. Discrimination of paramagnetic Fe(III)–Mimochrome I Δ and Λ isomers by NMR spectroscopy, **278**, 76
spectroscopic and magnetic properties of the pyridine-2-carbaldehyde thiosemicarbazonateiron(III) complexes: $[Fe(C_7H_7N_4S)_2]X \cdot nH_2O$ (X = Cl, ClO_4, NO_3, PF_6). Crystal structure of the hexafluorophosphate compound, **278**, 150
the effect of end-group substitutions and use of a mixed solvent system on β-diketones and their iron complexes, **278**, 170
spin crossover in bis(2,6-bis(benzimidazol-2-yl)pyridine) iron(II) tetraphenylborate, **278**, 190
synthesis and characterization of a neutral, low spin iron(III) complex of a hexadentate tripodal ligand containing three imidazolate arms. Use as a dinucleating agent, **278**, 197
preparation and some properties of perfluoroalkoxy-substituted phthalocyanine complexes of iron(III), nickel(II) and zinc(II), **278**, 253
structural information on *trans*-1,2-diaminocyclohexane-*N,N′*-tetraacetateferrate(III) in the solid and aqueous phase, **279**, 37
synthesis and structural characterization of metal complexes of an unsymmetric N_3O tetradentate Schiff base ligand, **279**, 111
synthesis, crystal structures and magnetic properties of $[Fe_2(bpym)(C_5O_5)_2(H_2O)_4] \cdot 2H_2O$ and two polymorphs of $[Fe_2(bpym)(C_4O_4)_2(H_2O)_6] \cdot 2H_2O$ (bpym = 2,2′-bipyrimidine), **279**, 127
synthesis and characterisation of mononuclear and binuclear iron(II) complexes of pentadentate and bis(pentadentate) ligands derived from 1,4,7-triazacyclononane, **279**, 192
iron and manganese complexes of benzothiazolylformazans, **280**, 30
transition metal complexes with sulfur ligands. Part CXXIX. Retention and reactivity of the [Fe⎓NH⎓NH⎓Fe] chromophore in the iron sulfur diazene complex $[\mu\text{-}N_2H_2\{Fe(PPr_3)('S_4')\}_2]$ in exchange and oxidation processes. ($'S_4'^{2-}$ = 1,2-bis(2-mercaptophenylthio)ethane(2−)), **280**, 39
metal ion-coordinating properties of imidazole and derivatives in aqueous solution: interrelation between complex stability and ligand basicity, **280**, 50
chiroptical properties of heterotrimetallic 'wing-tip' butterflies; synthesis and crystal structure of $(\mu\text{-}H)Ru_2Fe_2PdC(CO)_{12}(\eta^3\text{-}\beta\text{-}C_{10}H_{15})$, **280**, 275
molecular structure and reactions of azobenzene complexes with iron–lithium compounds, **280**, 288
a novel iron sequestering agent: synthesis and iron-chelating properties of 1,1,1-tris(3-hydroxy-2-oxo-1,2-dihydro-1-pyridylpropoxymethyl)ethane, **281**, 60
ferrocenyl-substituted imine ligands and their reactivity towards $Fe_2(CO)_9$; crystal structure of Fc–C(H)=C(H)–C(H)=N–C_6H_{11} and $(\eta^4\text{-}R^1\text{–}C(H)=C(H)\text{–}C(H)=N\text{–}R^2)Fe(CO)_3$ ($R^1 = C_6H_5$, R^2 = Fc; R^1 = Fc, $R^2 = C_6H_{11}$, $R^1 = R^2$ = Fc; Fc = $(\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4)$), **282**, 111
electron donating substituent effects on redox and spin state properties of iron(II) bis-terpyridyl complexes, **282**, 131
a μ-oxo diiron complex of a new binucleating tetrapyridyl ligand, **282**, 243
folding properties of iron–sulfur proteins, **283**, 12
structure and properties of tetraphenylporphinate iron(III) complexes with an intramolecular NH···S benzenethiolate or NH···O phenolate hydrogen bond, **283**, 91
molecular structure of the methoxy-iron(III) derivative of 5,15-(*o,o′*(2-methyl-2′-hydroxy-3,3′-diamidobiphenyl)-diphenylporphyrin and CO binding properties of iron(II)–pyridine complexes of 2,2′-substituted biphenyl strapped porphyrins, **283**, 98
a synthetic iron-containing prismane complex: peculiar electron-delocalization and spin-coupling properties deduced from Mössbauer and magnetization studies, **283**, 111
catalytic oxygenation of olefin with dioxygen and tetra-t-butylphthalocyanine complexes in the presence of sodium borohydride, **283**, 275

Iron ions
angular overlap calculations of the spin Hamiltonian parameters of transition metal ions in low symmetry environments. High spin iron(II), iron(III) and manganese(III), **274**, 90

Iron-oxo porphyrin complexes

reversible formation of high-valent-iron-oxo porphyrin intermediates in heme-based catalysis: revisiting the kinetic model for horseradish peroxidase, **275–276**, 98
IR spectroelectrochemistry
infrared spectroelectrochemical and electrochemical kinetics studies of the reaction of nickel cluster radicals $[Ni_3(\mu_2\text{-dppm})_3(\mu_3\text{-L})(\mu_3\text{-I})]^{\bullet}$ (L = CNR, R = CH_3, i-C_3H_7, C_6H_{11}, $CH_2C_6H_5$, t-C_4H_9, 2,6-$Me_2C_6H_3$; L = CO) with carbon dioxide, **270**, 111
IR spectroscopy
ground and electronically excited states of $Cr(CO)_4$(bipyridine): energy factored force field analysis of CO stretching vibrations and resonance Raman study, **278**, 83
Isocyanate
mechanistic details for metathetical exchange between XCO (X = O and RN) and the tin(II) dimer, $\{Sn[N(SiMe_3)_2](\mu\text{-}OBu^t)\}_2$, **270**, 118
Isocyanide complexes
synthesis of trichlorooxo-bis(2,6-dimethylphenylisocyanide) rhenium(V) and crystal structure of μ-oxo-bis[dichlorooxo-bis(2,6-dimethylphenylisocyanide)dirhenium(V)], **269**, 347
three reactions of Ru_2^{5+} compounds of the paddlewheel type that lead to cleavage of the Ru–Ru bond, **275–276**, 541
ammonia ruthenium complexes for the access to new hydrido, carbonyl and isocyanide ruthenium-alkynyl derivatives, **280**, 118
Isodiazene complexes
the synthesis and characterization of a technetium(III) isodiazene complex. The X-ray crystal structure of $[TcCl_3(N{=}NPh_2)(PPh_3)_2]$, **271**, 191
cis and *trans* Isomers
coordination chemistry of polyhydroxy acids: role of the hydroxy groups, **275–276**, 130
Isothiocyanate complexes
crystal and molecular structures of some tri-aryltin isothiocyanates (Ar_3SnNCS; Ar = C_6H_5, *p*-$CH_3C_6H_4$, *o*-$CH_3C_6H_4$, *o*-$CH_3OC_6H_4$) and of the adduct hexamethylphosphoramide(*N*-thiocyanato)-tris(*o*-tolyl)tin, **282**, 30
Itaconic acid complexes
synthesis and structural properties of (diamine)platinum(II) complexes of itaconic acid, **279**, 116

Jahn–Teller coupling
the effects of d–s mixing in low symmetry transition metal complexes, **268**, 55
Jahn–Teller effects
chemical applications of topology and group theory. 30. Distortions from idealized symmetrical coordination polyhedra by lifting molecular orbital degeneracies, **270**, 68

Kanamycin complexes
kanamycin revisited: a combined potentiometric and spectroscopic study of copper(II) binding to kanamycin B, **275–276**, 546
Kinetics
fast water exchange on a Pt(II) center: variable temperature and pressure ^{17}O NMR study at 14.1 tesla of the ionic arylplatinum species $[Pt\{C_6H_3(CH_2NMe_2)_2\text{-}2,6\}(OH_2)]^+(OSO_2CF_3)$, **269**, 322
the dynamics of formation of the O_2–Co^{II} bond in the cobalt(II) cyclidene complexes, **270**, 151
the dissociation kinetics of the ethylenediaminetetrakis(methylenephosphonato) complex of europium(III), **278**, 185
ricochet inter-ring haptotropic rearrangement of σ-methyl-(η^5-indenyl)chromium tricarbonyls. Experimental kinetic and theoretical DFT study, **280**, 328
substitution reaction of aqua(ethylenediaminetetraacetato) ruthenium(III) with 2-mercaptobenzoic acid, **282**, 38
Kinetics and mechanism
the kinetics and mechanism of ligand exchange in photogenerated (η^1-bromobenzene)$Cr(CO)_5$/alkene systems, **267**, 49
4- and 3-hydroxypyridine complexes of pentacyanoferrate(II, III), **267**, 81
cyanide binding to phthalocyaninatoiron(II) in dimethyl sulfoxide in the presence of carbon monoxide: kinetic and equilibrium study, **267**, 109
the kinetics and mechanism of the oxidation of 1,4-thioxane by potassium ferrate, **267**, 159
properties and kinetics of dihydroxy- and diaminoanthraquinone ruthenium bipyridyl dimers, **268**, 31
substituent effects of *cis*-dioxobis(dithiocarbamato)molybdenum(VI) on redox properties: redox potentials for one-electron reduction and second-order rate constants for oxygen atom transfer, **269**, 260
kinetics and mechanism of the oxidation of ammineruthenium(II) complexes by bromine, **270**, 189
rate constants and activation parameters for organo–cobalt porphyrin bond homolysis from NMR relaxation times, **270**, 197
kinetics and mechanism of the axial substitution reactions of (ligand)bis(4,5-dichloro-1,2-benzosemiquinonediiminato)-cobalt(III) complexes, **270**, 326
model reactions of a carbonylation catalyst: phosphite induced migratory CO insertion in $[MeIr(CO)_2I_3]^-$, **270**, 382
monovalent copper as a potential catalyst for formation of acetaldehyde via the migration of methyl radicals to the coordinated carbonyl in the complex $(CO)Cu^{II}$–CH_3^+, **270**, 440
first- and second-order mechanisms for oxidative addition for bound methyl disulfide in the complex $W(CO)_3$(phen)(MeSSMe), **270**, 488
bonding modes of arenes in photogenerated (arene)$Cr(CO)_5$ complexes, **271**, 105
stopped-flow spectrophotometric study on the reactions between copper(II) ions and some dithiophosphates, **272**, 291
kinetics and mechanism of ester hydrolysis by metal complexes of 2,6-diacetylpyridine dioxime, **273**, 167
stereoselectivity in reactions of metal complexes. Part XXI. Kinetics and mechanism of electron transfer between *rac*-$[Co(L)H_2O]^+$ (L = *N,N'*-[(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]) and optically active iron(II) complexes, **275–276**, 1
stereoselectivity in reactions of metal complexes. Part XXII. Kinetics and mechanism of electron-transfer reactions between *rac*-$[Co(L)X]^+$ (L = [(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]; X = pyridine, imidazole, 1-methylimidazole and pyrazine) and optically active iron(II) complexes, **275–276**, 9
electrophilic substitution of metal-coordinated pyrazole: nitration, sulfonation and bromination of $[Co(NH_3)_5(pyzH)]^{3+}$ (pyzH = pyrazole), **277**, 89
kinetic studies using Co(III) and Cr(III) complexes of 1,4,7,11-tetraazaundecane (2,2,3-tet) and the crystal structure of *trans*-$[CrCl_2(2,2,3\text{-}tetH)(OH_2)]Cl\cdot ClO_4$, **278**, 24
the redox kinetics of platinum(II)/(IV) complexes, **278**, 32
synthesis, characterization and kinetics of the metallation of the intermediate compound $[Rh_2(\mu\text{-}O_2CCH_3)_3(O_2CCH_3)\{\eta^2\text{-}\underline{P}(2\text{-}(CH_3\underline{O})C_6H_4)Ph_2\}]\cdot H_2O$ to produce compound $[Rh_2(\mu\text{-}O_2CCH_3)_3\text{-}\{\mu\text{-}(\underline{C}C_5H_4)\underline{P}(2\text{-}(CH_3O)C_6H_4)Ph\}]\cdot(H_2O)$ from Rh(II) tetraacetate, **278**, 61
bonding of halogenated arenes in photogenerated (arene)$M(CO)_5$ complexes (M = Cr, Mo, W), **278**, 202
enantioselective cleavage of activated amino acid esters promoted by chiral palladacycles, **280**, 57
intramolecular protonation and other mechanisms for substitution reactions of hydrido(thiolato)– and di(thiolato)–ruthenium(II) phosphine complexes, **280**, 75
the kinetics and mechanism of oxidation of 3-mercaptopropionic acid, 2-mercaptoethanesulfonic acid and 2-mercaptobenzoic acid by potassium ferrate, **281**, 43

kinetics and mechanism for reduction of *trans*-dichloro-tetracyano-platinate(IV) by tetraammineplatinum(II) and bis(ethylene-diamine)platinum(II), **282**, 55
kinetics of the photolysis of bis(triethylphosphine)dichloro-platinum(II) in chloroform, **282**, 217
hydrolysis of phosphate esters catalyzed by copper(II)-triamine complexes. The effect of triamine ligands on the reactivity of the copper(II) catalysts, **283**, 51
spectroscopic characterization and kinetic studies of a novel plastocyanin from the green alga *Ulva pertusa*, **283**, 184

Lanthanide complexes
complexes of lead(II) and lanthanide(III) ions with two novel 26-membered-imine and -amine macrocycles derived from 2,6-bis(2-formylphenoxymethyl)pyridine, **267**, 55
role of the intramolecular hydrogen bond and ligand rigidity in the complexation of trifluoroacetylcycloalkanones with lanthanides: novel strategy for the design of organic ligands of high selectivity, **267**, 201
phosphide complexes of divalent europium: synthesis and structural characterization of $Eu[PPh_2]_2(N\text{-MeIm})_4$, **267**, 309
homodinuclear lanthanide complexes: Ln_2L_3 (H_2L = tetradentate Schiff bases). Magnetic properties (solid state) and spectroscopic studies (solution), **268**, 125
lanthanide complexes with 1-hydroxyethane-1,1-diphosphonic acid: solvent organization and coordination geometry in crystalline and amorphous solids, **269**, 211
lanthanide metal complexes of *N,N'*-ethylenebis(pyridin-2-one); from chelates to metallacyclic frameworks, **272**, 131
functionalized acyclic Schiff bases and related complexes with d- and f-metal ions, **272**, 235
different picrate coordination modes along the lanthanide series: synthesis and molecular structure of two isomorphous compounds $[Ln_2(pic)_4(TDTD)_3(H_2O)_2](pic)_2(H_2O)_2$ (Ln = Tm, Lu; pic = picrate; TDTD = *trans*-1,4-dithiane-*S,S'*-dioxide), **274**, 122
synthesis, structure and properties of lanthanide dithionates and their triphenylphosphine oxide complexes, **274**, 177
synthesis, UV–visible and electrochemical studies of lipophilic and hydrophilic lanthanide(III) bis(porphyrinates), **275–276**, 182
potentiometric and NMR spectroscopic study of protonations and amide hydrogen exchange rates of DTPA-bis(butylamide), DTPA-bis(glucamide), and their lanthanide(III) complexes, **277**, 193
synthesis and X-ray crystal determination of four novel lanthanide(III) oxydiacetate complexes, **281**, 18
lanthanide(III) complexes of two oxaazadiamine macrocyclic ligands derived from 2,6-diformylpyridine: the crystal structures of a reduced macrocyclic ligand and the corresponding diprotonated macrocycle, **282**, 42
Lanthanum complexes
multinuclear NMR study of the interactions between the La(III) complex of DTPA-bis(glucamide) and Zn(II) or borate, **268**, 249
synthesis and crystal structure of a new polynuclear lanthanum picrate complex with *trans*-1,4-dithiane-1,4-dioxide, **274**, 236
Lead complexes
complexes of lead(II) and lanthanide(III) ions with two novel 26-membered-imine and -amine macrocycles derived from 2,6-bis(2-formylphenoxymethyl)pyridine, **267**, 55
complexes of thallium(I) and lead(II) with the potentially tetradentate ligand bis[3-(2-pyridyl)-pyrazolyl]dihydroborate, **267**, 323
Lewis base complexes
Lewis base adducts of halogenorhenium(VII) oxides: ^{17}O NMR spectroscopy, structural aspects and catalysis, **279**, 44
Ligand exchanges
imidazole release from the antitumor-active ruthenium complex imidazolium *trans*-tetrachlorobis(imidazole)ruthenate(III) by biologically occurring nucleophiles, **267**, 137
salicylaldehyde-2-mercaptoanilato(2−)-2-hydroxyphenyl-benzthiazolato-nitrosyl-molybdenum(II) — a chelate which simultaneously contains an azomethine ligand and its cyclic oxidation product, **269**, 350
Ligand photolysis
ligand photolysis and recombination of (dimethylsulfoxide)$_2$Fe(II)-porphyrin complexes, **271**, 119
Ligand rigidity
role of the intramolecular hydrogen bond and ligand rigidity in the complexation of trifluoroacetylcycloalkanones with lanthanides: novel strategy for the design of organic ligands of high selectivity, **267**, 201
Lightly bonded ligand complexes
dissociative kinetics of 'lightly bonded ligands' *L* from the clusters $Rh_6(CO)_{15}L$ (*L* = DMSO, NCMe, cyclooctene, THF and EtOH), **268**, 189
Linear compounds
organic–inorganic composite materials: structural archetypes of linear polymeric chlorocadmates(II), **272**, 252
Linkage isomerism
linkage isomers of alizarin-bis(bipyridine)ruthenium(II), **281**, 126
Linkage isomers
synthesis and rearrangements of the pentaamminecobalt(III) linkage isomers of some amidic acids, **268**, 85
Lipophilic complexes
synthesis, UV–visible and electrochemical studies of lipophilic and hydrophilic lanthanide(III) bis(porphyrinates), **275–276**, 182
Lithium complexes
syntheses and structure of dimeric yttrium tris[bis(trimethylsilyl)-phosphanide]; X-ray structures of $[(thf)LiCl\cdot(thf)_2LiP^tBu_2]_2$ and $Y[CH(SiMe_3)_2]_3$, **269**, 91
molecular structure and reactions of azobenzene complexes with iron–lithium compounds, **280**, 288
Low-dimensional magnetics
chain-like crystal structure and magnetic properties of [catena-bis(1,2-diaminoethane)copper(II)-μ-dicyano-argentate]-dicyanoargentate, **281**, 134
Low-spin complexes
crystal structure and molecular stereochemistry of the 1:2 adduct of anionic complex, dicyano-(5,10,15,20-tetramesityleneporphinato)-iron(II) with bis(triphenylphosphine)nitrogen (1+)·I_2, $CDCl_3$ solvate $\{[(CN)_2Fe^{II}(TMP)][PPN]_2\}\cdot I_2\cdot 2CDCl_3$, **277**, 242
synthesis and characterization of a neutral, low spin iron(III) complex of a hexadentate tripodal ligand containing three imidazolate arms. Use as a dinucleating agent, **278**, 197
Low symmetry ligand fields
the effects of d–s mixing in low symmetry transition metal complexes, **268**, 55
Low-valence complexes
polymeric low-valence platinum–phenanthroline complexes as precursors of platinum colloids, **280**, 339
Lumazine complexes
synthesis and structural studies on three-dimensionally hydrogen-bonded metal perchlorate complexes with 1,3-dimethyllumazine and 1,3,6,7-tetra-methyllumazine. Crystal structure of diaqua-bis-(1,3,6,7-tetramethyl-pteridine-2,4(1*H*,3*H*)-dione-O^4,N^5)-copper(II) perchlorate dihydrate, **268**, 77
Lumazine derivatives complexes
synthesis and structural studies on metal nitrate complexes with 1,3-dimethyllumazine and 1,3,6,7-tetramethyllumazine: crystal structure of two new cobalt(II) and copper(II) three-dimensionally hydrogen-bonded complexes and a cadmium(II) complex with unusual geometry, **277**, 103
Luminescence
evaluation of solution structures of highly luminescent europium(III) chelates by using laser induced excitation of the $^7F_0 \rightarrow {}^5D_0$ transition, **267**, 63

remarkable substituent effects on the photophysics of Pt(4′-X-trpy)Cl$^+$ systems (trpy = 2,2′; 6′,2″-terpyridine), **273**, 346
photoinduced processes in 4′-(9-anthryl)-2,2′:6′,2″-terpyridine, its protonated forms and Zn(II), Ru(II) and Os(II) complexes, **277**, 225

Lutetium complexes
C_{60}-fullerenides of ytterbium and lutetium, **280**, 138

Macrocycle complexes
electronic spectra and structures of Cu^{2+} and Ni^{2+} complexes with dioxotetraazacycloalkanediacetates and their diester derivatives, **269**, 224
the dynamics of formation of the O_2–Co^{II} bond in the cobalt(II) cyclidene complexes, **270**, 151
synthesis and coordinating properties of the functionalized macrocyclic ligand 1,4-bis(1-methylimidazol-2-ylmethyl)-7-carboxymethyl-1,4,7-triazacyclononane. Crystal structures of cobalt(III), nickel(II) and zinc(II) complexes, **273**, 151
a large cavity reinforced cryptand for the binding of metal cations and anions, **273**, 326
synthesis and characterization of nickel(II) complexes of di-*N*-alkylated 14-membered tetraaza macrocycles containing cyclohexane rings via regioselective alkylation reaction, **279**, 238
new cyclam-type copper(II) complexes with amide molecular padlock: synthesis, properties and crystal structure, **282**, 142
synthetic models for biological trinuclear copper clusters. Trinuclear and binuclear complexes derived from an octadentate tetraamine-tetrabenzimidazole ligand, **282**, 180
binuclear nickel(II) and copper(II) complexes of amide-containing two-armed and macrocyclic ligands, **282**, 222

Macrocyclic ligand complexes
cooperative bimetallic reactivity, **270**, 13
structure of an indium(III) complex with a 15-membered macrocycle, dioxopentaazacyclopentadecanetriacetate, **271**, 207
macrocyclic ligand design: structure–function relationships involving the interaction of cobalt(II), nickel(II) and copper(II) with mixed donor macrocyclic ligands, **273**, 372
siderophore analogues. A new macrocyclic *bis*-(amine, amide, hydroxamate) ligand. Synthesis, solution chemistry, electrochemistry and molecular mechanics calculations for the iron complex, **278**, 51
synthesis, structure and characterization of zinc(II), copper(II), zinc(II)barium(II) and copper(II)barium(II) complexes of macrocyclic heteronucleating ligands based on isothiosemicarbazide, **282**, 61

Macrocyclic tetraamine complexes
a new macrocyclic tetraamine, 2,4-dinitrophenylcyclen (= 1-(2,4-dinitrophenyl)-1,4,7,10-tetraazacyclododecane): synthesis, cation reporter properties, and recognition of 1-methylthymine by its zinc(II) complex, **270**, 424

Magnesium activation
magnesium activation of nuclease enzymes — the importance of water, **275–276**, 24

Magnesium complexes
structure and magnetic properties of a novel chloro-bridged polymeric cadmium(II) complex with pyridyl-substituted nitronyl nitroxide, **279**, 233

Magnetic circular dichroism
spectroscopic study of the compound ES and the oxoferryl compound II states of cytochrome *c* peroxidase: comparison with the compound II of horseradish peroxidase, **275–276**, 250

Magnetic circular dichroism spectroscopy
spectroscopic distinction between two Co(II) ions substituted for types 1 and 2 Cu in nitrite reductase, **275–276**, 289
electronic absorption and MCD spectra for the Pt(PPh$_3$)(AuPPh$_3$)$_6^{2+}$ and Pt(H)(PPh$_3$)(AuPPh$_3$)$_7^{2+}$ cluster complexes, **278**, 113

Magnetic interactions
transition metal complexes with nitronyl nitroxides. Part 1. Crystal structure and properties of square-planar *trans* radical–copper(II)–radical complexes, **279**, 69
transition metal complexes with nitronyl nitroxides. Part 2. Crystal structure and magnetism of manganese(II) compounds including radicals and their reduced derivatives, **279**, 76
dinuclear $Cu^{II}M^{II}$ complexes of a phenol-based macrocycle with N(amine)$_2$O$_2$ and N(imine)$_2$O$_2$ metal-binding sites: the effect of chloride ligation upon the site selectivity of metal ions, **283**, 72

Magnetic properties
crystal structure, spectroscopic, magnetochemical, thermoanalytical and electrochemical properties of binuclear copper(II) complexes of Suprofen, **268**, 239
structure and magnetic properties of a linear oximato-bridged tetranuclear copper(II) complex, **268**, 263
synthesis and characterization of electric and magnetic properties of the new intermetallic compounds M$_3$(GaTe$_3$)$_2$ (M = Cr, Fe, Co), **268**, 271
{*catena*-($\mu_{N,S}$-NCS)[Ni(Medien)(NCS)]}: a new 1D nickel(II) complex with bridging thiocyanato ligands in *cis* position. Crystal structure and magnetic properties, **269**, 313
stepwise substituting magnetic centres in a giant cluster host system, **271**, 9
synthesis, structure and magnetic properties of one-dimensional azide-bridged manganese(III) uniform chain complex Mn(salpn)N$_3$, **271**, 99
synthesis, crystal structure and magnetic properties of bis[μ-bis(3-(pyridin-2-yl)-1,2,4-triazole-*N′*,*N*1,*N*2)] bis[triaqua nickel(II)] tetranitrate, **273**, 54
synthesis, structure characterization and magnetic properties of tellurostannates [M(en)$_3$]$_2$(Sn$_2$Te$_6$) (M = Mn, Zn), **273**, 310
spin crossover in six-coordinate [Fe(L)$_2$(NCX)$_2$] compounds with L = DPQ = 2,3-bis-(2′-pyridyl)-quinoxaline, ABPT = 4-amino-3,5-bis(pyridin-2-yl)-1,2,4-triazole and X = S, Se: synthesis, magnetic properties and single crystal studies, **274**, 1
synthesis, crystal structures and magnetic properties of [Fe$_2$(bpym)(C$_5$O$_5$)$_2$(H$_2$O)$_4$]·2H$_2$O and two polymorphs of [Fe$_{\text{character.2}}$(bpym)(C$_4$O$_4$)$_2$(H$_2$O)$_6$]·2H$_2$O (bpym = 2,2′-bipyrimidine), **279**, 127
synthesis, crystal structure and magnetic properties of the first single azido-bridged copper(II) chain [Cu(bpym)(N$_3$)$_2$]$_n$ (bpym = 2,2′-bipyrimidine), **282**, 252
synthesis, crystal structure and magnetic properties of the one-dimensional chain compound *catena*-poly[aquabis(4,4′-bis-1,2,4-triazole-*N*1)(thiocyanato-*N*)-manganese-μ-[4,4′-bis-1,2,4-triazole-*N*1,*N*1′]thiocyanate], **282**, 257

Magnetic resonance imaging
a tripod gadolinium(III) poly(amino carboxylate) relevant to magnetic resonance imaging: structural and dynamical ^{17}O NMR and ^{1}H NMRD studies, **275–276**, 15
new non-ionic water-soluble porphyrins: evaluation of manganese(III) polyhydroxylamide porphyrins as MRI contrast agents, **275–276**, 106

Magnetic resonance imaging contrast agents
multinuclear NMR study of the interactions between the La(III) complex of DTPA-bis(glucamide) and Zn(II) or borate, **268**, 249
potentiometric and NMR spectroscopic study of protonations and amide hydrogen exchange rates of DTPA-bis(butylamide), DTPA-bis(glucamide), and their lanthanide(III) complexes, **277**, 193

Magnetic susceptibility
polymeric and dimeric magnetic properties of square planar Cu(II) species controlled by hydrogen bond networks: [Cu^{II}{OCO-2,6-(CH$_3$CONH)$_2$C$_6$H$_3$}$_2$(H$_2$O)$_2$]·nH$_2$O (n = 1, 4), **275–276**, 43

Magnetism
synthesis, X-ray structure, spectroscopic and magnetic properties of [bis(4-amino-3,5-bis(pyridin-2-yl)-1,2,4-triazole-*N′*,*N*1)-(aqua)copper(II)]bis(hydrogensulfate), **268**, 37

crystal structure, EPR and magnetic susceptibility studies of tetrakis[μ-(β-alanine)-*O,O′*]dichlorodicopper(II) dichloromonohydrate, **275–276**, 28

spectroscopic and magnetic properties and structure of a five-coordinate, O_5-binding cobalt(II) Schiff base complex and of the copper(II) analogue, **278**, 127

spin crossover in bis(2,6-bis(benzimidazol-2-yl)pyridine) iron(II) tetraphenylborate, **278**, 190

electron donating substituent effects on redox and spin state properties of iron(II) bis-terpyridyl complexes, **282**, 131

synthesis, structure and characterization of dinuclear nickel(II) complexes containing the derivatives of *N,N,N′,N′*-tetrakis[(2-benzimidazolyl)methyl]-2-hydroxy-1,3-diaminopropane and an acetate or a propionate ion, **283**, 118

Main group metal complexes

nucleophilic attack within Ge, Sn and Pb complexes containing $Me_2N(CH_2)_3$– as a potential intramolecular donor ligand, **275–276**, 203

Maleimide complexes

synthesis of two maleimide derivatives of *cis*-configurated platinum(II) complexes for the preparation of chemoimmunoconjugates, **267**, 133

Maleonitriledithiolate complexes

the synthesis and characterization of Ni, Pd and Pt maleonitriledithiolate complexes: X-ray crystal structures of the isomorphous Ni, Pd and Pt $(Ph_2PCH_2CH_2PPh_2)M$(maleonitriledithiolate) congeners, **282**, 155

Maltolate complexes

the formation of ternary complexes between VO(maltolate)$_2$ and small bioligands, **283**, 202

Manganese carbonyls

nucleophilic attack at the five-coordinate anion $[Mn(CO)_3$(3,5-di-tert-butyl-catecholate)$]^-$ controlled by electronic and steric effects, **268**, 1

Manganese catalase

structural and spectroscopic models of the manganese catalase active site. Isolation and structures of the asymmetric $[(H_2O)Mn^{III}(\mu\text{-}O)(\mu\text{-}O_2CR)_2Mn^{III}(L)]$ ($L = Cr_2O_7^{2-}$, CH_3OH) cores: analogs of a substrate-bound catalase active site intermediate, **281**, 25

Manganese complexes

NMR spectra of dinuclear manganese and iron compounds containing 1,4,7-triazacyclononane and 1,4,7-trimethyl-1,4,7-triazacyclononane, **268**, 43

EXAFS studies on Mn(III) tetraphenylporphyrin halides in Nafion, **268**, 231

the cyanide ligand as an efficient bridge in mixed-valence complexes, **269**, 246

synthesis, structure and magnetic properties of one-dimensional azide-bridged manganese(III) uniform chain complex Mn(salpn)N_3, **271**, 99

a new manganese-β-heptanitro-porphyrin with extreme redox potentials: spectral, electrochemical and catalytic properties, **272**, 4

metal complexes of Imazapyr, a herbicide provided with efficient metal-chelating ability: crystal structure of the cobalt(III) and manganese(II) complexes, **272**, 68

comparison of the neutral benzyldiphenylphosphine complexes $C_6H_6Cr(CO)_2PPh_2Bz$ and $C_5H_5Mn(CO)_2PPh_2Bz$ with the isoelectronic manganiobenzyldiphenylphosphonium salts $[C_5R_5Mn(CO)(NO)PPh_2Bz]BF_4$, **272**, 101

spectroscopic characterization of fluorinated/hydrogenated mixed vesicles containing fluorinated Mn(III)-porphyrin, **272**, 274

a ^{31}P and ^1H NMR relaxometric study of the interaction between adenosine triphosphate (ATP) and paramagnetic ions (Gd^{3+} and Mn^{2+}), **273**, 92

synthesis, structure characterization and magnetic properties of tellurostannates $[M(en)_3]_2(Sn_2Te_6)$ (M = Mn, Zn), **273**, 310

new non-ionic water-soluble porphyrins: evaluation of manganese(III) polyhydroxylamide porphyrins as MRI contrast agents, **275–276**, 106

characterisation of covalent copper and manganese organometallic complexes with Schiff bases by ionspray mass spectrometry, **275–276**, 462

crystal structure and properties of a new five-coordinate manganese superoxide dismutase mimic, **277**, 21

synthesis and catalytic activity of manganese(II) complexes of pentanedioic acid; X-ray crystal structure of $[Mn(phen)_2(H_2O)_2]$-$[Mn(O_2C(CH_2)_3CO_2)(phen)_2H_2O](O_2C(CH_2)_3CO_2)\cdot 12H_2O$ (phen = 1,10-phenanthroline), **277**, 257

tuning up the T_C in Mn(II)Cu(II) bimetallic planes and design of molecular-based magnets, **278**, 159

syntheses and X-ray crystal structures of *cis*-$[Mn(bipy)_2Cl_2]\cdot 2H_2O\cdot EtOH$ and *cis*-$[Mn(phen)_2Cl_2]$ (bipy = 2,2′-bipyridine; phen = 1,10-phenanthroline); catalysts for the disproportionation of hydrogen peroxide, **279**, 24

transition metal complexes with nitronyl nitroxides. Part 2. Crystal structure and magnetism of manganese(II) compounds including radicals and their reduced derivatives, **279**, 76

synthesis and characterization of the new complex tris[(2-oxazolinyl)phenolato] manganese(III) and its use as a catalyst in oxidation reactions, **279**, 217

iron and manganese complexes of benzothiazolylformazans, **280**, 30

chelated alkyl halide manganese complexes $(\eta^5{:}\eta^1\text{-}C_5H_4CH_2CH_2X)$-$Mn(CO)_2$ from photolysis of $(\eta^5\text{-}C_5H_4CH_2CH_2X)Mn(CO)_3$, **280**, 316

structural and spectroscopic models of the manganese catalase active site. Isolation and structures of the asymmetric $[(H_2O)Mn^{III}(\mu\text{-}O)(\mu\text{-}O_2CR)_2Mn^{III}(L)]$ ($L = Cr_2O_7^{2-}$, CH_3OH) cores: analogs of a substrate-bound catalase active site intermediate, **281**, 25

complexes of the bis(trimethylsilyl)-benzamidinato ligand '*siam*': synthesis and X-ray structures of (*siam*)$_2$M, (*siam*)(*siam*H)MX (M = Ni, Pd), (*siam*)$_2$MnI and (*siam*)ReO_3, and their reactivity towards CO_2, **281**, 181

synthesis of manganese tricarbonyl cationic complexes of ferrocenyl substituted arenes via a manganese tricarbonyl cation transfer reaction, **281**, 229

relative reactivities of hydroxy compounds with short-lived manganese(V), **281**, 235

synthesis, crystal structure and magnetic properties of the one-dimensional chain compound *catena*-poly[aquabis(4,4′-bis-1,2,4-triazole-*N*1)(thiocyanato-*N*)-manganese-μ-[4,4′-bis-1,2,4-triazole-*N*1,*N*1′]thiocyanate], **282**, 257

catalytic oxygenation of olefin with dioxygen and tetra-t-butylphthalocyanine complexes in the presence of sodium borohydride, **283**, 275

Manganese ions

angular overlap calculations of the spin Hamiltonian parameters of transition metal ions in low symmetry environments. High spin iron(II), iron(III) and manganese(III), **274**, 90

Mass spectrometry

generation of mixed iron chalcogenide clusters $MFeX_n^+$ (M = V, Fe; X = O, S; $n = 2, 3$) in the gas phase, **273**, 111

liquid secondary ion mass spectrometric study of diruthenium(II,III) complexes, **277**, 76

Melanostatin complexes

ESR and NMR investigation of structure and dynamics of the Cu(II) complex of melanostatin in DMSO solution, **275–276**, 274

Membrane-bound bc_1 complexes

spectroscopic investigations on the water-soluble fragment of the Rieske [2Fe–2S] protein from *Paracoccus denitrificans*, **275–276**, 493

Membranous enzymes

desulfoviridin, the dissimilatory sulfite reductase from *Desulfovibrio desulfuricans* (Essex): new structural and functional aspects of the membranous enzyme, **275–276**, 52

α-Mercaptoacid complexes
synthesis and characterization of molybdenum(VI)-dioxo complexes containing both coordinated thiolate and carboxylate groups. Reactions with their own free ligands, **268**, 145

Mercaptopurine complexes
molybdenum complexes of the anticancer drug 6-mercaptopurine, **279**, 136

Mercury complexes
synthesis and X-ray structure of the thioethereal mercury(II) complex $[N(CH_2CH_2SCHMe_2)_3HgCl]_2(Hg_2Cl_6)$, **269**, 274
interaction of Hg(II) with tetrapeptides containing cysteinyl and histidinyl residues, **272**, 7
preparation and X-ray crystal structures of $Li_4[Hg_2(edta)_2]\cdot 8H_2O$ and $Ba_2[Hg_2(teta)_2]\cdot 9H_2O$ (teta = *N,N,N′,N′*-diaminobutanetetraacetate), **274**, 219
complexes of heterocyclic thiones and group twelve metals. Part 1. Preparation and characterisation of 1:1 complexes of mercury(II) halides with 1-methylimidazoline-2(3H)-thione: the crystal structure of [(μ_2-dibromo) bis(*trans*{(bromo)(1-methyl-imidazoline-2(3H)-thione)}mercury(II))] at 160 K, **277**, 14
synthesis and structure of (1,3-dimethyluracil-5-yl)mercury(II) complexes with aromatic nitrogen donor ligands, **282**, 237

Mercury(II) complexes
a bis(9-methyladeninium) complex of Hg(II) with a highly irregular coordination geometry: $[Hg(9\text{-MeAH-}N7)_2(H_2O)(NO_3)_3]ClO_4$, **267**, 87

Mesityl compounds
alkoxymesitylchlorogermanes and aryloxymesitylchlorogermanes, **277**, 98

Meso-helicate complexes
a linear open-chain piperazine–pyridine ligand and its *meso*-helical Co complex, **277**, 55

Metal binding
metal binding properties of three Cys_2X_2 (X = His, Asp) metallothionein-related peptides, **278**, 10

Metal complexes
reinforced piperazine rings containing polyamines: metal complex equilibria and structural studies, **268**, 63
synthesis and structural studies on three-dimensionally hydrogen-bonded metal perchlorate complexes with 1,3-dimethyllumazine and 1,3,6,7-tetra-methyllumazine. Crystal structure of diaqua-bis-(1,3,6,7-tetramethyl-pteridine-2,4(1*H*,3*H*)-dione-O^4,N^5)-copper(II) perchlorate dihydrate, **268**, 77
kinetics and mechanism of ester hydrolysis by metal complexes of 2,6-diacetylpyridine dioxime, **273**, 167
interaction of metal compounds with 'double-hydrophilic' block copolymers in aqueous medium and metal colloid formation, **280**, 348

Metal hydroxide compounds
X-ray absorption fine structure study of the atomic and electronic structure of molybdenum disulfide intercalation compounds with transition metals, **280**, 211

Metal ions
an electrospray mass spectrometric and voltammetric study of horse heart cytochrome *c* in the presence of metal ions, **267**, 281
interaction of manganese(II), cobalt(II) and nickel(II) with DNA oligomers studied by ^1H NMR spectroscopy, **273**, 41

Metallacycle complexes
mononuclear and binuclear *o*-xylylene derivatives of Pd(II), **269**, 191
lanthanide metal complexes of *N,N′*-ethylenebis(pyridin-2-one); from chelates to metallacyclic frameworks, **272**, 131

Metallated complexes
synthesis, characterization and kinetics of the metallation of the intermediate compound $[Rh_2(\mu\text{-}O_2CCH_3)_3(O_2CCH_3)\{\eta^2\text{-}\underline{P}(2\text{-}(CH_3\underline{O})C_6H_4)Ph_2\}]\cdot H_2O$ to produce compound $[Rh_2(\mu\text{-}O_2CCH_3)_3\text{-}\{\mu\text{-}(\underline{C}\underline{C}_5H_4)\underline{P}(2\text{-}(CH_3O)C_6H_4)Ph\}]\cdot(H_2O)$ from Rh(II) tetraacetate, **278**, 61

Metallophthalocyanines
azide promotion of alkane oxidations catalyzed by metal complexes in solution, **270**, 162

Metalloporphyrins
azide promotion of alkane oxidations catalyzed by metal complexes in solution, **270**, 162
conductive polymeric films of *meso*-tetrakis[2,2]-*para*-cyclophanylmetalloporphyrins — spectroelectrochemistry and catalytic properties, **271**, 167

Metalloprotein complexes
probing the metal site in *Rhus vernicifera* stellacyanin by Ni(II) substitution and paramagnetic NMR spectroscopy, **273**, 367
miniaturized hemoproteins: design, synthesis and characterization of mimochrome II, **275–276**, 301
selective versus non-selective T_1 experiments to determine metal–nucleus distances in paramagnetic metalloproteins, **275–276**, 373

Metal–metal bonds
trimethylphosphine halide complexes of titanium, zirconium and hafnium in oxidation states IV, III and II. Crystal structure of $MI_4(PMe_3)_n$ (M = Zr, n = 3; M = Hf, n = 2.5), $M_2X_6(PMe_3)_4$ (M = Hf, Ti, X = I; M = Zr, X = Br) and $(\mu\text{-}\eta^6,\eta^6\text{-}C_6H_6)\text{-}Hf_2I_4(PMe_3)_4\cdot C_6H_6$, **271**, 180
synthesis and gas-occlusion properties of dinuclear molybdenum(II) dicarboxylates (fumarate, terephthalate, *trans-trans*-muconate, pyridine-2,5-dicarboxylate, and *trans*-1,4-cyclohexanedicarboxylate), **283**, 268

Metal–metal bond cleavage
three reactions of Ru_2^{5+} compounds of the paddlewheel type that lead to cleavage of the Ru–Ru bond, **275–276**, 541

Metal–metal bond complexes
synthesis of tetracarboxylatediruthenium(II) compounds: reduction and disproportionation reactions of diruthenium(II,III) units. Crystal structure of $Ru_2(\mu\text{-}O_2CCOPh)_4(thf)_2$, **279**, 30

Metal–metal multiply-bonded complexes
metal–metal multiply-bonded complexes of technetium. 7. Oxidative decomposition of tetrachlorotetrakis(dimethylphenylphosphino)-ditechnetium(II) by an aminium hexachloroantimonate(V), **271**, 187

Metal-porphines
a catalytic activity of glass beads, silica gels and anion-exchange resins modified with metal-porphines in oxidative reactions of ascorbic acid, **283**, 44

Metal surfaces
carbonyl bonding on ZnO(0001), CuCl(111) and Cu_2O(111) $3d^{10}$ metal ion surfaces: photoelectron spectroscopy of CO/Cu_2O(111) and electronic structure calculations, **275–276**, 327

Methemoglobin
interspin distances determined by time domain EPR of spin-labeled high-spin methemoglobin, **273**, 354

Methionine sulfur
detection of two novel lowest lying singlet–triplet transitions of the C–S–C chromophore in L-methionine, **272**, 150

Methyl cobalt complexes
the Co–CH_3 bond in Schiff base B_{12} models: influence of the *trans* and equatorial ligands as assessed by Fourier transform Raman spectroscopy, **275–276**, 90

Methyl complexes
thermodynamics of the formation of *cis*-bis(dialkylsulfide)dimethylplatinum(II) complexes from bis(μ-dialkylsulfide)bisdimethylplatinum(II) and free dialkylsulfide, **268**, 311
selective C–H bond activation of arenes catalyzed by methylrhenium trioxide, **270**, 55
model reactions of a carbonylation catalyst: phosphite induced migratory CO insertion in $[MeIr(CO)_2I_3]^-$, **270**, 382

monovalent copper as a potential catalyst for formation of acetaldehyde via the migration of methyl radicals to the coordinated carbonyl in the complex $(CO)Cu^{II}–CH_3^+$, **270**, 440

intramolecular and intermolecular C–H activation at a cationic Pt^{II} center, **270**, 467

1-Methylcytosine complexes

tris- and tetrakis(1-methylcytosine) complexes of Pt(II): syntheses and X-ray structures of $[Pt(1\text{-MeC-}N3)_3Cl]^+$ and $[Pt(1\text{-MeC-}N3)_4]^{2+}$ compounds, **279**, 152

Methylene-bridged complexes

dialkyl-μ-ethylidene-μ-methylene-bis(pentamethylcyclopentadienyl)-dirhodium complexes: synthesis, spectra and decomposition reactions, **280**, 62

Methyl ligation

a theoretical study of the methyl ligation in tetraaza macrocycle nickel complexes which model the acetyl-CoA synthase active site, **273**, 379

Methyl migration

monovalent copper as a potential catalyst for formation of acetaldehyde via the migration of methyl radicals to the coordinated carbonyl in the complex $(CO)Cu^{II}–CH_3^+$, **270**, 440

Methylphosphonylphosphate complexes

stability of binary and ternary copper(II) complexes of the diphosphate analogue, methylphosphonylphosphate, in aqueous solution, **273**, 101

Methyl transfer

alkyl exchange between aluminum trialkyls and zirconocene dichloride complexes — a measure of electron densities at the Zr center, **270**, 376

Methyl-viologen

photoreduction of methyl-viologen by zeolite-Y stabilized silver clusters, **269**, 253

Microemulsions

hydrolysis of nerve agent simulants by synergistic effects of tetraminecobalt(III) and microemulsions, **269**, 287

Microporous material

synthesis and gas-occlusion properties of dinuclear molybdenum(II) dicarboxylates (fumarate, terephthalate, *trans-trans*-muconate, pyridine-2,5-dicarboxylate, and *trans*-1,4-cyclohexanedicarboxylate), **283**, 268

Misdirected ligands

misdirected π-donor ligands: $(\eta^5\text{-}C_5H_5)_2Zr(Cl)(SR)$ with sterically more demanding R groups have lower rotational barriers, **270**, 227

Mixed ligand complexes

stability of binary and ternary copper(II) complexes of the diphosphate analogue, methylphosphonylphosphate, in aqueous solution, **273**, 101

ternary complexes in solution. Intramolecular stacking interactions in mixed ligand complexes formed by copper(II), 2,2′-bipyridyl or 1,10-phenanthroline and a pyrimidine-nucleoside 5′-diphosphate (CDP^{3-}, UDP^{3-}, $dTDP^{3-}$), **283**, 193

Mixed-metal complexes

on the mechanism of silylformylation catalyzed by Rh–Co mixed metal complexes, **270**, 279

generation of mixed iron chalcogenide clusters $MFeX_n^+$ (M = V, Fe; X = O, S; $n = 2, 3$) in the gas phase, **273**, 111

Mixed valence complexes

synthesis and properties of the unsymmetrical binuclear complexes: $Na[(CN)_5Fe(dtdp)Ru(NH_3)_5]6H_2O$ and $K[(CN)_5Ru(dtdp)Ru(NH_3)_5]6H_2O$, dtdp = 4,4′-dithiodipyridine, **267**, 93

assembly of a mixed-valence Cu(I/II) system coupled by multiple hydrogen bonding through tetrahydroxybenzoquinone, **268**, 317

the cyanide ligand as an efficient bridge in mixed-valence complexes, **269**, 246

resonance Raman investigation of mixed-valence dinuclear and trinuclear complexes of Ru, Os and Re, **275–276**, 380

mixed valence, tetradisordered structure of iodine-doped bismuth(III) diphthalocyanine, **281**, 195

synthesis and characterization of mixed valence μ-alkoxo-diiron-(II,III) complexes with an unsymmetric dinucleating ligand, **283**, 124

Mixed vesicles

spectroscopic characterization of fluorinated/hydrogenated mixed vesicles containing fluorinated Mn(III)-porphyrin, **272**, 274

MNDO method

simulation of the molecular and electronic structure of heterofullerenes $C_{55}Y_5$ (Y = Si, Ge, Sn, B, Al, N, P, SiH, GeH, SnH) and their η^5-π-complexes with Li by the MNDO method, **280**, 219

MO calculations

ternary copper(II) complexes with the versatile 4,7-dihydro-5-methyl-7-oxo-[1,2,4]triazolo[1,5-*a*]pyrimidine ligand, **274**, 53

Model heme complexes

ROESY spectra of paramagnetic model heme complexes: unambiguous differentiation of cross-peaks due to chemical exchange and the nuclear Overhauser effect in the same molecule, **272**, 204

Modified resins

a catalytic activity of glass beads, silica gels and anion-exchange resins modified with metal-porphines in oxidative reactions of ascorbic acid, **283**, 44

Molecular design

role of the intramolecular hydrogen bond and ligand rigidity in the complexation of trifluoroacetylcycloalkanones with lanthanides: novel strategy for the design of organic ligands of high selectivity, **267**, 201

Molecular dynamics

molecular dynamics of *N,N′*-di-*p*-fluorophenyltriazenido complex of platinum(II), **280**, 238

Molecular mechanics

structural preferences of η^4-dienecyclopentadienyl complexes: molecular mechanics, molecular orbital and crystallographic studies, **275–276**, 263

ESR and NMR investigation of structure and dynamics of the Cu(II) complex of melanostatin in DMSO solution, **275–276**, 274

Molecular orbital calculations

hydrogen site exchange in $CpIr(PR_3)H_3^+$ complexes, **272**, 95

Molecular vibrations

Mössbauer spectroscopy with synchrotron radiation: a new technique entering biological inorganic chemistry, **275–276**, 334

Molybdacarborane

oxidation of tricarbonylmolybdacarborane. 2. Oxidative decarbonylation with various sulfur donors, **283**, 145

Molybdate complexes

synthesis and molecular structure of the molybdate-bridged cyclic imido complex $[Mo(S_2CNEt_2)_2(NR)(\mu\text{-}MoO_4)]_2$ (R = 2,6-$^iPr_2C_6H_3$), **274**, 243

preparation of tetrabutylammonium octa-μ_3-bromohexa(trifluoromethanesulfonato)octahedro-hexamolybdate(2−), $(Bu_4N)_2[Mo_6Br_8^i(CF_3SO_3)_6^a]$ and other derivatives containing the $Mo_6Br_8^{4+}$ core, **278**, 241

Molybdenum complexes

synthesis and characterization of molybdenum(II) and molybdenum(V) monothiocarbamate complexes. Crystal and molecular structures of tetrakis(isopropylmonothiocarbamato)dimolybdenum(II) and oxobis[di(oxoisopropylmonothiocarbamato)-molybdenum(V)], **267**, 101

synthesis and characterization of molybdenum(VI)-dioxo complexes containing both coordinated thiolate and carboxylate groups. Reactions with their own free ligands, **268**, 145

reactivity of molybdenum complexes containing both NHNPhR and NNPhR ligands (R = Ph, Me) toward hydridotris(pyrazolyl)-borate, Tp and hydridotris(3,5-dimethylpyrazolyl)borate, Tp*, ligands. X-ray crystal structure of $[TpMo(NNMePh)_2Cl]$, **268**, 199

tripod ligands of constitutional C_1 symmetry containing a deliberate set of mixed P/S donor groups: synthesis and coordination chemistry, **269**, 143
substituent effects of *cis*-dioxobis(dithiocarbamato)molybdenum(VI) on redox properties: redox potentials for one-electron reduction and second-order rate constants for oxygen atom transfer, **269**, 260
salicylaldehyde-2-mercaptoanilato(2−)-2-hydroxyphenyl-benzthiazolato-nitrosyl-molybdenum(II) — a chelate which simultaneously contains an azomethine ligand and its cyclic oxidation product, **269**, 350
catalytic reduction of acetylene in the presence of molybdenum and iron clusters, including FeMo cofactor of nitrogenase, **270**, 221
chemistry of the azine phosphine ligand *Z,E*-$PPh_2CH_2C(Bu^t)$=N–N=$CMe(C_6H_4NO_2$-4): crystal structure of $[Mo(CO)_4$-{$PPh_2CH_2C(Bu^t)$=N–N=$CMe(C_6H_4NO_2$-4)}], **270**, 312
modeling for the active site nitrate reductase. Oxidation of the complex $[Mo^{V}O(O_2CC(S)CH_3Ph)_2]^-$ by nitrate and nitrite in methanol, **271**, 174
synthesis and reactivity of bis(ethylene)-phosphite complexes of molybdenum(0), **272**, 125
synthesis of chloro and methyl imido cyclopentadienyl molybdenum and tungsten complexes. X-ray molecular structures of $[WCp^*Cl_3(N^tBu)]$, $[MoCp^*ClMe_2(N^tBu)]$ and $[WCp^*ClMe_2(N^tBu)]$, **273**, 270
quadruply bonded complexes containing η^2- or η^3-tridentate phosphine ligands; syntheses, structures and $^{31}P\{^1H\}$ NMR spectra of $Mo_2Cl_4(\eta^3$-etp)(L) and $Mo_2Cl_4(\eta^2$-etp)$_2$ (L = CH_3OH, PMe_3 and PEt_3; etp = $Ph_2PCH_2CH_2P(Ph)CH_2CH_2PPh_2$), **274**, 7
synthesis and molecular structure of the molybdate-bridged cyclic imido complex $[Mo(S_2CNEt_2)_2(NR)(\mu$-$MoO_4)]_2$ (R = 2,6-$^iPr_2C_6H_3$), **274**, 243
structural preferences of η^4-dienecyclopentadienyl complexes: molecular mechanics, molecular orbital and crystallographic studies, **275–276**, 263
preparation and electrochemistry of six-coordinate monooxo molybdenum(VI) complexes containing bidentate catecholate and tridentate NOS-donor Schiff base ligands, **278**, 1
synthesis and crystal structure of a new salt of the complex anion $[(TCNQ)_3]^{2-}$ $[Mo_2(\mu_2$-$S_2)_2(S_2CNMe_2)_4](TCNQ)_3$, **278**, 108
molybdenum complexes of the anticancer drug 6-mercaptopurine, **279**, 136
solid-state synthesis of cluster compound $[MoCu_3OS_3(PPh_3)_3$-$(CH_3COO)]$ with its structure, reactivity and optical non-linearities, **279**, 172
reactivities of the coordinated organonitriles in molybdenum(0) and tungsten(0) phosphine complexes: protonation of the nitrile carbon and cleavage of the C≡N triple bond, **280**, 163
synthesis and characterization of arene, mono- and dihydrido-arene, monohydrido-cyclopentadienyl, and phosphite complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$, **281**, 153
investigations into the engineering of inorganic/organic solids: hydrothermal synthesis and crystal structures of aromatic ammonium sodium hydrogen phosphomolybdates(VI), **282**, 10
preparation and characterization of twin-cubane-type clusters with molybdenum–copper–sulfur cores: X-ray structure determination of $[\{(NH_3)_6(H_2O)(OH)Mo_3CuS_4(O)\}_2]Cl_4 \cdot 8H_2O$, **282**, 50
mononuclear, dinuclear, and polynuclear dioxomolybdenum(VI)–tridentate Schiff base complexes: crystal structures and properties, **282**, 209
synthesis of double incomplete-cubane clusters $[PPh_4]_2[(\eta^5$-$C_5Me_5)MS_3Cu_3X_3]_2$ (M = Mo, W; X = Cl, Br) and reactions with PPh_3, dppm (1,2-bis(diphenylphosphino)methane), and $AsPh_3$, **283**, 136
Molybdenum compounds
X-ray absorption fine structure study of the atomic and electronic structure of molybdenum disulfide intercalation compounds with transition metals, **280**, 211
Molybdenum(II) dicarboxylate
synthesis and gas-occlusion properties of dinuclear molybdenum(II) dicarboxylates (fumarate, terephthalate, *trans-trans*-muconate, pyridine-2,5-dicarboxylate, and *trans*-1,4-cyclohexanedicarboxylate), **283**, 268
Molybdotellurate complexes
Anderson–Evans type molybdotellurates with diazolium cations: preparation, structure and properties of $(H_2imz)_6[TeMo_6O_{24}] \cdot 4H_2O$ and $(H_2pyz)_6[TeMo_6O_{24}] \cdot Te(OH)_6$, **277**, 139
Mononuclear complexes
structures of $HRh(PMe_2Ph)_4$ and $HRh(PPh_2Me)_4$, **269**, 162
Monophosphate complexes
interaction of mono- and triphosphate anions with a protonated polyaza macrocycle and its Cu(II) complexes, **270**, 207
Mössbauer spectroscopy
reductive changes to polynuclear iron(III) clusters in iron-loaded human spleen tissue, **267**, 7
Mössbauer spectroscopy with synchrotron radiation: a new technique entering biological inorganic chemistry, **275–276**, 334
a synthetic iron-containing prismane complex: peculiar electron-delocalization and spin-coupling properties deduced from Mössbauer and magnetization studies, **283**, 111
MO study
the role of cyclo-triphosphorus (P_3) in-plane σ orbitals in coordinating transition metal fragments, **275–276**, 366
Multi-copper enzyme
anions binding to bilirubin oxidase from *Trachyderma tsunodae* K-2593, **273**, 204
Multidentate amine complexes
synthesis of new copper(II) complexes of polydentate ligands containing pyridyl pendant arms. Crystal structure of [*N,N,N',N'*-tetrakis(2-pyrid-2-yl-ethyl)1,2-ethylenediaminecopper(II)]-bis(hexafluorophosphate) · 1.5 acetone, **278**, 237
Multiple bonded complexes
the synthesis and characterization of α-$Re_2Cl_4(depe)_2$ and α-$W_2Cl_4(depe)_2$, **271**, 49

Nafion complexes
EXAFS studies on Mn(III) tetraphenylporphyrin halides in Nafion, **268**, 231
Nanotubes
the immobilisation of proteins in carbon nanotubes, **272**, 261
N-donor ligand complexes
nucleophilic attack within Ge, Sn and Pb complexes containing $Me_2N(CH_2)_3$– as a potential intramolecular donor ligand, **275–276**, 203
Nerve agent simulant
hydrolysis of nerve agent simulants by synergistic effects of tetraminecobalt(III) and microemulsions, **269**, 287
Neutron diffraction
protonation of Fischer-type alkylidyne carbonyltungsten complexes. Structural comparison of alkylidyne and alkylidene metal complexes, including a neutron diffraction study of $[W(CHCH_3)$-$Cl_2(CO)(PMe_3)_2]$, **279**, 7
Neutron scattering
taranakite – the mineral with the longest crystallographic axis, **269**, 47
spectroscopic characterization of fluorinated/hydrogenated mixed vesicles containing fluorinated Mn(III)-porphyrin, **272**, 274
N-heterocycles complexes
the effect of net charge and π-backbonding contribution of ML_5 fragments containing Fe(II), Ru(II), Os(II), Co(III) and Rh(III) on the ^1H NMR shifts of coordinated *N*-heterocycles, **267**, 11
Nickel-catalysed Grignard reagents

enantioselective catalysis. XVI: regio- and enantioselectivity in nickel-catalysed cross-coupling reactions of allylic substrates with Grignard reagents, **269**, 34

Nickel complexes

effect of substitution on packing in *trans*-di(*Z*-chlorobenzoato-*O*)-di(1,3-diaminopropane-*N*,*N*′)nickel(II) (*Z* = 2, 3 or 4) complexes, **267**, 249

synthesis, electrical properties of $[Ni(dmit)_2]^-$ complexes and X-ray crystal structure of $[EDA][Ni(dmit)_2]$ (dmit = 1,3-dithiol-2-thione-4,5-dithiolate), **268**, 103

electronic spectra and structures of Cu^{2+} and Ni^{2+} complexes with dioxotetraazacycloalkanediacetates and their diester derivatives, **269**, 224

{*catena*-($\mu_{N,S}$-NCS)[Ni(Medien)(NCS)]}: a new 1D nickel(II) complex with bridging thiocyanato ligands in *cis* position. Crystal structure and magnetic properties, **269**, 313

infrared spectroelectrochemical and electrochemical kinetics studies of the reaction of nickel cluster radicals $[Ni_3(\mu_2\text{-dppm})_3(\mu_3\text{-L})(\mu_3\text{-I})]^+$ (L = CNR, R = CH_3, i-C_3H_7, C_6H_{11}, $CH_2C_6H_5$, t-C_4H_9, 2,6-$Me_2C_6H_3$; L = CO) with carbon dioxide, **270**, 111

transition metal complexes. IX. Nickel complexes with a chiral azaphosphole ligand, **270**, 273

synthesis, spectroscopic and electrochemical study of nickel(II) complexes with tetradentate asymmetric Schiff bases derived from salicylaldehyde and methyl-2-amino-1-cyclopentenedithiocarboxylate, **271**, 83

synthesis, characterization and reactivity of bis(diphenylphosphino)-amine nickel complexes, **271**, 112

nickel(II) 1,1-dithiolates and related complexes. Solid-state and solution structural evidence for mixed-ligand complexes with substituted ethylenediamines, **271**, 195

Ni(II) and Cu(II) Schiff base complexes with an extended H-bond network, **272**, 283

synthesis, crystal structure and magnetic properties of bis[μ-bis(3-(pyridin-2-yl)-1,2,4-triazole-*N*′,*N*1,*N*2)] bis[triaqua nickel(II)] tetranitrate, **273**, 54

synthesis and coordinating properties of the functionalized macrocyclic ligand 1,4-bis(1-methylimidazol-2-ylmethyl)-7-carboxymethyl-1,4,7-triazacyclononane. Crystal structures of cobalt(III), nickel(II) and zinc(II) complexes, **273**, 151

novel oxidation and reduction products of the neutral nickel-dithiolene $[Ni(Pr^i_2timdt)_2]$ (Pr^i_2timdt is the monoanion of 1,3-diisopropylimidazolidine-2,4,5-trithione), **273**, 175

formation and reactivity of paramagnetic organometallic nickel complexes of 21-oxa- and 21-selenaporphyrins — ^1H NMR and EPR investigations, **273**, 184

the oxidation of nickel^{II}(1,4,8,11-tetraazacyclotetradecane) by hydroperoxyl radicals: a pulse radiolysis study, **273**, 266

metal complexes of $P(Ph_2)CH_2(Ph_2)PSe$. Crystal structure of $[Ni\{P(Ph_2)CH_2(Ph_2)PSe\}_2]Cl_2 \cdot 2EtOH$, **273**, 320

probing the metal site in *Rhus vernicifera* stellacyanin by Ni(II) substitution and paramagnetic NMR spectroscopy, **273**, 367

a theoretical study of the methyl ligation in tetraaza macrocycle nickel complexes which model the acetyl-CoA synthase active site, **273**, 379

synthesis and properties of a new tetraaza macrocyclic ligand containing one N-acetic acid pendant arm and its copper(II) and nickel(II) complexes, **274**, 24

structural and electronic studies of the coordination of 6-(thiazol-2-yl)-2,2′-bipyridine and related systems to Fe(II), Co(II) and Ni(II), **274**, 32

nickel(II), palladium(II), platinum(II) and platinum(IV) complexes of *cis*-1,2-bis(diphenylphosphino)ethene and nitrogen-containing heterocycles or sulfur ligands, **274**, 143

potentiometric and spectroscopic studies on the copper(II) and nickel(II) complexes of tripeptides of methionine, **275–276**, 440

preparation of soluble tetrabenzoporphyrins with substituents at the peripheral positions, **277**, 151

ligand exchange reactions of $ReCl_4(PPh_3)_2$, $ReOCl_3(PPh_3)_2$ and $NiCl_2(PPh_3)_2$ with acetylacetone thiobenzoylhydrazone. An unexpected template reaction at the ReO^{3+} core. Molecular structures of the resulting oxorhenium(V) and nickel complexes, **277**, 219

synthesis, structural characterization and magnetic behaviour of two new $(\mu_{1,3}\text{-}N_3)_2$ dinuclear nickel compounds: study of the magnetic transition at 235 K for the *cis*-$(\mu_{1,3}\text{-}N_3)_2[Ni(dl\text{-}cth)]_2(ClO_4)_2$ compound, **278**, 43

some hexadentate Ni(II)-edta-type complexes containing five-membered diamine rings. The molecular and crystal structure of the *trans*(O_5) isomer of barium(ethylenediamine-*N*,*N*′-diacetato-*N*,*N*′-di-3-propionato)nickelate(II) hexahydrate, *trans*(O_5)-$Ba[Ni(eddadp)] \cdot 6H_2O$, and strain analysis of edta-type chelates in relation to their octahedral distortion, **278**, 66

bis(1,2-diselenooxalato)nickelate(II) and -cuprate(II) — the first diselenooxalate complexes, **278**, 118

the 1,8-bis(2′-pyridyl)-3,6-dithiaoctane complex of nickel(II): X-ray crystal structure and borohydride adduct formation, **278**, 217

preparation and some properties of perfluoroalkoxy-substituted phthalocyanine complexes of iron(III), nickel(II) and zinc(II), **278**, 253

synthesis and metal complexes of a chelating ligand containing thiomorpholine and pyridine: 2,6-bis(thiomorpholinomethyl)pyridine, **279**, 221

synthesis and characterization of nickel(II) complexes of di-*N*-alkylated 14-membered tetraaza macrocycles containing cyclohexane rings via regioselective alkylation reaction, **279**, 238

synthesis and characterization of homologous nickel(II) and palladium(II) complexes with biphosphine monoxide ligands $Ph_2P(CH_2)_nP(O)Ph_2$ (n = 1–3) and pTol$_2$P(CH$_2$)P(O)pTol$_2$, **280**, 150

bimetallic cyclooligosiloxanolate complexes of copper and nickel, **280**, 282

complexes of the bis(trimethylsilyl)-benzamidinato ligand '*siam*': synthesis and X-ray structures of (*siam*)$_2$M, (*siam*)(*siam*H)MX (M = Ni, Pd), (*siam*)$_2$MnI and (*siam*)ReO$_3$, and their reactivity towards CO_2, **281**, 181

preparation and spectroscopic properties of five 7,16-bis(azophenyl)-tetraaza[14]annulene nickel(II) complexes, **281**, 190

the effect of aromatic–aromatic interactions on the crystallization of helical nickel(II) complexes, **282**, 71

the phenomenon of conglomerate crystallization. Part 51. The crystallization behavior of nickel dinitro amine complexes. *Trans*-$[Ni(3,2,3\text{-}tet)(NO_2)_2] \cdot H_2O$ (**I**), *mer*-[Ni(N5-Me-1,5,9-triazanonane)$(NO_2)_2$ (**II**), *cis*-$[Ni(tren)(NO_2)_2] \cdot H_2O$ (**III**) and *cis*-$[Ni(edda)(H_2O)_2] \cdot H_2O$ (**IV**) and control of the bonding mode of the nitrite ligand by the choice of the amine, **282**, 96

the synthesis and characterization of Ni, Pd and Pt maleonitriledithiolate complexes: X-ray crystal structures of the isomorphous Ni, Pd and Pt ($Ph_2PCH_2CH_2PPh_2$)M(maleonitriledithiolate) congeners, **282**, 155

binuclear nickel(II) and copper(II) complexes of amide-containing two-armed and macrocyclic ligands, **282**, 222

synthesis, structure and characterization of dinuclear nickel(II) complexes containing the derivatives of *N*,*N*,*N*′,*N*′-tetrakis[(2-benzimidazolyl)methyl]-2-hydroxy-1,3-diaminopropane and an acetate or a propionate ion, **283**, 118

Nickel(III) complexes

synthesis and properties of nickel(III) complexes of hexaazamacrocyclic ligands, **269**, 337

N–I compounds

the cations $[ICNI]^+$ and $[H_3N\text{–}NH_3]^{2+}$ — natural bond orbital analysis and some valence bond considerations, **269**, 1

Niobium complexes

eight-coordinate chelate complexes of zirconium(IV) and niobium(IV): X-ray diffractometric and EPR investigations, **270**, 177
synthesis of a new series of Nb(IV) β-diketonates with the $Nb_2S_4^{4+}$ core: crystal structure of $Nb_2S_4(CF_3COCHCOCH_3)_4$, **271**, 222
synthesis and properties of $[M_6X_{12}(C_3H_7NO)_6]X_2$ (M = Nb, Ta; X = Cl, Br), **271**, 239
structural characterization of an eight-coordinate Nb(IV)–edta complex, $[Nb(edta)(H_2O)_2]\cdot 2H_2O$, **274**, 103
complexes of niobium and tantalum with chloride, sulfide and oxide ligands: synthesis and crystal structure of the (tetrachloro)bis-(η^2-disulfido)(μ-oxo)(μ-η^2:η^2-disulfido) bismetallates $[Cl_2(S_2)Nb(\mu\text{-}O)(\mu\text{-}S_2)Nb(S_2)Cl_2]^{2-}$ and $[Cl_2(S_2)Ta(\mu\text{-}O)(\mu\text{-}S_2)Ta(S_2)Cl_2]^{2-}$ as their tetraphenylphosphonium salts, **274**, 251
tetravalent niobium alkyls and enolate stabilized by anionic organic amides, **280**, 143

Nitrate complexes
the chemistry of dinitrato-2,2′-bipyridinecopper(II): preparation and characterization of binuclear complexes having a $Cu(\mu\text{-}OH)_2Cu$ and $Cu(\mu\text{-}N_3)_2Cu$ core, **278**, 143

Nitrate reductases
modeling for the active site nitrate reductase. Oxidation of the complex $[Mo^{V}O(O_2CC(S)CH_3Ph)_2]^-$ by nitrate and nitrite in methanol, **271**, 174

Nitration
electrophilic substitution of metal-coordinated pyrazole: nitration, sulfonation and bromination of $[Co(NH_3)_5(pyzH)]^{3+}$ (pyzH = pyrazole), **277**, 89

Nitric oxide complexes
application of the electrocatalytic reduction of nitric oxide mediated by ferrioxamine B to the determination of nitric oxide concentrations in solution, **273**, 316
adducts of nitric oxide with cobaltous tetraphenylporphyrin and phthalocyanines: potential nitric oxide sorbents, **277**, 1

Nitride complexes
nephelauxetic series of in-plane ligands for square-pyramidal nitridochromium(V) complexes as studied by means of ESR spectroscopy, **281**, 207

Nitrido complexes
synthesis of nitridoosmium(VI) dithiolene complexes, **267**, 335
transition metal complexes with sulfur ligands. Part CXXVII. Azido, halido and nitrido ruthenium complexes with sulfur-rich coordination spheres, **269**, 63
an electron paramagnetic resonance study of tetra-*n*-butylammonium-tetrabromo-nitridorhenate(VI), $[(n\text{-}C_4H_9)_4N][ReNBr_4]$, **271**, 199

Nitrite reductase
spectroscopic distinction between two Co(II) ions substituted for types 1 and 2 Cu in nitrite reductase, **275–276**, 289

Nitro complexes
the phenomenon of conglomerate crystallization. Part 51. The crystallization behavior of nickel dinitro amine complexes. *Trans*-$[Ni(3,2,3\text{-}tet)(NO_2)_2]\cdot H_2O$ (**I**), *mer*-[Ni(N5-Me-1,5,9-triazanonane)$(NO_2)_2$ (**II**), *cis*-$[Ni(tren)(NO_2)_2]\cdot H_2O$ (**III**) and *cis*-$[Ni(edda)(H_2O)_2]\cdot H_2O$ (**IV**) and control of the bonding mode of the nitrite ligand by the choice of the amine, **282**, 96

Nitrogenase
protonation of cyclopropene complexes of platinum(0) and the reduction of cyclopropene by nitrogenases, **280**, 193

Nitrogen base complexes
crystal structure and ^{63}Cu NQR of bis[1-aza-2-(1-pyrrolidinyl)]-copper(I) chloride and bromide, **271**, 231

Nitrogen fixation
transformations of molecular nitrogen into aromatic amines under the action of titanium compounds, **280**, 264

Nitrogen heterocycle complexes
synthesis and crystal structure of a new isomer of the μ-oxo-bis[dichlorooxobis(pyridine)rhenium(V)] complex $\{OReCl_2py_2\}_2O$, **279**, 159

Nitrogen ligand complexes
tetrahedral coordination by a seven-membered chelate ring, **275–276**, 279
thermodynamics of the binding of nitrogenous ligands to β-trifluoromethylcobinamide and β-cyanomethylcobinamide, **279**, 178

Nitrogen transfer
nitrogen atom transfer and redox chemistry of terpyridyl phosphoraniminato complexes of osmium (IV), **270**, 511

Nitronyl nitroxide complexes
transition metal complexes with nitronyl nitroxides. Part 1. Crystal structure and properties of square-planar *trans* radical–copper(II)–radical complexes, **279**, 69
transition metal complexes with nitronyl nitroxides. Part 2. Crystal structure and magnetism of manganese(II) compounds including radicals and their reduced derivatives, **279**, 76
structure and magnetic properties of a novel chloro-bridged polymeric cadmium(II) complex with pyridyl-substituted nitronyl nitroxide, **279**, 233

Nitrosamine complexes
nitrosamine complexes of $SbCl_5$ and $CuCl_2$, **269**, 7

Nitrosyl-bridged complexes
redox behavior of a binuclear ruthenium complex having a di-μ-nitrosyl ligand, $[\{Ru(acac)_2\}_2(\mu\text{-}NO)_2]$ (acac = acetylacetonato), **269**, 342

Nitrosyl complexes
interaction between the NO^+ ligand and *trans* π-donor ligands in chloro(2-methyl-8-quinolinolato)nitrosyl(8-quinolinolato)-ruthenium(II) complexes, **267**, 147
the synthesis and characterization of a technetium nitrosyl complex with *cis*-{2-pyridyl,diphenylphosphine} coligands. The X-ray crystal structure of $[TcCl_2(NO)(pyPPh_2\text{-}P,N)(pyPPh_2\text{-}P)]$, **267**, 319
synthesis and structures of the cuboidal iron–sulfur–nitrosyl–phosphine clusters $[Fe_4S_3(NO)_4(PR_3)_3]^{0,1+}$ (R = Et, Pr^i, C_6H_{11}), **270**, 46
η^3-allyl–cobalt(II) complexes, **270**, 60
the reaction chemistry of a technetium(I) nitrosyl complex with potentially chelating organohydrazines: the X-ray crystal structure of $[TcCl_2(NO)(HN{=}NC_5H_4N)(PPh_3)]$, **279**, 206
electrophilic abstractions on $Os(H)_2X(NO)L_2$: selective access to new unsaturated cationic Os(II) hydrides, **280**, 125

Nitrous oxide complexes
preparation of trimethylsilyl alkyne complexes of bis(pentamethylcyclopentadienyl)zirconium, $(\eta^5\text{-}C_5Me_5)_2Zr(RC{\equiv}CSiMe_3)$, and their regioselective reactions with nitrous oxide, **270**, 399

NMR
NMR conformational study of diamagnetic complexes of some triazatriacetate macrocycles, **273**, 288
Boc-Glu-Thr-Ile-His-OMe/Zn^{2+} in terms of the zinc-binding sites of proteases, **283**, 243

^{11}B NMR
^1H and ^{11}B NMR evidence for specific binding of borate ion to cytochrome *c*, **273**, 388

^{13}C-CP/MAS NMR
synthesis and gas-occlusion properties of dinuclear molybdenum(II) dicarboxylates (fumarate, terephthalate, *trans-trans*-muconate, pyridine-2,5-dicarboxylate, and *trans*-1,4-cyclohexanedicarboxylate), **283**, 268

^{113}Cd NMR
the use of ^{113}Cd NMR chemical shifts as a structural probe in tetrathiolate metalloproteins, **273**, 279

NMR relaxation
ESR and NMR investigation of structure and dynamics of the Cu(II) complex of melanostatin in DMSO solution, **275–276**, 274

selective versus non-selective T_1 experiments to determine metal–nucleus distances in paramagnetic metalloproteins, **275–276**, 373

NMR spectroscopy
oligonuclear ferrocenolato and 1,1′-ferrocenediolato derivatives of phosphorus: synthesis, structure and electrochemical behaviour, **273**, 24
Pd(gly-L-his-L-lys)Cl: solution structure and ternary complex formation with mono- and tetranucleotides, **273**, 31
interaction of manganese(II), cobalt(II) and nickel(II) with DNA oligomers studied by ^{1}H NMR spectroscopy, **273**, 41
nucleophilic attack within Ge, Sn and Pb complexes containing $Me_2N(CH_2)_3$– as a potential intramolecular donor ligand, **275–276**, 203
hemoprotein models based on a covalent helix–heme–helix sandwich 4. Discrimination of paramagnetic Fe(III)–Mimochrome I Δ and Λ isomers by NMR spectroscopy, **278**, 76
NMR evidence for formation of new alcohol rhenium complexes as intermediates in ionic hydrogenations of carbonyl groups with systems composed of $ReH_2(NO)(CO)(PR_3)_2$ (R = Pr^i, CH_3, OPr^i) and CF_3COOH, **278**, 122

N,O-heterocycles
palladium-catalyzed seven-membered N,O-heterocycle ring formation, **270**, 123

Non-conjugated dienes
zirconium-catalyzed and zirconium-promoted cyclization reactions of non-conjugated dienes with alkylmagnesium halides to give cycloalkylmethylmagnesium derivatives, **280**, 8

Non-heme complexes
sterically hindered iron(II) complex of a new tripodal polyimidazole ligand: structure and reactivity toward superoxide, **275–276**, 234

Nonlinear optics
solid-state synthesis of cluster compound $[MoCu_3OS_3(PPh_3)_3(CH_3COO)]$ with its structure, reactivity and optical non-linearities, **279**, 172
preparation and properties of ferrocenyl bimetallic compounds for non-linear optics, **279**, 243

Non-oxo complexes
the preparation and synthetic potential of chlorovanadium(V and IV) complexes supported by enamines and bis(enamines), **267**, 229

N-protected aminoacids–Me(II) complexes
palladium(II) complexes of *N*-sulfonyl-asparagine and glutamine. Evidence for metal coordination of the deprotonated amide nitrogen of the side-chain, **273**, 397

N,S-bidentate ligand complexes
novel palladium(II) allyl complexes with nitrogen–sulfur donor bidentate ligands. Mechanism of allyl amination of $[Pd(\eta^3\text{-allyl})(N\text{-}SR)]ClO_4$ (allyl = C_3H_5; N-SR = C_5H_4N-2-CH_2SR, R = C_6H_5, C_2H_5) in the presence of activated olefins. X-ray structure determination and fluxional behavior, **275–276**, 385

N,S-ligand complexes
the effect of aromatic–aromatic interactions on the crystallization of helical nickel(II) complexes, **282**, 71

N-substituted aminomethyl complexes
crystal and molecular structure of protonated (*N*-propyl)aminomethyl ferrocene, a proton-sensitive redox-responsive fragment, **267**, 177

Nuclear inelastic scattering
Mössbauer spectroscopy with synchrotron radiation: a new technique entering biological inorganic chemistry, **275–276**, 334

Nuclear magnetic resonance studies
a tripod gadolinium(III) poly(amino carboxylate) relevant to magnetic resonance imaging: structural and dynamical ^{17}O NMR and ^{1}H NMRD studies, **275–276**, 15

Nuclear resonant forward scattering
Mössbauer spectroscopy with synchrotron radiation: a new technique entering biological inorganic chemistry, **275–276**, 334

Nuclease enzymes
magnesium activation of nuclease enzymes — the importance of water, **275–276**, 24

Nucleic acid complexes
synthesis of binuclear platinum(II) organometallic complexes as potential cross-linking reagents, **275–276**, 141
interaction of organometallic cationic complex ions containing terpyridine ligands with nucleic acids: an investigation on aggregative phenomena, **275–276**, 242

Nucleobase complexes
a bis(9-methyladeninium) complex of Hg(II) with a highly irregular coordination geometry: $[Hg(9\text{-MeAH-}N7)_2(H_2O)(NO_3)_3]ClO_4$, **267**, 87
tris- and tetrakis(1-methylcytosine) complexes of Pt(II): syntheses and X-ray structures of $[Pt(1\text{-MeC-}N3)_3Cl]^+$ and $[Pt(1\text{-MeC-}N3)_4]^{2+}$ compounds, **279**, 152
synthesis and structure of (1,3-dimethyluracil-5-yl)mercury(II) complexes with aromatic nitrogen donor ligands, **282**, 237
interactions between $[AuX_4]^-$ (X = Cl, CN) and cytosine and guanine model nucleobases: salt formation with (hemi-) protonated bases, coordination, and oxidative degradation of guanine, **283**, 223
can Pt(IV)–amine complexes act as 'prodrugs'?, **283**, 251

Nucleophilic and electrophilic attack
nucleophilic attack at the five-coordinate anion $[Mn(CO)_3(3,5\text{-di-tert-butyl-catecholate})]^-$ controlled by electronic and steric effects, **268**, 1

Nucleoside complexes
comparison of the reactivity of *cis*-$[RuCl_2(DMSO)_4]$ and *trans*-$[RuCl_2(DMSO)_4]$ towards nucleosides, **281**, 10

Nucleotide complexes
ternary complexes in solution. Intramolecular stacking interactions in mixed ligand complexes formed by copper(II), 2,2′-bipyridyl or 1,10-phenanthroline and a pyrimidine-nucleoside 5′-diphosphate (CDP^{3-}, UDP^{3-}, $dTDP^{3-}$), **283**, 193

Olefin complexes
synthesis, solution behavior and molecular structure of 16-electron *closo*-2-(Ph_3P)-1-N,2-$[\mu\text{-}(\eta^2\text{-}CH_2CH{=}CH_2)]$-1-N-$(\sigma\text{-}CH_2CH{=}CH_2)$-2,1-$RhCB_{10}H_{10}$, the first η^2-olefinic monocarbon metallacarborane, **280**, 233

Olefin oxidation
catalysis of aerobic olefin oxidation by a ruthenium perhaloporphyrin complex, **270**, 433

Oligonucleotide complexes
monofunctional *trans*-$Pt^{II}(NH_3)_2$ modification of pyrimidine-rich deoxyoligonucleotides: direct platination and use of a protective group, **269**, 135
Pd(gly-L-his-L-lys)Cl: solution structure and ternary complex formation with mono- and tetranucleotides, **273**, 31
interaction of manganese(II), cobalt(II) and nickel(II) with DNA oligomers studied by ^{1}H NMR spectroscopy, **273**, 41

Oligopyridine complexes
4′-(4-fluorophenyl)-2,2′:6′,2″-terpyridine – an aryl-2,2′:6′,2″-terpyridine with an electron-releasing aryl substituent, **275–276**, 359

One-dimensional chain complexes
synthesis, structure and magnetic properties of one-dimensional azide-bridged manganese(III) uniform chain complex Mn(salpn)N_3, **271**, 99

O–O distances
role of the intramolecular hydrogen bond and ligand rigidity in the complexation of trifluoroacetylcycloalkanones with lanthanides: novel strategy for the design of organic ligands of high selectivity, **267**, 201

Organic hydroperoxides
an ab initio MO-LCAO investigation of the electronic structure of organic hydroperoxides and platinum(II) hydroperoxo complexes:

a contribution to the knowledge of the mechanism of olefin epoxidation, **270**, 479
gas-phase positive and negative ion chemistry of methyl hydroperoxide, **275–276**, 192
Organoammonium salt complexes
investigations into the engineering of inorganic/organic solids: hydrothermal synthesis and crystal structures of aromatic ammonium sodium hydrogen phosphomolybdates(VI), **282**, 10
Organocobalt complexes
alkylcobalt chelates with Schiff bases derived from a β-diketone bearing both alkyl and aryl groups, **270**, 169
rate constants and activation parameters for organo–cobalt porphyrin bond homolysis from NMR relaxation times, **270**, 197
Organodiazene complexes
the synthesis and characterization of $[Tc(SPh)_2(N{=}NC_5H_4N)(HN{=}NC_5H_4N)]_2$, a paramagnetic dimer with bridging phenylthiolate ligands. The X-ray structure of $[Tc(SPh)_2(N{=}NC_5H_4N)(HN{=}NC_5H_4N)]_2$, **267**, 165
Organohydrazine complexes
the reaction chemistry of a technetium(I) nitrosyl complex with potentially chelating organohydrazines: the X-ray crystal structure of $[TcCl_2(NO)(HN{=}NC_5H_4N)(PPh_3)]$, **279**, 206
Organoiron complexes
synthesis and crystal structures of complexes containing an Fe→Cd donor–acceptor bond formed by reaction of *trans*-$(CO)_3Fe(Ph_2Ppym)_2$ (Ph_2Ppym = 2-(diphenylphospino)pyrimidine) with CdX_2 (X = Cl, Br, I, SCN, ClO_4), **268**, 177
Organolead complexes
synthesis and structures of tin(II) and lead(II) 1-aza-allyls; the $[N(SiMe_3)C(Ph)C(SiMe_3)_2]^-$ ligand, **269**, 181
Organomercury(II) complexes
synthesis and NMR spectroscopy of dichlorobis(acetophenonethiosemicarbazone)mercury(II) formed from phenylmercury(II) chloride and acetophenonethiosemicarbazone: the first example of symmetrisation in organomercury(II)-thiosemicarbazone chemistry, **267**, 169
Organometallic complexes
organometallic Rh(I) and Rh(III) complexes with [2-(dimethylamino)ethyl]cyclopentadiene, **283**, 24
Organonitrile complexes
reactivities of the coordinated organonitriles in molybdenum(0) and tungsten(0) phosphine complexes: protonation of the nitrile carbon and cleavage of the C≡N triple bond, **280**, 163
Organotin complexes
synthesis and structures of tin(II) and lead(II) 1-aza-allyls; the $[N(SiMe_3)C(Ph)C(SiMe_3)_2]^-$ ligand, **269**, 181
Ornithine complexes
copper(II) complexes of (*R*,*S*)-α-hydroxymethylornithine and its N^{δ}-benzoyl derivative, **269**, 279
Osmacyclopentadiene complexes
steric limitations in associative substitution reactions of $Os_3(CO)_9(\mu\text{-}C_4Ph_4)$, **274**, 82
Osmium complexes
synthesis of nitridoosmium(VI) dithiolene complexes, **267**, 335
reversible β-hydride elimination of amine ligands forming stable *cis*-η^2-iminium hydride Os^{II} complexes, **270**, 337
nitrogen atom transfer and redox chemistry of terpyridyl phosphoraniminato complexes of osmium (IV), **270**, 511
crystalline dihydrogen complexes. Intramolecular and intermolecular interactions and dynamic behavior, **273**, 116
resonance Raman investigation of mixed-valence dinuclear and trinuclear complexes of Ru, Os and Re, **275–276**, 380
NMR characterization of four new isomers of $H(\mu\text{-}H)Os_3(CO)_{10}$(phosphine), **275–276**, 521
photoinduced processes in 4′-(9-anthryl)-2,2′:6′,2″-terpyridine, its protonated forms and Zn(II), Ru(II) and Os(II) complexes, **277**, 225
electrophilic abstractions on $Os(H)_2X(NO)L_2$: selective access to new unsaturated cationic Os(II) hydrides, **280**, 125
Oxaazadiamine complexes
lanthanide(III) complexes of two oxaazadiamine macrocyclic ligands derived from 2,6-diformylpyridine: the crystal structures of a reduced macrocyclic ligand and the corresponding diprotonated macrocycle, **282**, 42
Oxalate complexes
alternative reaction path for the photochemical reaction of tris(oxalato)cobaltate(III) in various aprotic solvents in the presence of oxygen, **271**, 29
Oxamato complexes
the preparation and X-ray structure of $[Ho(oxam)_3(H_2O)_3]_4 \cdot 2.75H_2O$, **267**, 299
tuning up the T_C in Mn(II)Cu(II) bimetallic planes and design of molecular-based magnets, **278**, 159
Oxamidate complexes
synthesis of polynuclear copper(II) complexes by controlling the conformation (*cis* or *trans*) of *N*,*N*′-bis(3-aminopropyl)oxamidate and the coordination mode of 2,2′-bipyrimidine, **281**, 95
Oxamido complexes
tuning up the T_C in Mn(II)Cu(II) bimetallic planes and design of molecular-based magnets, **278**, 159
Oxazoline complexes
synthesis and characterization of the new complex tris[(2-oxazolinyl)phenolato] manganese(III) and its use as a catalyst in oxidation reactions, **279**, 217
Oxidation
the kinetics and mechanism of the oxidation of 1,4-thioxane by potassium ferrate, **267**, 159
three dinuclear copper(II) carboxylates with the paddle-wheel cage structure as intermediates in copper(II) catalyzed oxidations of carboxylic acids. X-ray crystal structures of [tetrakis(diphenyl acetato-μ-*O*,*O*′)bis(acetonitrile-*N*)dicopper(II)] tetrakis(acetonitrile), [tetrakis(diphenylacetato-μ-*O*,*O*′)bis(acetone-*O*)dicopper(II)] and [tetrakis(1-phenyl-1-carboxylato-μ-*O*,*O*′-*cyclo*-pentane)-bis(ethanol-*O*)dicopper(II)] bis(ethanol), **267**, 183
kinetics and mechanism of the oxidation of ammineruthenium(II) complexes by bromine, **270**, 189
oxidation of amino esters to imine complexes with chiral high-valent oxoruthenium porphyrins, **273**, 142
the oxidation of nickel^{II}(1,4,8,11-tetraazacyclotetradecane) by hydroperoxyl radicals: a pulse radiolysis study, **273**, 266
relative reactivities of hydroxy compounds with short-lived manganese(V), **281**, 235
Oxidation catalysis
a new manganese-β-heptanitro-porphyrin with extreme redox potentials: spectral, electrochemical and catalytic properties, **272**, 4
sulfonated and acetamidosulfonylated tetraarylporphyrins as biomimetic oxidation catalysts under aqueous conditions, **272**, 228
Oxidation of arenes
selective C–H bond activation of arenes catalyzed by methylrhenium trioxide, **270**, 55
Oxidation of thiols
the kinetics and mechanism of oxidation of 3-mercaptopropionic acid, 2-mercaptoethanesulfonic acid and 2-mercaptobenzoic acid by potassium ferrate, **281**, 43
Oxidative addition
first- and second-order mechanisms for oxidative addition for bound methyl disulfide in the complex $W(CO)_3(phen)(MeSSMe)$, **270**, 488
unexpected bell-shaped effect of the ligand on the rate of the oxidative addition to palladium(0) complexes generated in situ from mixtures of $Pd(dba)_2$ and *para*-substituted triarylphosphines, **273**, 76

oxidative addition of methyl iodide and iodine to new binuclear rhodium(I) and iridium(I) compounds containing diaminoanthraquinonate-bridging ligands. Crystal structure of $[Rh_2(\mu\text{-}1,4\text{-}DA)(CO)_2(PPh_3)_2]$ (1,4-H_2DA = 1,4-diaminoanthraquinone), **274**, 15

Oxidative carbonylation

mechanisms of ≡C–H bond activation in oxidative carbonylations of alkynes catalyzed by palladium complexes, **280**, 202

Oxidative decarbonylation

oxidation of tricarbonylmolybdacarborane. 2. Oxidative decarbonylation with various sulfur donors, **283**, 145

Oxidative dimerization

homogeneous catalysis by tetra-t-butylphthalocyanine iron: intermediates in the oxidative dimerization of 2,6-di-t-butylphenol, **274**, 126

Oxidative reactions

a catalytic activity of glass beads, silica gels and anion-exchange resins modified with metal-porphines in oxidative reactions of ascorbic acid, **283**, 44

Oxide complexes

complexes of niobium and tantalum with chloride, sulfide and oxide ligands: synthesis and crystal structure of the (tetrachloro)bis-(η^2-disulfido)(μ-oxo)(μ-η^2:η^2-disulfido) bismetallates $[Cl_2(S_2)Nb(\mu\text{-}O)(\mu\text{-}S_2)Nb(S_2)Cl_2]^{2-}$ and $[Cl_2(S_2)Ta(\mu\text{-}O)(\mu\text{-}S_2)Ta(S_2)Cl_2]^{2-}$ as their tetraphenylphosphonium salts, **274**, 251

Lewis base adducts of halogenorhenium(VII) oxides: ^{17}O NMR spectroscopy, structural aspects and catalysis, **279**, 44

Oximato-bridged complexes

structure and magnetic properties of a linear oximato-bridged tetranuclear copper(II) complex, **268**, 263

Oxime complexes

synthesis and crystal structure of rubidium hydrogen oximate complexes with 18-crown-6: is it possible to reach a perfect fit of the rubidium atom inside the crown ether cavity?, **268**, 93

X-ray structures of Cu(poph)Cl_2, $[Cu(pxo)Cl_2]_2$ and $[Cu(pxo)(pxo\text{–}H)Cl]_2$ (poph = 2-pyridinecarboxaldehyde 1-oxide 2′-pyridinylhydrazone; pxo = 2-acetylpyridine 1-oxide oxime). One monomeric structure and two *N*-oxide-bridged dimeric structures, involving multidentate pyridine *N*-oxide ligands, **268**, 135

kinetics and mechanism of ester hydrolysis by metal complexes of 2,6-diacetylpyridine dioxime, **273**, 167

manifestation of redox duality of 2-propanone oxime: Pt(II)-assisted reduction versus Pt(IV)-mediated oxidation of $Me_2C{=}NOH$ species, **277**, 83

Oxoacetate complexes

chemistry of rare earth hypocoordinate oxocluster cations: formation of novel oxoacetate clusters, **281**, 201

Oxoalkoxide complexes

synthesis through an in situ esterification process and characterization of oxo isopropoxo titanium clusters, **279**, 144

Oxo-bridged complexes

synthesis of trichlorooxo-bis(2,6-dimethylphenylisocyanide) rhenium(V) and crystal structure of μ-oxo-bis[dichlorooxo-bis(2,6-dimethylphenylisocyanide)dirhenium(V)], **269**, 347

unexpected assembly of a (μ-oxo)(μ-formato)diiron(III) complex from an aerobic methanolic solution of Fe(III) and the TPA ligand, **273**, 393

binuclear oxo-bridged iron(III) complexes of chiral tetradentate ligands, **274**, 155

synthesis and characterization of a new unsymmetrical μ-oxo-bridged iron(III) complex with triphenylarsine oxide ligands, **277**, 127

structural and spectroscopic models of the manganese catalase active site. Isolation and structures of the asymmetric $[(H_2O)Mn^{III}(\mu\text{-}O)(\mu\text{-}O_2CR)_2Mn^{III}(L)]$ ($L = Cr_2O_7^{2-}$, CH_3OH) cores: analogs of a substrate-bound catalase active site intermediate, **281**, 25

a μ-oxo diiron complex of a new binucleating tetrapyridyl ligand, **282**, 243

preparation and properties of a series of (μ-oxo)di(μ-acetato) diruthenium(III) complexes, $[Ru_2(\mu\text{-}O)(\mu\text{-}CH_3COO)_2(pyridine)_4(L)_2]^{2+}$ and $[Ru_2(\mu\text{-}O)(\mu\text{-}CH_3COO)_2(AA)_2(L)_2]^{2+}$, where AA is 2,2′-bipyridine or 1,10-phenanthroline, and L is a monodentate ligand, **283**, 151

Oxo complexes

ligand exchange reactions of $ReCl_4(PPh_3)_2$ with tridentate diacidic ligands with the donor set O–N–O(S). Molecular structures of the resulting oxo-rhenium(V) complexes, **268**, 109

synthesis, characterization and study of the redox properties of rhenium(V) diolates, **271**, 65

the coordination mode of 3-hydroxypicolinic acid in oxorhenium(V) complexes. Crystal and molecular structures of $(n\text{-}Bu_4N)[ReOCl_3(HO\text{–}C_5H_3N\text{–}CO_2)]$ and $[ReOCl(HO\text{–}C_5H_3N\text{–}CO_2)_2]$, **269**, 235

oxidation of amino esters to imine complexes with chiral high-valent oxoruthenium porphyrins, **273**, 142

ligand exchange reactions of $ReCl_4(PPh_3)_2$, $ReOCl_3(PPh_3)_2$ and $NiCl_2(PPh_3)_2$ with acetylacetone thiobenzoylhydrazone. An unexpected template reaction at the ReO^{3+} core. Molecular structures of the resulting oxorhenium(V) and nickel complexes, **277**, 219

preparation and electrochemistry of six-coordinate monooxo molybdenum(VI) complexes containing bidentate catecholate and tridentate NOS-donor Schiff base ligands, **278**, 1

synthesis and crystal structure of a new isomer of the μ-oxo-bis[dichlorooxobis(pyridine)rhenium(V)] complex $\{OReCl_2py_2\}_2O$, **279**, 159

synthesis and characterization of two novel monooxovanadium(V) complexes with bidentate benzohydroxamate ligand, **282**, 149

mononuclear, dinuclear, and polynuclear dioxomolybdenum(VI)–tridentate Schiff base complexes: crystal structures and properties, **282**, 209

Oxodithiocarboxylic ester complexes

synthesis and coordination properties of amphiphilic 3-oxodithiocarboxylic esters, **269**, 83

Oxoferryl complexes

spectroscopic study of the compound ES and the oxoferryl compound II states of cytochrome *c* peroxidase: comparison with the compound II of horseradish peroxidase, **275–276**, 250

Oxo-pyrimidine

electrochemistry of 2-thio- and 2-oxo-pyrimidines in dimethyl formamide in the presence of dioxygen, **270**, 145

Oxorhenium complexes

the synthesis, characterization and X-ray crystal structures of the *fac-cis* and *mer-trans* isomers of $ReOCl_3(PMe_3)_2$, **267**, 123

synthesis and structural characterization of the [2,9-dimethyl-4,7-diaza-2,9-decanedithiolato]oxorhenium(V) complex, **268**, 169

synthesis and X-ray crystal structure of an oxorhenium(V) complex with a double Schiff base, **275–276**, 551

Oxo transfer

substituent effects of *cis*-dioxobis(dithiocarbamato)molybdenum(VI) on redox properties: redox potentials for one-electron reduction and second-order rate constants for oxygen atom transfer, **269**, 260

Oxovanadium complexes

oxovanadium(IV) complexes of imidazole-4-acetic, imidazole-4,5-dicarboxylic and pyrazole-3,5-dicarboxylic acids, **268**, 297

a bimetallic main group oxide cluster of the oxovanadium borate system: $(H_3NCH_2CH_2NH_3)_4[(VO)_{12}O_4\{B_8O_{17}(OH)_4\}_2\{Mn(H_2O)_2\}_2]\cdot H_2O$, **282**, 123

the formation of ternary complexes between VO(maltolate)$_2$ and small bioligands, **283**, 202

Oxovanadium compounds

hydrothermal synthesis and structural characterization of the three-dimensional framework solids $(NH_4)[V_2(OH)(PO_4)_2(H_2O)]\cdot H_2O$ and $Rb_6[(Mo_9V_3O_6)(PO_4)_{10}(H_2PO_4)_3(OH)_9]\cdot 8.5H_2O$, **275–276**, 122

Oxygen activation

integrated photocatalysts for hydrocarbon oxidation: polyoxotungstates/iron porphyrins systems in the reductive activation of molecular oxygen, **272**, 197
Oxygen binding
the dynamics of formation of the O_2–Co^{II} bond in the cobalt(II) cyclidene complexes, **270**, 151
Oxygen exchange
reversible formation of high-valent-iron-oxo porphyrin intermediates in heme-based catalysis: revisiting the kinetic model for horseradish peroxidase, **275–276**, 98

Packing
effect of substitution on packing in *trans*-di(*Z*-chlorobenzoato-*O*)-di(1,3-diaminopropane-*N*,*N'*)nickel(II) (*Z* = 2, 3 or 4) complexes, **267**, 249
Palladacycles
enantioselective cleavage of activated amino acid esters promoted by chiral palladacycles, **280**, 57
Palladium-catalyzed
synthesis of porphyrin–spacer–quinone compounds via metal-mediated cross-coupling: new systems for probing the relative magnitudes of axial and equatorial electronic coupling at the porphyrin macrocycle in thermal and photoactivated electron transfer reactions, **275–276**, 427
Palladium complexes
some oxidative addition reactions with a diphosphadibora[1.1.0]bicyclobutane: formation of cage compounds and of an unusual trinuclear coordination compound of palladium, **269**, 101
palladium(II)-catalyzed alternating copolymerization of alkenes with sulfur dioxide, **270**, 8
palladium-catalyzed seven-membered N,O-heterocycle ring formation, **270**, 123
an unusual cyanide-bridged extended A-frame complex: the molecular structure of $[\{Pd_2Br_2(\mu\text{-}PPh_2)(\mu\text{-}dppm)\}_2(\mu\text{-}CN)_2]$, **271**, 236
cyclopalladated complexes of Schiff bases of homoveratrylamine and tryptamine. Synthesis and CO insertion, **272**, 18
Pd(gly-L-his-L-lys)Cl: solution structure and ternary complex formation with mono- and tetranucleotides, **273**, 31
comparative DFT and HF-MP2 study on structure, relative stabilities and rotational barriers of end-on (η^1) and side-on (η^2) $(PH_3)_2Pd$ HPPH model complexes, **273**, 72
unexpected bell-shaped effect of the ligand on the rate of the oxidative addition to palladium(0) complexes generated in situ from mixtures of $Pd(dba)_2$ and *para*-substituted triarylphosphines, **273**, 76
nickel(II), palladium(II), platinum(II) and platinum(IV) complexes of *cis*-1,2-bis(diphenylphosphino)ethene and nitrogen-containing heterocycles or sulfur ligands, **274**, 143
novel palladium(II) allyl complexes with nitrogen–sulfur donor bidentate ligands. Mechanism of allyl amination of $[Pd(\eta^3\text{-allyl})(N\text{-}SR)]ClO_4$ (allyl = C_3H_5; N-SR = C_5H_4N-2-CH_2SR, R = C_6H_5, C_2H_5) in the presence of activated olefins. X-ray structure determination and fluxional behavior, **275–276**, 385
palladium(II) and platinum(II) complexes containing new phosphinoxy chiral P,N-ligands, **277**, 119
crystal structure and fluxional behavior of novel Pd(II) allyl complexes containing a potentially tridentate S,N,S ligand: an example of selective *syn–anti* isomerism, **277**, 247
complexes of a new bidentate chelating pyridyl/sulfonamide ligand with copper(II), cobalt(II) and palladium(II): crystal structures and spectroscopic properties, **278**, 178
synthesis and metal complexes of a chelating ligand containing thiomorpholine and pyridine: 2,6-bis(thiomorpholinomethyl)pyridine, **279**, 221
structures, dynamic behaviour in solution and cross-coupling reactions of the α-palladiated (η^6-alkylarene)tricarbonylchromium complexes containing a palladium–chromium bond, **280**, 71
the *trans* influence of F, Cl, Br and I ligands in a series of square-planar Pd(II) complexes: relative affinities of halide anions for the metal centre in *trans*-$[(Ph_3P)_2Pd(Ph)X]^+$, **280**, 87
synthesis and characterization of homologous nickel(II) and palladium(II) complexes with biphosphine monoxide ligands $Ph_2P(CH_2)_nP(O)Ph_2$ (n = 1–3) and $pTol_2P(CH_2)P(O)pTol_2$, **280**, 150
mechanisms of ≡C–H bond activation in oxidative carbonylations of alkynes catalyzed by palladium complexes, **280**, 202
chiroptical properties of heterotrimetallic 'wing-tip' butterflies; synthesis and crystal structure of $(\mu\text{-}H)Ru_2Fe_2PdC(CO)_{12}(\eta^3\text{-}\beta\text{-}C_{10}H_{15})$, **280**, 275
synthesis and crystal structure of dinuclear palladium(I) dppm complex with 4-mercaptopyridine in the thione mode. $[Pd_2(\mu\text{-}dppm)_2(HpyS)_2](ClO_4)_2 \cdot CH_3OH$, **281**, 116
reactivity of $[NBu_4]_2[\{(C_6X_5)_2M(\mu\text{-}OH)\}_2]$ (X = F or Cl; M = Pd or Pt) towards heterocyclic thiones: crystal structure of $[NBu_4]_2[\{(C_6F_5)_2Pd(\mu\text{-}\eta^2\text{-}LL')\}_2]$ (LL' = methylimidazole-2-thiolate), **281**, 165
complexes of the bis(trimethylsilyl)-benzamidinato ligand '*siam*': synthesis and X-ray structures of $(siam)_2M$, $(siam)(siamH)MX$ (M = Ni, Pd), $(siam)_2MnI$ and $(siam)ReO_3$, and their reactivity towards CO_2, **281**, 181
electrocrystallization and characterization of phosphonium-based radical anion salts $[MePh_3P][Pd(dmit)_2]_3$ and $[Ph_4P][Pd(dmit)_2]_3$ (dmit = 1,3-dithiole-2-thione-4,5-dithiolate). Comparison with related Pd and Ni isologues, **282**, 90
the synthesis and characterization of Ni, Pd and Pt maleonitriledithiolate complexes: X-ray crystal structures of the isomorphous Ni, Pd and Pt $(Ph_2PCH_2CH_2PPh_2)M$(maleonitriledithiolate) congeners, **282**, 155
Palladium(II) complexes
palladium(II) complexes of *N*-sulfonyl-asparagine and glutamine. Evidence for metal coordination of the deprotonated amide nitrogen of the side-chain, **273**, 397
Palladium–platinum/C catalysts
selective oxidation of glucose on bismuth-promoted Pd–Pt/C catalysts prepared from $NOct_4Cl$-stabilized Pd–Pt colloids, **270**, 95
Paramagnetic complexes
the synthesis and characterization of $[Tc(SPh)_2(N{=}NC_5H_4N)(HN{=}NC_5H_4N)]_2$, a paramagnetic dimer with bridging phenylthiolate ligands. The X-ray structure of $[Tc(SPh)_2(N{=}NC_5H_4N)(HN{=}NC_5H_4N)]_2$, **267**, 165
formation and reactivity of paramagnetic organometallic nickel complexes of 21-oxa- and 21-selenaporphyrins — ^1H NMR and EPR investigations, **273**, 184
selective versus non-selective T_1 experiments to determine metal–nucleus distances in paramagnetic metalloproteins, **275–276**, 373
Paramagnetic NMR
NMR spectra of dinuclear manganese and iron compounds containing 1,4,7-triazacyclononane and 1,4,7-trimethyl-1,4,7-triazacyclononane, **268**, 43
probing the metal site in *Rhus vernicifera* stellacyanin by Ni(II) substitution and paramagnetic NMR spectroscopy, **273**, 367
structure and reactivity of reconstituted myoglobins: interaction between protein and polar side chain of chemically modified hemin, **275–276**, 159
spectroscopic investigations on the water-soluble fragment of the Rieske [2Fe–2S] protein from *Paracoccus denitrificans*, **275–276**, 493
Paramagnetic relaxation
a ^{31}P and ^1H NMR relaxometric study of the interaction between adenosine triphosphate (ATP) and paramagnetic ions (Gd^{3+} and Mn^{2+}), **273**, 92
Paramagnetic shifts
calculation of NMR paramagnetic shift temperature dependence in actinide methyl borohydrides, **272**, 43

the orientation of the iron axial ligands in the low-potential cytochrome c_{549} from *Synechocystis sp.* PCC 6803 studied by NMR, **273**, 196

P450cam
roles of valine-98 and glutamic acid-72 of putidaredoxin in the electron-transfer complexes with NADH-putidaredoxin reductase and P450cam, **272**, 80

P-donor substituents
steric limitations in associative substitution reactions of $Os_3(CO)_9(\mu\text{-}C_4Ph_4)$, **274**, 82

Pentacyanoferrate complexes
4- and 3-hydroxypyridine complexes of pentacyanoferrate(II, III), **267**, 81

Pentalene complexes
bis(η^3-allyl)(η^8-pentalene)zirconium: preparation, structure and structural dynamics, **270**, 555

Pentamethylcyclopentadienyl complexes
trimethylsilyl iodide as a halide exchange reagent in inorganic chemistry: synthesis and structure of $Cp^*_2ThI_2$ ($Cp^* = \eta^5\text{-}C_5Me_5$), **274**, 232
pentamethylcyclopentadienyl ruthenium(II) complexes containing chiral diphosphines: synthesis, characterisation and electrochemical behaviour. X-ray structure of $(\eta^5\text{-}C_5Me_5)Ru\{(S,S)\text{-}Ph_2PCH(CH_3)CH(CH_3)PPh_2\}Cl$, **282**, 163

Pentanedioic acid complexes
synthesis and catalytic activity of manganese(II) complexes of pentanedioic acid; X-ray crystal structure of $[Mn(phen)_2(H_2O)_2]\text{-}[Mn(O_2C(CH_2)_3CO_2)(phen)_2H_2O](O_2C(CH_2)_3CO_2)\cdot 12H_2O$ (phen = 1,10-phenanthroline), **277**, 257

Peptide complexes
miniaturized hemoproteins: design, synthesis and characterization of mimochrome II, **275–276**, 301
metal binding properties of three Cys_2X_2 (X = His, Asp) metallothionein-related peptides, **278**, 10
hemoprotein models based on a covalent helix–heme–helix sandwich 4. Discrimination of paramagnetic Fe(III)–Mimochrome I Δ and Λ isomers by NMR spectroscopy, **278**, 76
how non-bonding amino acid side-chains may enormously increase the stability of a Cu(II)–peptide complex, **283**, 1

Peptide synthesis
salt-induced peptide formation from amino acids in the presence of clays and related catalysts, **272**, 89

Perfluoroalkoxy-substituted complexes
preparation and some properties of perfluoroalkoxy-substituted phthalocyanine complexes of iron(III), nickel(II) and zinc(II), **278**, 253

Perfluoroarene complexes
a ferrocene–perfluoroarene molecular complex, **270**, 559

Perhalogenated cyclopentadiene complexes
coordination chemistry of perhalogenated cyclopentadienes. XXII. Reaction of tetrabromodiazocyclopentadiene with some halide bridged Rh(I), Ru(II) and Ru(III) complexes. Molecular structure of $[C_5Br_5]Rh(COD)$, a complex with an η^1,η^4-coordinated cyclopentadienyl ligand, **269**, 111

Perhaloporphyrin complexes
catalysis of aerobic olefin oxidation by a ruthenium perhaloporphyrin complex, **270**, 433

Peroxidase
reversible formation of high-valent-iron-oxo porphyrin intermediates in heme-based catalysis: revisiting the kinetic model for horseradish peroxidase, **275–276**, 98

Peroxo complexes
alternative reaction path for the photochemical reaction of tris(oxalato)cobaltate(III) in various aprotic solvents in the presence of oxygen, **271**, 29

Peroxovanadate complexes
solvent effects on ^{51}V NMR chemical shifts: characterization of vanadate and peroxovanadate complexes in mixed water/acetonitrile solvent, **283**, 37

Peroxovanadium complexes
solvation, preferential solvation and complexation by the solvent of peroxovanadium complexes studied by ^{51}V NMR spectroscopy. Correlations with the oxidative reactivity, **272**, 62

Peroxydisulfate
properties and kinetics of dihydroxy- and diaminoanthraquinone ruthenium bipyridyl dimers, **268**, 31

pH dependence
thermodynamics of reduction of *Chromatium vinosum* high-potential iron–sulfur protein and its histidine depleted H42Q mutant, **275–276**, 230

Phenanthroline complexes
structural study and electron exchange kinetics of the bis(2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline)copper(II)/(I) couple in acetonitrile, **277**, 31
syntheses and X-ray crystal structures of *cis*-$[Mn(bipy)_2Cl_2]\cdot 2H_2O\cdot EtOH$ and *cis*-$[Mn(phen)_2Cl_2]$ (bipy = 2,2′-bipyridine; phen = 1,10-phenanthroline); catalysts for the disproportionation of hydrogen peroxide, **279**, 24
polymeric low-valence platinum–phenanthroline complexes as precursors of platinum colloids, **280**, 339
synthesis and characterization of mono-, bi- and tetra-ruthenium(II)-phenanthroline complexes with dipyrido(2,3-*a*:2′,3′-*h*)phenazine (dpop), **281**, 1

Phenylalanine complexes
a ternary complex of carbonic anhydrase: X-ray crystallographic structure of the adduct of human carbonic anhydrase II with the activator phenylalanine and the inhibitor azide, **275–276**, 295

Phosphanide complexes
syntheses and structure of dimeric yttrium tris[bis(trimethylsilyl)-phosphanide]; X-ray structures of $[(thf)LiCl\cdot(thf)_2LiP^tBu_2]_2$ and $Y[CH(SiMe_3)_2]_3$, **269**, 91

Phosphate-bridged complexes
complexes of the terdentate N-donor ligand 6-(2-aminophenyl)-2,2′-bipyridine (L): crystal structures of mononuclear $[ZnL_2][PF_6]_2$ and tetranuclear $[\{CuL(MeCN)\}_4(\mu^4\text{-}PO_4)][PF_6]_5$ containing an unusual μ^4-bridging phosphate ion, **267**, 239

Phosphate complexes
hydrothermal synthesis and structural characterization of a three-dimensional organically-templated vanadium phosphate $[H_3NCH_2CH_2NH_3][(VO)_3(H_2O)_2(PO_4)_2(HPO_4)]$, **268**, 257

Phosphate compounds
hydrothermal synthesis and structural characterization of the three-dimensional framework solids $(NH_4)[V_2(OH)(PO_4)_2(H_2O)]\cdot H_2O$ and $Rb_6[(Mo_9V_3O_6)(PO_4)_{10}(H_2PO_4)_3(OH)_9]\cdot 8.5H_2O$, **275–276**, 122

Phosphate esters
hydrolysis of phosphate esters catalyzed by copper(II)-triamine complexes. The effect of triamine ligands on the reactivity of the copper(II) catalysts, **283**, 51

Phosphides
phosphide complexes of divalent europium: synthesis and structural characterization of $Eu[PPh_2]_2(N\text{-}MeIm)_4$, **267**, 309

Phosphido-bridged complexes
synthesis of a dinuclear, phosphido-bridged complex of rhenium, X-ray crystal structure of $Re_2(\mu\text{-}H)[\mu\text{-}(t\text{-}Bu)PC_6H_4P(H)t\text{-}Bu](CO)_7$, **277**, 253

Phosphine complexes
an infrequent geometry for an element of the first transition series: structural characterization of a distorted eight-coordinate compound $TiCl_4[C_2H_4(PMe_2)_2]_2$, **267**, 173
NMR characterization of four new isomers of $H(\mu\text{-}H)Os_3(CO)_{10}$(phosphine), **275–276**, 521

the dicarbonylation of diiodomethane to malonate esters catalysed by triethylphosphine complexes of rhodium, **280**, 99
syntheses and crystal structures of geometrical isomeric pairs: *trans*- and *cis*-$(PPh_4)[Co(acac)_2(CN)_2]$ and *trans*- and *cis*-$[Co(acac)_2(PMe_3 \text{ or } PEt_3)_2]PF_6$ (acac = pentane-2,4-dionate), **281**, 77
kinetics of the photolysis of bis(triethylphosphine)dichloroplatinum(II) in chloroform, **282**, 217

Phosphine halide complexes
trimethylphosphine halide complexes of titanium, zirconium and hafnium in oxidation states IV, III and II. Crystal structure of $MI_4(PMe_3)_n$ (M = Zr, n = 3; M = Hf, n = 2.5), $M_2X_6(PMe_3)_4$ (M = Hf, Ti, X = I; M = Zr, X = Br) and $(\mu\text{-}\eta^6,\eta^6\text{-}C_6H_6)\text{-}Hf_2I_4(PMe_3)_4 \cdot C_6H_6$, **271**, 180

Phosphine oxide complexes
synthesis, structure and properties of lanthanide dithionates and their triphenylphosphine oxide complexes, **274**, 177

Phosphino-amine complexes
synthesis and reactivity of Ru(II) complexes containing the phosphino-amine $Ph_2PCH_2CH_2NMe_2$, **268**, 69
substitution reactions of $CpRu(PPh_3)_2Cl$ with PPh_2H and dppa, **269**, 157
synthesis, characterization and reactivity of bis(diphenylphosphino)-amine nickel complexes, **271**, 112
copper(I) and copper(II) complexes of polydentate phosphinoamines, **275–276**, 401
phosphinoamino complexes of copper(I) and silver(I): a combined NMR and IR spectroscopy, and electrospray ionization and fast atom bombardment mass spectrometry analysis, **282**, 119

Phosphite complexes
synthesis and reactivity of bis(ethylene)-phosphite complexes of molybdenum(0), **272**, 125
synthesis and characterization of arene, mono- and dihydrido-arene, monohydrido-cyclopentadienyl, and phosphite complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$, **281**, 153

Phosphonic acid complexes
lanthanide complexes with 1-hydroxyethane-1,1-diphosphonic acid: solvent organization and coordination geometry in crystalline and amorphous solids, **269**, 211

Phosphonium complexes
electrocrystallization and characterization of phosphonium-based radical anion salts $[MePh_3P][Pd(dmit)_2]_3$ and $[Ph_3P][Pd(dmit)_2]_3$ (dmit = 1,3-dithiole-2-thione-4,5-dithiolate). Comparison with related Pd and Ni isologues, **282**, 90

Phosphopolyoxometallate complexes
investigations into the engineering of inorganic/organic solids: hydrothermal synthesis and crystal structures of aromatic ammonium sodium hydrogen phosphomolybdates(VI), **282**, 10

Phosphoraniminato complexes
nitrogen atom transfer and redox chemistry of terpyridyl phosphoraniminato complexes of osmium (IV), **270**, 511

Photocatalysis
integrated photocatalysts for hydrocarbon oxidation: polyoxotungstates/iron porphyrins systems in the reductive activation of molecular oxygen, **272**, 197

Photochemistry
alternative reaction path for the photochemical reaction of tris(oxalato)cobaltate(III) in various aprotic solvents in the presence of oxygen, **271**, 29
the use of photogenerated intermediates in the study of cluster build-up reactions: the generation of $Ru_6C(CO)_{17}$ from $Ru_3(CO)_{12}$, **278**, 250

Photolysis
intramolecularly alkylated Cosalen complexes: thermolysis and photolysis of coenzyme B_{12} models with a cobalt-to-ligand carbon bridge, **269**, 203
synthesis and characterization of copper(I) complexes with triphenylphosphine and heterocyclic thione ligands: the crystal structure of (thiazolidine-2-thione)(bis-triphenylphosphine)copper(I) chloride, **271**, 243
kinetics of the photolysis of bis(triethylphosphine)dichloroplatinum(II) in chloroform, **282**, 217

Photolysis of water
preparation and characterization of the catalytically active heteropolyoxotungstate anion $[Co^{3+}ZnW_{11}O_{40}H_2]^{7-}$, **278**, 91

Photophysics
remarkable substituent effects on the photophysics of Pt(4′-X-trpy)Cl^+ systems (trpy = 2,2′; 6′,2″-terpyridine), **273**, 346
photophysical properties of porphyrins, phthalocyanines, and benzochlorins, **279**, 226
intercalation into calf thymus DNA of 2-(4-arylphenyl)imidazo-[4,5-f]-[1,10]phenanthroline (aryl = -OMe, $-NMe_2$ or $-NO_2$) in its bipyridyl ruthenium(II) complex, **283**, 17

Photoreduction
photoreduction of methyl-viologen by zeolite-Y stabilized silver clusters, **269**, 253

Photosensitizers
protonation of the ground states of ruthenium(II) photosensitizers, **279**, 200

pH potentiometry
oxovanadium(IV) complexes of imidazole-4-acetic, imidazole-4,5-dicarboxylic and pyrazole-3,5-dicarboxylic acids, **268**, 297

Phthalocyaninato complexes
a new synthetic route to unsymmetrical bis(phthalocyaninato)-europium(III) complexes, **268**, 141

Phthalocyanine complexes
cyanide binding to phthalocyaninatoiron(II) in dimethyl sulfoxide in the presence of carbon monoxide: kinetic and equilibrium study, **267**, 109
heteroleptic triple-decker (phthalocyaninato)(porphyrinato)-europium(III) complexes: synthesis and electrochemical study, **268**, 49
homogeneous catalysis by tetra-t-butylphthalocyanine iron: intermediates in the oxidative dimerization of 2,6-di-t-butylphenol, **274**, 126
adducts of nitric oxide with cobaltous tetraphenylporphyrin and phthalocyanines: potential nitric oxide sorbents, **277**, 1
fixation of carbon dioxide with 1,2-epoxypropane catalysed by tetra-t-butylphthalocyaninatoaluminium(III) acetate, tetra-t-butylphthalocyaninatoaluminium(III) trimethylsilanolate and tetra-t-butylphthalocyaninatoaluminium(III) triphenylsilanolate, **278**, 223
preparation and some properties of perfluoroalkoxy-substituted phthalocyanine complexes of iron(III), nickel(II) and zinc(II), **278**, 253
photophysical properties of porphyrins, phthalocyanines, and benzochlorins, **279**, 226
mixed valence, tetradisordered structure of iodine-doped bismuth(III) diphthalocyanine, **281**, 195
catalytic oxygenation of olefin with dioxygen and tetra-t-butylphthalocyanine complexes in the presence of sodium borohydride, **283**, 275

Picrate complexes
different picrate coordination modes along the lanthanide series: synthesis and molecular structure of two isomorphous compounds $[Ln_2(pic)_4(TDTD)_3(H_2O)_2]$ $(pic)_2(H_2O)_2$ (Ln = Tm, Lu; pic = picrate; TDTD = *trans*-1,4-dithiane-*S*,*S*′-dioxide), **274**, 122
synthesis and crystal structure of a new polynuclear lanthanum picrate complex with *trans*-1,4-dithiane-1,4-dioxide, **274**, 236

Piperazine complexes
boat versus chair conformation in *N*-methyl- and *N*,*N*′-dimethylpiperazine platinum(II) complexes studied by X-ray analysis. A rare example of metal chelate piperazine: *cis*-$[PtCl_2(Me_2ppz)]$. Part I, **275–276**, 410

synthesis and molecular structure of *N*-methylpiperazinium(+1) and *N,N'*-dimethylpiperazinium(+2) platinum(II) and platinum(IV) complexes. Part II, **275–276**, 419

a linear open-chain piperazine–pyridine ligand and its *meso*-helical Co complex, **277**, 55

pK_a values

pK_a values of aqua ligands of platinum(II) anticancer complexes: [^{1}H, ^{15}N] and ^{195}Pt NMR studies of *cis*- and *trans*-[$PtCl_2(NH_3)$-(cyclohexylamine)], **273**, 8

Plastocyanin

homology predicted structure and comparison with the secondary structure from NMR data for plastocyanin from the cyanobacterium *Synechocystis* sp. PCC 6803, **275–276**, 73

spectroscopic characterization and kinetic studies of a novel plastocyanin from the green alga *Ulva pertusa*, **283**, 184

Platinum-195 NMR

the synthesis and characterization of platinum monodithiolene complexes containing 1,3-dithiole-2-oxo-4,5-dithiolate ($dmid^{2-}$) and 1,3-dithiole-2-thione-4,5-dithiolate ($dmit^{2-}$), **282**, 200

Platinum colloids

polymeric low-valence platinum–phenanthroline complexes as precursors of platinum colloids, **280**, 339

Platinum complexes

synthesis of two maleimide derivatives of *cis*-configurated platinum(II) complexes for the preparation of chemoimmunoconjugates, **267**, 133

thermodynamics of the formation of *cis*-bis(dialkylsulfide)dimethylplatinum(II) complexes from bis(μ-dialkylsulfide)bisdimethylplatinum(II) and free dialkylsulfide, **268**, 311

monofunctional *trans*-$Pt^{II}(NH_3)_2$ modification of pyrimidine-rich deoxyoligonucleotides: direct platination and use of a protective group, **269**, 135

fast water exchange on a Pt(II) center: variable temperature and pressure ^{17}O NMR study at 14.1 tesla of the ionic arylplatinum species $[Pt\{C_6H_3(CH_2NMe_2)_2\text{-}2,6\}(OH_2)]^+(OSO_2CF_3)$, **269**, 322

hydride complexes of platinum(II) containing unsymmetrical di(phosphine) ligands: synthesis, characterization and evidence for pairwise addition of hydrogen based on parahydrogen induced polarization, **270**, 363

intramolecular and intermolecular C–H activation at a cationic Pt^{II} center, **270**, 467

an ab initio MO-LCAO investigation of the electronic structure of organic hydroperoxides and platinum(II) hydroperoxo complexes: a contribution to the knowledge of the mechanism of olefin epoxidation, **270**, 479

substitution of inosine for chloride in $[Pt_2(hdta)Cl_2]^{2-}$ ($hdta^{4-}$ = 1,6-hexanediamine-*N,N,N',N'*-tetraacetate), **271**, 124

cytotoxic anionic tribromo platinum(II) complexes containing benzothiazole and benzoxazole donors: synthesis, characterization, and structure–activity correlation, **271**, 137

comparative time-resolved IR and UV spectroscopic study of monophosphine and diphosphine platinum(II) azido complexes, **271**, 160

synthesis and antitumor activity of platinum(II) complexes of aminocyclodextrin, **272**, 162

pK_a values of aqua ligands of platinum(II) anticancer complexes: [^{1}H, ^{15}N] and ^{195}Pt NMR studies of *cis*- and *trans*-[$PtCl_2(NH_3)$-(cyclohexylamine)], **273**, 8

oxidative addition of 2,3,4-benzotrithiepin 1-oxide to platinum(0) compounds: a convenient synthetic route to platinum(II) thiosulfinato complexes, **273**, 106

the role of $O_2^{\bullet-}$ radicals in DNA–metal complexes, **273**, 201

remarkable substituent effects on the photophysics of Pt(4′-X-trpy)Cl^+ systems (trpy = 2,2′; 6′,2″-terpyridine), **273**, 346

nickel(II), palladium(II), platinum(II) and platinum(IV) complexes of *cis*-1,2-bis(diphenylphosphino)ethene and nitrogen-containing heterocycles or sulfur ligands, **274**, 143

synthesis of binuclear platinum(II) organometallic complexes as potential cross-linking reagents, **275–276**, 141

interaction of organometallic cationic complex ions containing terpyridine ligands with nucleic acids: an investigation on aggregative phenomena, **275–276**, 242

boat versus chair conformation in *N*-methyl- and *N,N'*-dimethylpiperazine platinum(II) complexes studied by X-ray analysis. A rare example of metal chelate piperazine: *cis*-[$PtCl_2(Me_2ppz)$]. Part I, **275–276**, 410

synthesis and molecular structure of *N*-methylpiperazinium(+1) and *N,N'*-dimethylpiperazinium(+2) platinum(II) and platinum(IV) complexes. Part II, **275–276**, 419

influence of *cis*- and *trans*-diamminedichloroplatinum(II) binding on the helix-coil transition of DNAs with different GC content, **275–276**, 510

reactions of platinum(II)–alkyne and –alkynyl complexes with the hydroxy compounds $HO(CH_2)_2X$ (X = Br, I, OH), **275–276**, 515

toward a novel metal based chemotherapy against tropical diseases. 4. Synthesis and characterization of new metal-clotrimazole complexes and evaluation of their activity against *Trypanosoma cruzi*, **275–276**, 528

manifestation of redox duality of 2-propanone oxime: Pt(II)-assisted reduction versus Pt(IV)-mediated oxidation of $Me_2C{=}NOH$ species, **277**, 83

palladium(II) and platinum(II) complexes containing new phosphinoxy chiral P,N-ligands, **277**, 119

trans influence of triphenylstibine. Crystal and molecular structures of *cis*-[$PtCl_2(SbPh_3)_2$] and *trans*-[$PtI_2(SbPh_3)_2$], **277**, 236

the redox kinetics of platinum(II)/(IV) complexes, **278**, 32

electronic absorption and MCD spectra for the $Pt(PPh_3)(AuPPh_3)_6^{2+}$ and $Pt(H)(PPh_3)(AuPPh_3)_7^{2+}$ cluster complexes, **278**, 113

intercalator-linked cisplatin: synthesis and antitumor activity of *cis*-dichloroplatinum(II) complexes connected to acridine and phenylquinolines by one methylene chain, **279**, 51

a phosphane analog of nedaplatin: preparation and structural characterization of *cis*-bis(trimethylphosphane)(glycolato-*O,O'*)-platinum(II), **279**, 105

synthesis and structural properties of (diamine)platinum(II) complexes of itaconic acid, **279**, 116

tris- and tetrakis(1-methylcytosine) complexes of Pt(II): syntheses and X-ray structures of $[Pt(1\text{-}MeC\text{-}N3)_3Cl]^+$ and $[Pt(1\text{-}MeC\text{-}N3)_4]^{2+}$ compounds, **279**, 152

preparation and properties of $C_8H_4S_8$-platinum(II) complexes and electrical conductivities of their oxidized species and X-ray crystal structure of $C_8H_4S_8(CH_2CH_2CN)_2$ as a pro-ligand compound, **279**, 165

protonation of cyclopropene complexes of platinum(0) and the reduction of cyclopropene by nitrogenases, **280**, 193

molecular dynamics of *N,N'*-di-*p*-fluorophenyltriazenido complex of platinum(II), **280**, 238

polymeric low-valence platinum–phenanthroline complexes as precursors of platinum colloids, **280**, 339

organometallic polymers containing coordinatively saturated platinum(II) complexes as pendants, **281**, 141

reactivity of $[NBu_4]_2[\{(C_6X_5)_2M(\mu\text{-}OH)\}_2]$ (X = F or Cl; M = Pd or Pt) towards heterocyclic thiones: crystal structure of $[NBu_4]_2[\{(C_6F_5)_2Pd(\mu\text{-}\eta^2\text{-}LL')\}_2]$ (LL′ = methylimidazole-2-thiolate), **281**, 165

kinetics and mechanism for reduction of *trans*-dichloro-tetracyanoplatinate(IV) by tetraammineplatinum(II) and bis(ethylenediamine)platinum(II), **282**, 55

the synthesis and characterization of Ni, Pd and Pt maleonitriledithiolate complexes: X-ray crystal structures of the isomorphous Ni, Pd and Pt ($Ph_2PCH_2CH_2PPh_2$)M(maleonitriledithiolate) congeners, **282**, 155

the synthesis and characterization of platinum monodithiolene complexes containing 1,3-dithiole-2-oxo-4,5-dithiolate ($dmid^{2-}$) and 1,3-dithiole-2-thione-4,5-dithiolate ($dmit^{2-}$), **282**, 200
kinetics of the photolysis of bis(triethylphosphine)dichloroplatinum(II) in chloroform, **282**, 217
synthesis and crystal structure of a dinuclear platinum(II) bis(*N*,*N*-diethylamino)carbeniumdithiocarboxylate complex $Pt_2\{\mu\text{-}(Et_2N)_2C_2S_2\}_2Cl_4\cdot 0.5CH_2Cl_2$: structural study of the coordinated inner salt, **283**, 256

Platinum(II) complexes
the preparation and crystal and molecular structure of the complex *cis*-$PtCl_2$(razoxane), **268**, 313
five-coordinate platinum(II) alkyne complexes: synthesis, ab initio calculations and crystal and molecular structure of $[PtI_2(Me_2phen)(\eta^2\text{-}PhC{\equiv}CPh)]\cdot CHCl_3$, **275–276**, 500

Platinum(IV) complexes
can Pt(IV)–amine complexes act as 'prodrugs'?, **283**, 251

P,N-donor ligand complexes
some chlorocarbonylruthenium(II) complexes of P,N-donor ligands: crystal structures of $[RuCl(CO)\{PPh_2CH_2C(Bu^t){=}NNH_2\}_2]Cl$ and *fac*,*cis*-$[RuCl_2(CO)\{PPh_2CH_2C(Bu^t){=}N\text{–}N{=}C(Bu^t)\text{-}CH_2PPh_2\}]$, **279**, 95

P,N-ligand complexes
the synthesis and characterization of a technetium nitrosyl complex with *cis*-{2-pyridyl,diphenylphosphine} coligands. The X-ray crystal structure of $[TcCl_2(NO)(pyPPh_2\text{-}P,N)(pyPPh_2\text{-}P)]$, **267**, 319
synthesis and physical studies of copper(I) complexes with the ambidentate ligand (1-piperidyl-2-diphenylphosphino)ethane. X-ray structural characterization of a copper(I) dimer, **271**, 217

Polyamine complexes
reinforced piperazine rings containing polyamines: metal complex equilibria and structural studies, **268**, 63
anionopentaaminecobalt(III) complexes with polyamine ligands. 29. Relative substitution lability in structurally related chloropentaaminecobalt(III) complexes, **274**, 42
gold(III) compounds as potential antitumor agents: cytotoxicity and DNA binding properties of some selected polyamine–gold(III) complexes, **281**, 90

Polyamine ligand complexes
synthesis and crystal structure of the *trans*-dichloro(*N*,*N*′-bis-[2(*S*)-pyrrolidylmethyl]-ethane-1,2-diamine)cobalt(III) tetrachlorocobaltate(II) complex, **267**, 127

Poly(amino carboxylate) complexes
a tripod gadolinium(III) poly(amino carboxylate) relevant to magnetic resonance imaging: structural and dynamical ^{17}O NMR and ^{1}H NMRD studies, **275–276**, 15

Polyazamacrocycle complexes
interaction of mono- and triphosphate anions with a protonated polyaza macrocycle and its Cu(II) complexes, **270**, 207
synthesis and properties of a new tetraaza macrocyclic ligand containing one N-acetic acid pendant arm and its copper(II) and nickel(II) complexes, **274**, 24

Polydentate ligand complexes
square-pyramidal copper(II) complexes of linear tetradentate edda-type ligands forming six-membered rings. Molecular structures of $[Cu(1,3\text{-}pdda)(H_2O)]$ and $[Cu(eddp)(H_2O)]\cdot 3.5H_2O$, **268**, 221
syntheses and properties of bivalent metal complexes with *N*-(2-hydroxyethyl)ethylenediamine-*N*,*N*′,*N*′-triacetic acid and the crystal structure of *N*-(2-hydroxyethyl)ethylenediamine-*N*,*N*′,*N*′-triacetato-copper(II) monohydrate, **271**, 13
synthesis and characterisation of mononuclear and binuclear iron(II) complexes of pentadentate and bis(pentadentate) ligands derived from 1,4,7-triazacyclononane, **279**, 192

Polyfluorooxo anions
preparation and characterization of two novel heteropolyfluorooxotungstate anions $[NaCoW_{11}O_{43}FH_{12}]^{6-}$ and $[NaNiW_{11}O_{43}FH_{12}]^{6-}$, **282**, 136

Polyfunctional tridentate ligand complexes
a new class of biomimetically relevant 'Scorpionate' ligands. III. The bis(pyrazolyl)methane(phen-2′-ol)s: synthesis and structural characterization of mono and dinuclear copper(II) complexes, **273**, 14

Polyhedra
chemical applications of topology and group theory. 30. Distortions from idealized symmetrical coordination polyhedra by lifting molecular orbital degeneracies, **270**, 68

Polyhydride complexes
hydrogen site exchange in $CpIr(PR_3)H_3^+$ complexes, **272**, 95

Polyhydroxy acid complexes
coordination chemistry of polyhydroxy acids: role of the hydroxy groups, **275–276**, 130

Polyimidazole ligands
sterically hindered iron(II) complex of a new tripodal polyimidazole ligand: structure and reactivity toward superoxide, **275–276**, 234

Polyiodide complexes
synthesis, X-ray and spectroscopic characterization of $[SnI_2(mbit)_2]\text{-}(I_3)_2\cdot 2/3I_2$ obtained through the one-step reaction of $mbit\cdot 2I_2$ with tin metal powder (mbit = 1,1′-bis(3-methyl-4-imidazoline-2-thione)methane), **267**, 115

Polymer-anchored complexes
the hydrogenation and hydroformylation of alkenes as catalyzed by polymer-anchored rhodium trichloride under water gas shift reaction conditions, **270**, 345

Polymer complexes
lanthanide metal complexes of *N*,*N*′-ethylenebis(pyridin-2-one); from chelates to metallacyclic frameworks, **272**, 131
weak intermolecular antiferromagnetic exchange in $\{[Ru_2(O_2CCH_3)_4(L)]X\}_n$ polymers (L = 4,4′-dipyridine and 1,4-diazabicyclooctane, X = PF_6^- or BPh_4^-), **279**, 121
structure and magnetic properties of a novel chloro-bridged polymeric cadmium(II) complex with pyridyl-substituted nitronyl nitroxide, **279**, 233
organometallic polymers containing coordinatively saturated platinum(II) complexes as pendants, **281**, 141
spectroscopic characterization of coaxially and π–π stacked binuclear copper(II) complexes of ω,ω′-bis(pyridine-2-carboxamido)-alkanes in acidic solution, **283**, 61

Polymeric compounds
organic–inorganic composite materials: structural archetypes of linear polymeric chlorocadmates(II), **272**, 252

3,2′-Polymethylene-6-(2″-pyridyl)-2-phenylpyridines
synthesis and properties of 3,2′-polymethylene-6-(2″-pyridyl)-2-phenylpyridines and their cycloplatinated complexes, **267**, 265

Polynuclear complexes
resonance Raman investigation of mixed-valence dinuclear and trinuclear complexes of Ru, Os and Re, **275–276**, 380
synthesis of polynuclear copper(II) complexes by controlling the conformation (*cis* or *trans*) of *N*,*N*′-bis(3-aminopropyl)oxamidate and the coordination mode of 2,2′-bipyrimidine, **281**, 95
synthesis, crystal structure and magnetic properties of the one-dimensional chain compound *catena*-poly[aquabis(4,4′-bis-1,2,4-triazole-*N*1)(thiocyanato-*N*)-manganese-μ-[4,4′-bis-1,2,4-triazole-*N*1,*N*1′]thiocyanate], **282**, 257

Polyoxometalate compounds
a new form of decavanadate: hydrothermal synthesis and structural characterization of $Ba_{5.33}(H_3O)_{0.33}[V_{10}O_{30}]$, **277**, 263

Polyoxomolybdate compounds
stepwise substituting magnetic centres in a giant cluster host system, **271**, 9

Polyoxotungstates
integrated photocatalysts for hydrocarbon oxidation: polyoxotungstates/iron porphyrins systems in the reductive activation of molecular oxygen, **272**, 197

study of polyoxotungstates with the Keggin structure by cyclic voltammetry, **281**, 225

Polypyrazolylborato complexes
syntheses, properties and structural characteristics of several new (polypyrazolylborato)samarium(III) complexes: agostic interaction of the BH_2 proton with samarium, **277**, 8

Polypyridyl complexes
crystal and molecular structure of $[Ru(bpy)_2(2,3\text{-dpp})]Cl_2 \cdot 3H_2O \cdot CH_3CN$ and ^1H and ^{99}Ru NMR spectra of $[Ru(bpy)_2(2,n\text{-dpp})][PF_6]_2$ (bpy = 2,2′-bipyridine, dpp = bis(2-pyridyl)-pyrazine, $n = 3$ or 5), **275–276**, 320
intercalation into calf thymus DNA of 2-(4-arylphenyl)imidazo-[4,5-f]-[1,10]phenanthroline (aryl = -OMe, $-NMe_2$ or $-NO_2$) in its bipyridyl ruthenium(II) complex, **283**, 17

Polypyridyl ligands
syntheses and structure of dinuclear cobalt(II) complexes of the polypyridine ligand 6,6′-bis[bis(2-pyridylmethyl)aminomethyl]-2,2′-bipyridine, **268**, 159

Porphyrinate complexes
synthesis, UV–visible and electrochemical studies of lipophilic and hydrophilic lanthanide(III) bis(porphyrinates), **275–276**, 182

Porphyrin complexes
preparation and characterization of water-soluble viologen-linked zinc porphyrin and bisviologen-linked zinc porphyrin, **267**, 257
heteroleptic triple-decker (phthalocyaninato)(porphyrinato)-europium(III) complexes: synthesis and electrochemical study, **268**, 49
EXAFS studies on Mn(III) tetraphenylporphyrin halides in Nafion, **268**, 231
rate constants and activation parameters for organo–cobalt porphyrin bond homolysis from NMR relaxation times, **270**, 197
synthesis of rhodium porphyrin aryls via intermolecular arene carbon–hydrogen bond activation, **270**, 551
ligand photolysis and recombination of $(dimethylsulfoxide)_2Fe(II)$-porphyrin complexes, **271**, 119
a new manganese-β-heptanitro-porphyrin with extreme redox potentials: spectral, electrochemical and catalytic properties, **272**, 4
integrated photocatalysts for hydrocarbon oxidation: polyoxotungstates/iron porphyrins systems in the reductive activation of molecular oxygen, **272**, 197
new non-ionic water-soluble porphyrins: evaluation of manganese(III) polyhydroxylamide porphyrins as MRI contrast agents, **275–276**, 106
synthesis of porphyrin–spacer–quinone compounds via metal-mediated cross-coupling: new systems for probing the relative magnitudes of axial and equatorial electronic coupling at the porphyrin macrocycle in thermal and photoactivated electron transfer reactions, **275–276**, 427
adducts of nitric oxide with cobaltous tetraphenylporphyrin and phthalocyanines: potential nitric oxide sorbents, **277**, 1
hemoprotein models based on a covalent helix–heme–helix sandwich 4. Discrimination of paramagnetic Fe(III)–Mimochrome I Δ and Λ isomers by NMR spectroscopy, **278**, 76
photophysical properties of porphyrins, phthalocyanines, and benzochlorins, **279**, 226
structure and properties of tetraphenylporphinate iron(III) complexes with an intramolecular NH···S benzenethiolate or NH···O phenolate hydrogen bond, **283**, 91
molecular structure of the methoxy-iron(III) derivative of 5,15-(*o*,*o*′(2-methyl-2′-hydroxy-3,3′-diamidobiphenyl)-diphenylporphyrin and CO binding properties of iron(II)–pyridine complexes of 2,2′-substituted biphenyl strapped porphyrins, **283**, 98

Porphyrin protein supramolecules
the synthesis and spectroscopic characterization of a chiral zinc(II) porphyrin dimer containing an amino acid bridge: model complexes for the induced circular dichroism of porphyrin protein supramolecules, **272**, 295

Potassium complexes
cation-π interactions between potassium ions and aromatic rings. Crystal structures of three potassium complexes of calix[6]arene, **281**, 36

Prismane complexes
a synthetic iron-containing prismane complex: peculiar electron-delocalization and spin-coupling properties deduced from Mössbauer and magnetization studies, **283**, 111

Prochiral ketones
the influence of the catalyst preparation protocol and silane structure on the rate and enantioselectivity of ansa-titanocene catalysed hydrosilation of prochiral ketones, **270**, 330

Protein complexes
the immobilisation of proteins in carbon nanotubes, **272**, 261
zinc as a structural and folding element of proteins which interact with DNA, **275–276**, 481
a μ-oxo diiron complex of a new binucleating tetrapyridyl ligand, **282**, 243
folding properties of iron–sulfur proteins, **283**, 12

Protonation
reinforced piperazine rings containing polyamines: metal complex equilibria and structural studies, **268**, 63
protonation of the ground states of ruthenium(II) photosensitizers, **279**, 200

Proton NMR
homology predicted structure and comparison with the secondary structure from NMR data for plastocyanin from the cyanobacterium *Synechocystis* sp. PCC 6803, **275–276**, 73

Pteridine complexes
synthesis and structural studies on three-dimensionally hydrogen-bonded metal perchlorate complexes with 1,3-dimethyllumazine and 1,3,6,7-tetra-methyllumazine. Crystal structure of diaqua-bis-(1,3,6,7-tetramethyl-pteridine-2,4(1*H*,3*H*)-dione-O^4,N^5)-copper(II) perchlorate dihydrate, **268**, 77

Putidaredoxin
roles of valine-98 and glutamic acid-72 of putidaredoxin in the electron-transfer complexes with NADH-putidaredoxin reductase and P450cam, **272**, 80

Pyrazine-bridged complexes
an asymmetrically-coordinated 2,3-dimethylpyrazine bridging ligand in the (II,II) binuclear ion, $\{[Ru^{II}(hedta)]_2(2,3\text{-}Me_2pz)\}^{2-}$ ($hedta^{3-}$ = *N*-hydroxyethylethylenediaminetriacetate), **268**, 279

Pyrazinecarboxamide complexes
ligand behaviour of a bis-pyrazinecarboxamide copper(II) complex towards a copper(II) β-diketonate. X-ray crystal structures of $[Cu(bpzpn)(H_2O)] \cdot 3H_2O$ and $[Cu(bpzpn)Cu(hfacac)_2]_2$ (H_2bpzpn = *N*,*N*′-bis(2′-pyrazinecarboxamide)-1,3-propane, Hhfacac = hexafluoroacetylacetone), **271**, 210

Pyrazole complexes
copper(II) complexes of pyrazol-3-yl amines. Synthesis, crystal and solution structure, **269**, 117
electrophilic substitution of metal-coordinated pyrazole: nitration, sulfonation and bromination of $[Co(NH_3)_5(pyzH)]^{3+}$ (pyzH = pyrazole), **277**, 89

Pyrazolylborate complexes
synthesis and characterization of new hydrotri(3,5-dimethyl-1-pyrazolyl)borate vanadium halfsandwich complexes, **267**, 19
complexes of thallium(I) and lead(II) with the potentially tetradentate ligand bis[3-(2-pyridyl)-pyrazolyl]dihydroborate, **267**, 323
syntheses and structural characterization of chromium(III) complexes containing hydrotris(1-pyrazolyl)borate, **269**, 302
syntheses of ruthenium hydridotris(1-pyrazoryl)borate complexes having sulfur-donor coligands, **273**, 238
synthesis and characterisation of rhodium(I) complexes containing the dihydrobis(pyrazolyl)borate ligands $[H_2B(pz')_2]^-$. The X-ray structures of $Bp^*Rh(C_2H_4)_2$ and $BpRh(PPh_3)_2$ ($Bp = H_2B(pz)_2$; $Bp^* = H_2B(3,5\text{-}Me_2\text{-}pz)_2$, **273**, 244

coordination chemistry of the sterically hindered N_3-donor hydrotris(3,5-diphenylpyrazol-1-yl)borate toward silver(I)triorganophosphino compounds. Synthesis, structural and spectroscopic characterization, **282**, 1

Pyridazine complexes
stereochemically rigid mono and bis pyridazine complexes of $[Ru^{II}(hedta)]^-$ ($hedta^{3-}$ = *N*-hydroxyethylethylenediaminetriacetate), **279**, 85

Pyridine
effect of pyridine on the alkaline isomerization of cytochrome *c*, **275–276**, 58

Pyridine ligand complexes
synthesis of new copper(II) complexes of polydentate ligands containing pyridyl pendant arms. Crystal structure of [*N*,*N*,*N*′,*N*′-tetrakis(2-pyrid-2-yl-ethyl)1,2-ethylenediaminecopper(II)]-bis(hexafluorophosphate)·1.5 acetone, **278**, 237

Pyridine *N*-oxide complexes
X-ray structures of $Cu(poph)Cl_2$, $[Cu(pxo)Cl_2]_2$ and $[Cu(pxo)(pxo–H)Cl]_2$ (poph = 2-pyridinecarboxaldehyde 1-oxide 2′-pyridinylhydrazone; pxo = 2-acetylpyridine 1-oxide oxime). One monomeric structure and two *N*-oxide-bridged dimeric structures, involving multidentate pyridine *N*-oxide ligands, **268**, 135

Pyridine-thionate complexes
complexes of copper(I), silver(I), nickel(II) and cadmium(II) with 6-tert-butyldimethylsilylpyridine-2-thionate (6-Bu^{t}SidmepyS$^-$): crystal structure of $[Cu_2(6\text{-Bu}^t\text{SidmepyS})_2dppm_2]$, **271**, 1

Pyridyl-sulfonamide ligand complexes
complexes of a new bidentate chelating pyridyl/sulfonamide ligand with copper(II), cobalt(II) and palladium(II): crystal structures and spectroscopic properties, **278**, 178

Pyridyl(thienylacetylene) complexes
coordination compounds of the rigid rod ligand 1-(2-pyridyl)-2-(2-thienyl)ethyne (pythio). Single crystal X-ray structures of $[Cu(pythio)_2(NO_3)_2]$ and $[Zn(pythio)_2Cl_2]$, **267**, 193

Pyridyl-thioether ligand complexes
the 1,8-bis(2′-pyridyl)-3,6-dithiaoctane complex of nickel(II): X-ray crystal structure and borohydride adduct formation, **278**, 217

Pyrimidine complexes
stereochemically-rigid and fluxional pyrimidines within the bis-$[Ru(hedta)(pym)_2]^-$ complex, **268**, 287

Pyruvic acid and pyridoxal thiosemicarbazones
synthetic, spectroscopic and X-ray crystallographic studies on copper(II) complexes with pyruvic acid and pyridoxal thiosemicarbazones, **269**, 297

Quadruply bonded complexes
quadruply bonded complexes containing η^2- or η^3-tridentate phosphine ligands; syntheses, structures and $^{31}P\{^1H\}$ NMR spectra of $Mo_2Cl_4(\eta^3\text{-etp})(L)$ and $Mo_2Cl_4(\eta^2\text{-etp})_2$ (L = CH_3OH, PMe_3 and PEt_3; etp = $Ph_2PCH_2CH_2P(Ph)CH_2CH_2PPh_2$), **274**, 7

Quinolinethiolato complexes
thermal and photochemical reactions of bis(8-quinolinolato)-copper(II), bis(8-quinolinethiolato)copper(II) and (8-quinolinolato,8-quinolinethiolato)copper(II) with chloroalkanes, **273**, 403

Racemic complexes
first solid state synthesis of the racemic compound and enantiomer of dinuclear Λ,Λ-di-μ-hydroxo-bis[bis((1*S*,2*S*)-1,2-*trans*-cyclohexanediamine)chromium(III)] and Δ,Δ-di-μ-hydroxo-bis-[bis((1*R*,2*R*)-1,2-*trans*-cyclohexanediamine)chromium(III)] bromides, **274**, 210

Racemization
stereochemistry of six-coordinated silicon complexes. V. Racemization and geometrical isomerization of tris(4-isopropyltropolonato)-silicon(IV), **271**, 24

isomerization and racemization of (acetylacetonato)bis(4,4′-dimethyl-2,2′-bipyridine 1,1′-dioxide)chromium(III), (acetylacetonato)bis(4,4′-dimethoxy-2,2′-bipyridine 1,1′-dioxide)-chromium(III) and (acetylacetonato)bis(4,4′-diethoxy-2,2′-bipyridine 1,1′-dioxide)chromium(III) complexes, **281**, 221

Radical complexes
infrared spectroelectrochemical and electrochemical kinetics studies of the reaction of nickel cluster radicals $[Ni_3(\mu_2\text{-dppm})_3(\mu_3\text{-L})(\mu_3\text{-I})]^+$(L = CNR, R = CH_3, i-C_3H_7, C_6H_{11}, $CH_2C_6H_5$, t-C_4H_9, 2,6-$Me_2C_6H_3$; L = CO) with carbon dioxide, **270**, 111

transition metal complexes with nitronyl nitroxides. Part 1. Crystal structure and properties of square-planar *trans* radical–copper(II)–radical complexes, **279**, 69

Radiopharmaceuticals
complexation of gallium(III) by 2-hydroxynicotinic acid and 3-hydroxypicolinic acid in aqueous solution, **273**, 47

Rare earth complexes
chemistry of rare earth hypocoordinate oxocluster cations: formation of novel oxoacetate clusters, **281**, 201

Rare-gas atom complexes
exohedral and endohedral complexation energies of complexes X@C_{70} between higher fullerene and rare-gas atoms (X = He, Ne, Ar, Kr, Xe), **272**, 33

Re_6S_8 core complexes
synthesis of the first sulfido-bridged octahedral rhenium(III) aqua ion $[Re_6S_8(H_2O)_6]$, **271**, 228

Reaction intermediate
homogeneous catalysis by tetra-t-butylphthalocyanine iron: intermediates in the oxidative dimerization of 2,6-di-t-butylphenol, **274**, 126

Reconstituted myoglobin
structure and reactivity of reconstituted myoglobins: interaction between protein and polar side chain of chemically modified hemin, **275–276**, 159

Redox chemistry
titanium–carbon functionalities on an oxo surface defined by a calix[4]arene moiety and its redox chemistry, **270**, 298

Redox reactions
the redox kinetics of platinum(II)/(IV) complexes, **278**, 32

Redox thermodynamics
effects of solvent on the redox properties of cytochrome *c*: cyclic voltammetry and ^1H NMR experiments in mixed water–dimethylsulfoxide solutions, **272**, 168

Reduction potential
thermodynamics of reduction of *Chromatium vinosum* high-potential iron–sulfur protein and its histidine depleted H42Q mutant, **275–276**, 230

the effect of surface charge on the reduction potential and heme propionate ionization of a monoheme cytochrome: spectroscopic and potentiometric studies of trifluoroacetylated cytochrome c_{551} from *Pseudomonas aeruginosa*, **275–276**, 256

Reductive elimination
a mechanistic study of thermal C–H reductive elimination from a six-coordinate d^6 iridium complex, **270**, 537

Resonance Raman spectroscopy
Co–C force constants from resonance Raman spectra of alkylcobalamins: insensitivity to dimethylbenzylimidazole coordination, **270**, 392

ground and electronically excited states of $Cr(CO)_4$(bipyridine): energy factored force field analysis of CO stretching vibrations and resonance Raman study, **278**, 83

Respiration
spectroscopic investigations on the water-soluble fragment of the Rieske [2Fe–2S] protein from *Paracoccus denitrificans*, **275–276**, 493

Restrained molecular dynamics calculations
Boc-Glu-Thr-Ile-His-OMe/Zn^{2+} in terms of the zinc-binding sites of proteases, **283**, 243

Rhenium cluster
synthesis of the first sulfido-bridged octahedral rhenium(III) aqua ion $[Re_6S_8(H_2O)_6]$, **271**, 228

Rhenium complexes
ligand exchange reactions of $ReCl_4(PPh_3)_2$ with tridentate diacidic ligands with the donor set O–N–O(S). Molecular structures of the resulting oxo-rhenium(V) complexes, **268**, 109
the synthesis and characterization of *trans*-$ReCl_2(dppe)_2$ and α-$Re_2Cl_4(dppe)_2$, **268**, 151
additions of carbon nucleophiles to acyclic imine complexes of the chiral rhenium Lewis acid $[(\eta^5\text{-}C_5H_5)Re(NO)(PPh_3)]^+$: controlling factors in 1,3-asymmetric induction and syntheses of non-racemic organic amines, **269**, 167
the coordination mode of 3-hydroxypicolinic acid in oxorhenium(V) complexes. Crystal and molecular structures of (n-Bu_4N)-$[ReOCl_3(HO\text{–}C_5H_3N\text{–}CO_2)]$ and $[ReOCl(HO\text{–}C_5H_3N\text{–}CO_2)_2]$, **269**, 235
synthesis of trichlorooxo-bis(2,6-dimethylphenylisocyanide) rhenium(V) and crystal structure of μ-oxo-bis[dichlorooxo-bis(2,6-dimethylphenylisocyanide)dirhenium(V)], **269**, 347
selective C–H bond activation of arenes catalyzed by methylrhenium trioxide, **270**, 55
thermal and photochemical substitution reactions of $CpRe(PPh_3)_2H_2$ and $CpRe(PPh_3)H_4$. Catalytic insertion of ethylene into the C–H bond of benzene, **270**, 77
the synthesis and characterization of α-$Re_2Cl_4(depe)_2$ and α-$W_2Cl_4(depe)_2$, **271**, 49
synthesis, characterization and study of the redox properties of rhenium(V) diolates, **271**, 65
an electron paramagnetic resonance study of tetra-*n*-butylammonium-tetrabromo-nitridorhenate(VI), $[(n\text{-}C_4H_9)_4N][ReNBr_4]$, **271**, 199
rhenium(VII) and (V) imido and oxo complexes containing 1,4,7-triazacyclononane: synthesis, electrochemistry and crystal structures, **272**, 176
technetium and rhenium carbonyl complexes with 1,4-benzodiazepine derivatives: X-ray crystal structure, **272**, 267
electrochemical and photophysical properties of two-component coordination compounds containing a metallocyclam and an $Re^I(bipy)(CO)_3Cl$ subunit, **275–276**, 117
resonance Raman investigation of mixed-valence dinuclear and trinuclear complexes of Ru, Os and Re, **275–276**, 380
ligand exchange reactions of $ReCl_4(PPh_3)_2$, $ReOCl_3(PPh_3)_2$ and $NiCl_2(PPh_3)_2$ with acetylacetone thiobenzoylhydrazone. An unexpected template reaction at the ReO^{3+} core. Molecular structures of the resulting oxorhenium(V) and nickel complexes, **277**, 219
synthesis of a dinuclear, phosphido-bridged complex of rhenium, X-ray crystal structure of $Re_2(\mu\text{-}H)[\mu\text{-}(t\text{-}Bu)PC_6H_4P(H)t\text{-}Bu](CO)_7$, **277**, 253
the reactivity of the $[Re{\equiv}O]^{3+}$ core toward aromatic 1,2-diamines: the formation of rhenium(V)-imido complexes, **278**, 96
NMR evidence for formation of new alcohol rhenium complexes as intermediates in ionic hydrogenations of carbonyl groups with systems composed of $ReH_2(NO)(CO)(PR_3)_2$ (R = Pr^i, CH_3, OPr^i) and CF_3COOH, **278**, 122
Lewis base adducts of halogenorhenium(VII) oxides: ^{17}O NMR spectroscopy, structural aspects and catalysis, **279**, 44
synthesis and crystal structure of a new isomer of the μ-oxo-bis[dichlorooxobis(pyridine)rhenium(V)] complex $\{ORecl_2py_2\}_2O$, **279**, 159
intermolecular Re–H···H–X hydrogen bonding (X = N, C) involving $ReH_5(PPh_3)_3$, **280**, 26
in situ IR and NMR study of the interactions between proton donors and the Re(I) hydride complex $[\{MeC(CH_2PPh_2)_3\}Re(CO)_2H]$. ReH···H bonding and proton-transfer pathways, **280**, 302
syntheses and properties of organocyanamide, cyanoguanidine and dinitrogen complexes of rhenium. Crystal structure of *mer*-$[ReCl_2(NCNEt_2)(PMePh_2)_3]$, **280**, 308
interaction of [ReO(SNS)(S)] and [^{99m}TcO(SNS)(S)] mixed ligand complexes with glutathione: isolation and characterization of the product, **281**, 148
complexes of the bis(trimethylsilyl)-benzamidinato ligand '*siam*': synthesis and X-ray structures of (*siam*)$_2$M, (*siam*)(*siam*H)MX (M = Ni, Pd), (*siam*)$_2$MnI and (*siam*)ReO_3, and their reactivity towards CO_2, **281**, 181

Rhenium(V) complexes
the synthesis, characterization and X-ray crystal structures of the *faccis* and *mer-trans* isomers of $Re(NEt)Cl_3(PMe_3)_2$, **278**, 245

Rheumatoid arthritis
the synthesis, NMR and X-ray analysis of tris(1-ethyl-2-isopropylimidazol-5-yl)phosphine gold(I)-2,3,4,6-tetra-O-acetyl-1-thio-β-D-glucopyranoside, a bifunctional auranofin analogue, **274**, 137

Rhodium complexes
syntheses, structures and properties of hydrosulfido-bridged diiridium and dirhodium complexes, $[Cp^*MCl(\mu_2\text{-}SH)_2MCp^*Cl]$ and $[Cp^*M(\mu_2\text{-}SH)_3MCp^*]^+$ (M = Ir, Rh; $Cp^* = \eta^5\text{-}C_5Me_5$), **267**, 73
NMR investigations of di-rhodium-tetraacetate complexes with cyclic phosphites, **267**, 293
dissociative kinetics of 'lightly bonded ligands' *L* from the clusters $Rh_6(CO)_{15}L$ (*L* = DMSO, NCMe, cyclooctene, THF and EtOH), **268**, 189
coordination chemistry of perhalogenated cyclopentadienes. XXII. Reaction of tetrabromodiazocyclopentadiene with some halide bridged Rh(I), Ru(II) and Ru(III) complexes. Molecular structure of $[C_5Br_5]Rh(COD)$, a complex with an η^1,η^4-coordinated cyclopentadienyl ligand, **269**, 111
structures of $HRh(PMe_2Ph)_4$ and $HRh(PPh_2Me)_4$, **269**, 162
rhodium aryloxide complexes containing terdentate nitrogen ligands, C–O bond formation, hydrogen bonding with phenol and oxidative addition reactions. Molecular structure of an Rh(III)–acetyl complex, **270**, 34
on the mechanism of silylformylation catalyzed by Rh–Co mixed metal complexes, **270**, 279
the hydrogenation and hydroformylation of alkenes as catalyzed by polymer-anchored rhodium trichloride under water gas shift reaction conditions, **270**, 345
the synthesis of the tripodal phosphine $CH_3C\{CH_2P(m\text{-}CF_3C_6H_4)_2\}_3$ and its rhodium coordination chemistry, **270**, 499
synthesis of rhodium porphyrin aryls via intermolecular arene carbon–hydrogen bond activation, **270**, 551
rhodium(III) ditertiary stibine complexes, **271**, 203
synthesis and characterization of heterobimetallic complexes containing C–S cleaved thiophenes, **272**, 55
transition metal complexes with sulfur ligands. Part CXXV. Synthesis and characterization of hydrido and chloro complexes with rhodium sulfur cores, **272**, 211
rhodium(III) analogues of antitumour-active ruthenium(III) compounds: the crystal structure of [ImH][*trans*-$RhCl_4(Im)_2$] (Im = imidazole), **273**, 62
synthesis and characterisation of rhodium(I) complexes containing the dihydrobis(pyrazolyl)borate ligands $[H_2B(pz')_2]^-$. The X-ray structures of $Bp^*Rh(C_2H_4)_2$ and $BpRh(PPh_3)_2$ (Bp = $H_2B(pz)_2$; Bp^* = $H_2B(3,5\text{-}Me_2\text{-}pz)_2$, **273**, 244
oxidative addition of methyl iodide and iodine to new binuclear rhodium(I) and iridium(I) compounds containing diaminoanthraquinonate-bridging ligands. Crystal structure of $[Rh_2(\mu\text{-}1,4\text{-}DA)(CO)_2(PPh_3)_2]$ (1,4-H_2DA = 1,4-diaminoanthraquinone), **274**, 15
study of the relative π-acceptor ability of unsaturated ligands $:C(=C)_nPh_2$ (n = 0, 1, 2 and 4) based on cyclic voltammetry of complexes *trans*-$[RhCl(L)(P^iPr_3)_2]$ (L = $:C(=C)_nPh_2$, C_2H_4 and CO), **275–276**, 435
toward a novel metal based chemotherapy against tropical diseases. 4. Synthesis and characterization of new metal-clotrimazole com-

plexes and evaluation of their activity against *Trypanosoma cruzi*, **275–276**, 528
synthesis, characterization and kinetics of the metallation of the intermediate compound $[Rh_2(\mu\text{-}O_2CCH_3)_3(O_2CCH_3)\{\eta^2\text{-}P(2\text{-}(CH_3O)C_6H_4)Ph_2\}]\cdot H_2O$ to produce compound $[Rh_2(\mu\text{-}O_2CCH_3)_3\text{-}\{\mu\text{-}(CC_5H_4)P(2\text{-}(CH_3O)C_6H_4)Ph\}]\cdot(H_2O)$ from Rh(II) tetraacetate, **278**, 61
solvent-induced reaction of rhodium(III) halide with N-heterocyclic ligands. Crystal structure of *trans*-dichlorotetrakis(*N*-methylimidazole)rhodium(III) chloride dihydrate, **278**, 209
synthesis, immobilization and catalytic activity of some silylated cyclopentadienyl rhodium(I) complexes, **280**, 21
dialkyl-μ-ethylidene-μ-methylene-bis(pentamethylcyclopentadienyl)-dirhodium complexes: synthesis, spectra and decomposition reactions, **280**, 62
the dicarbonylation of diiodomethane to malonate esters catalysed by triethylphosphine complexes of rhodium, **280**, 99
preparation, structure and properties of triphenylphosphine rhodium aryloxides, **280**, 172
synthesis, solution behavior and molecular structure of 16-electron *closo*-2-(Ph_3P)-1-N,2-$[\mu\text{-}(\eta^2\text{-}CH_2CH{=}CH_2)]$-1-N-$(\sigma\text{-}CH_2CH{=}CH_2)$-2,1-$RhCB_{10}H_{10}$, the first η^2-olefinic monocarbon metallacarborane, **280**, 233
pentacoordinated cationic rhodium(I) and iridium(I) organocomplexes with α-diimine ligands derived from glyoxal: crystal structure of [Rh(norbornadiene)(glyoxalbis(4-dimethylamino-anil))-$(PPh_3)]BF_4\cdot 1/2CH_2Cl_2$, **282**, 230

Rhodium(III) complexes
arm-off reactions of the tripodal polyphosphane $MeC(CH_2PPh_2)_3$ in octahedral rhodium(III) complexes of heterocyclic thioamides, **267**, 39

Rhodopseudomonas palustris
isolation and characterization of cytochrome c_2 from *Rhodopseudomonas palustris*, **269**, 125

Ribbon-sheet complexes
the structural characterization of the novel ribbon sheet, $[Cu_2Cl_2(pyz)]_n$ (pyz = pyrazine), **267**, 143

ROESY spectra
ROESY spectra of paramagnetic model heme complexes: unambiguous differentiation of cross-peaks due to chemical exchange and the nuclear Overhauser effect in the same molecule, **272**, 204

Rubidium complexes
synthesis and crystal structure of rubidium hydrogen oximate complexes with 18-crown-6: is it possible to reach a perfect fit of the rubidium atom inside the crown ether cavity?, **268**, 93

Rubredoxin
the use of ^{113}Cd NMR chemical shifts as a structural probe in tetrathiolate metalloproteins, **273**, 279
probing the iron environment in desulforedoxin. EXAFS of oxidized and reduced states, **273**, 409

Ruthenium clusters
reactivity studies on cationic non-hydridic triruthenium carbonyl clusters. Reactions of $[Ru_3(\mu_3\text{-ampy})(CO)_{10}][BF_4]$ with hydrogen, triethylsilane and triphenylstannane, **268**, 131

Ruthenium complexes
effects of ligand topology on the properties of dinuclear ruthenium complexes of bis-semiquinone bridging ligands, **267**, 1
synthesis and properties of the unsymmetrical binuclear complexes: $Na[(CN)_5Fe(dtdp)Ru(NH_3)_5]6H_2O$ and $K[(CN)_5Ru(dtdp)Ru(NH_3)_5]6H_2O$, dtdp = 4,4′-dithiodipyridine, **267**, 93
imidazole release from the antitumor-active ruthenium complex imidazolium *trans*-tetrachlorobis(imidazole)ruthenate(III) by biologically occurring nucleophiles, **267**, 137
interaction between the NO^+ ligand and *trans* π-donor ligands in chloro(2-methyl-8-quinolinolato)nitrosyl(8-quinolinolato)-ruthenium(II) complexes, **267**, 147
site-selective PMe_3 addition to the donor–acceptor complex $Ru_2(CO)_6[(Z)\text{-}Ph_2PCH{=}CHPPh_2]$. X-ray diffraction structure of $Ru_2(CO)_5(PMe_3)[(Z)\text{-}Ph_2PCH{=}CHPPh_2]$, **267**, 223
preparation and molecular structure of (*N*-benzyl-1,2-didehydroglycinato) bis(1,10-phenanthroline)ruthenium(II) chloride: $[Ru(N\text{-}Bngly\text{-}H_2)(phen)_2]Cl$, **267**, 329
properties and kinetics of dihydroxy- and diaminoanthraquinone ruthenium bipyridyl dimers, **268**, 31
synthesis and reactivity of Ru(II) complexes containing the phosphino-amine $Ph_2PCH_2CH_2NMe_2$, **268**, 69
an asymmetrically-coordinated 2,3-dimethylpyrazine bridging ligand in the (II,II) binuclear ion, $\{[Ru^{II}(hedta)]_2(2,3\text{-}Me_2pz)\}^{2-}$ ($hedta^{3-}$ = *N*-hydroxyethylethylenediaminetriacetate), **268**, 279
stereochemically-rigid and fluxional pyrimidines within the bis-$[Ru(hedta)(pym)_2]^-$ complex, **268**, 287
transition metal complexes with sulfur ligands. Part CXXVII. Azido, halido and nitrido ruthenium complexes with sulfur-rich coordination spheres, **269**, 63
coordination chemistry of perhalogenated cyclopentadienes. XXII. Reaction of tetrabromodiazocyclopentadiene with some halide bridged Rh(I), Ru(II) and Ru(III) complexes. Molecular structure of $[C_5Br_5]Rh(COD)$, a complex with an η^1,η^4-coordinated cyclopentadienyl ligand, **269**, 111
substitution reactions of $CpRu(PPh_3)_2Cl$ with PPh_2H and dppa, **269**, 157
redox behavior of a binuclear ruthenium complex having a di-μ-nitrosyl ligand, $[\{Ru(acac)_2\}_2(\mu\text{-}NO)_2]$ (acac = acetylacetonato), **269**, 342
ligand connected metal clusters. The molecular structures and solid state packing of $\{Ru_3(CO)_{11}\}_2$(bis(diphenylphosphino)ethane) and $\{Ru_3(CO)_{11}\}_2$(1,4-bis(diphenylphosphino)benzene), **270**, 28
second sphere coordination behavior of aquo and amine ligands bound to a η^6-benzeneruthenium(II) cation, **270**, 87
triply-bridged diruthenium(II) 1,4-bis(diphenylphosphino)butane (dppb) and (*R*)-2,2′-bis(diphenylphosphino)-1,1′-binaphthyl (binap) complexes, including structural characterisation of $[(dppb)ClRu(\mu\text{-}D_2O)(\mu\text{-}Cl)_2RuCl(dppb)]$, $[(\eta^2\text{-}H_2)\text{-}(dppb)Ru(\mu\text{-}Cl)_3RuCl(dppb)]$ and the $[(dppb)ClRu(\mu\text{-}Cl)_3RuCl(dppb)]^-$ anion, **270**, 130
kinetics and mechanism of the oxidation of ammineruthenium(II) complexes by bromine, **270**, 189
solution and crystal structure of the dihydrogen complex $[Ru(H_2)(H)(PMe_2Ph)_4]PF_6$, an active alkyne hydrogenation catalyst, **270**, 238
synthesis, characterization and application of Ru(BINAP)(Acac)-(MNAA)(MeOH), a highly effective catalyst for the asymmetric hydrogenation of 2-(6′-methoxynaphth-2′-yl)acrylic acid, **270**, 247
catalysis of aerobic olefin oxidation by a ruthenium perhaloporphyrin complex, **270**, 433
unique complexation by the tridentate ligands, 6-diphenylphosphino-2-pyridonate (pyphos) and 6-diphenylphosphino-2-pyridone (pyphosH), on ruthenium(II): crystal structures of (pyphosH)-$RuCl(\mu\text{-}Cl)Mo_2Cl(pyphos)_3$, $RuCl_2(CO)(pyphosH)_2$, $[Ru(pyphos)_3](\mu\text{-}Na)_2[Ru(pyphos)_3]$, **270**, 459
concomitant coordination of vinylidene and carbene ligands made possible at ruthenium(II), **272**, 1
the reaction of $Ru(CO)_2(MeCO_2)_2(PR_3)_2$ or $Ru_2(CO)_4\text{-}(MeCO_2)_2(PR_3)_2$ [R = Bu, Ph] with H_2 in the presence of Na_2CO_3: a stoichiometric carbon dioxide activation, **272**, 141
the stereochemistry of the oxidation of diamines and of the reduction of diimines in the coordination sphere of ruthenium(II), **272**, 153
mono- and bis-alkynyl ruthenium(II) complexes containing the ferrocenyl moiety; crystal structure of *trans*-$[Ru(C{\equiv}CC_5H_4FeC_5H_5)_2\text{-}(Ph_2PCH_2CH_2PPh_2)_2]$ and electrochemical studies, **272**, 188

chelate ring-opening ruthenium complexes: X-ray crystal structure and solution studies of *cis,trans*-bis(2-dimethyl-aminoethyl)diphenylphosphino(dichloro)ruthenium(II), **273**, 1
rhodium(III) analogues of antitumour-active ruthenium(III) compounds: the crystal structure of [ImH][*trans*-$RhCl_4(Im)_2$] (Im = imidazole), **273**, 62
crystalline dihydrogen complexes. Intramolecular and intermolecular interactions and dynamic behavior, **273**, 116
oxidation of amino esters to imine complexes with chiral high-valent oxoruthenium porphyrins, **273**, 142
syntheses of ruthenium hydridotris(1-pyrazoryl)borate complexes having sulfur-donor coligands, **273**, 238
synthesis, characterization and crystal structure of *trans*-aquatrichlorobis(5,7-dimethyl[1,2,4]triazolo[1,5-*a*]pyrimidine-N^3)-ruthenium(III) monohydrate, **273**, 259
photosubstitution reactivity, crystal structures, and electrochemistry of ruthenium(II)(III) complexes containing tetradentate (O_2N_2, S_2N_2, and P_2N_2) Schiff base ligands, **274**, 184
electrophilic reactivity of arene ligands coordinated to ruthenium carbido cluster complexes, **274**, 201
crystal and molecular structure of $[Ru(bpy)_2(2,3\text{-dpp})]Cl_2 \cdot 3H_2O \cdot CH_3CN$ and ^1H and ^{99}Ru NMR spectra of $[Ru(bpy)_2(2,n\text{-dpp})][PF_6]_2$ (bpy = 2,2′-bipyridine, dpp = bis(2-pyridyl)-pyrazine, $n = 3$ or 5), **275–276**, 320
4′-(4-fluorophenyl)-2,2′:6′,2″-terpyridine – an aryl-2,2′:6′,2″-terpyridine with an electron-releasing aryl substituent, **275–276**, 359
resonance Raman investigation of mixed-valence dinuclear and trinuclear complexes of Ru, Os and Re, **275–276**, 380
syntheses and structures of some Ru_2^{5+} complexes containing weakly coordinating anions as axial ligands, **275–276**, 447
toward a novel metal based chemotherapy against tropical diseases. 4. Synthesis and characterization of new metal-clotrimazole complexes and evaluation of their activity against *Trypanosoma cruzi*, **275–276**, 528
three reactions of Ru_2^{5+} compounds of the paddlewheel type that lead to cleavage of the Ru–Ru bond, **275–276**, 541
migration of $[Ru^{II}(hedta)]^-$ among three binding sites on 6-azauridine, **277**, 46
Fourier transform IR study of the interaction of $[(\eta^5\text{-}C_5H_5)_2FeRu(CO)_4]$ with hydroxylated alumina surfaces, **277**, 61
liquid secondary ion mass spectrometric study of diruthenium(II,III) complexes, **277**, 76
photoinduced processes in 4′-(9-anthryl)-2,2′:6′,2″-terpyridine, its protonated forms and Zn(II), Ru(II) and Os(II) complexes, **277**, 225
a simple and convenient synthesis of *cis*/*trans*-$RuH_2(Ph_2PCH_2PPh_2)_2$ and of *trans*-$RuHCl(Ph_2PCH_2PPh_2)_2$, **278**, 226
the use of photogenerated intermediates in the study of cluster build-up reactions: the generation of $Ru_6C(CO)_{17}$ from $Ru_3(CO)_{12}$, **278**, 250
synthesis of tetracarboxylatediruthenium(II) compounds: reduction and disproportionation reactions of diruthenium(II,III) units. Crystal structure of $Ru_2(\mu\text{-}O_2CCOPh)_4(thf)_2$, **279**, 30
stereochemically rigid mono and bis pyridazine complexes of $[Ru^{II}(hedta)]^-$ ($hedta^{3-}$ = *N*-hydroxyethylethylenediaminetriacetate), **279**, 85
some chlorocarbonylruthenium(II) complexes of P,N-donor ligands: crystal structures of $[RuCl(CO)\{PPh_2CH_2C(Bu^t)=NNH_2\}_2]Cl$ and *fac,cis*-$[RuCl_2(CO)\{PPh_2CH_2C(Bu^t)=N\text{–}N=C(Bu^t)\text{-}CH_2PPh_2\}]$, **279**, 95
weak intermolecular antiferromagnetic exchange in $\{[Ru_2(O_2CCH_3)_4(L)]X\}_n$ polymers (L = 4,4′-dipyridine and 1,4-diazabicyclooctane, X = PF_6^- or BPh_4^-), **279**, 121
protonation of the ground states of ruthenium(II) photosensitizers, **279**, 200
intramolecular protonation and other mechanisms for substitution reactions of hydrido(thiolato)– and di(thiolato)–ruthenium(II) phosphine complexes, **280**, 75
ammonia ruthenium complexes for the access to new hydrido, carbonyl and isocyanide ruthenium-alkynyl derivatives, **280**, 118
effect of ruthenium precursor on hydrogen-treated active carbon supported ruthenium catalysts for ammonia synthesis, **280**, 183
binuclear and tetranuclear complexes prepared by reactions of dodecacarbonyltriruthenium with the N-substituted cinnamic acid amides. Crystal and molecular structures of $[Ru_2(\mu\text{-}H)(\mu\text{-}\eta^3\text{-}PhC{=}CHC(O)N(CH_2)_4)(CO)_6]$ and $[Ru_4(\mu_3\text{-}H)_2(\mu\text{-}\eta^3\text{-}PhC{=}CHC(O)N(CH_2)_4)_2(CO)_{10}]$, **280**, 243
synthesis and properties of the dichloro[tetramethyl(trifluoromethyl)-cyclopentadienyl]ruthenium dimer: X-ray structure of $[Ru_2(\eta^5\text{-}C_5Me_4CF_3)_2Cl_2(\mu\text{-}Cl)_2]$ (M = Ru, Rh), **280**, 249
chiroptical properties of heterotrimetallic 'wing-tip' butterflies: synthesis and crystal structure of $(\mu\text{-}H)Ru_2Fe_2PdC(CO)_{12}(\eta^3\text{-}\beta\text{-}C_{10}H_{15})$, **280**, 275
synthesis and characterization of mono-, bi- and tetra-ruthenium(II)-phenanthroline complexes with dipyrido(2,3-*a*:2′,3′-*h*)phenazine (dpop), **281**, 1
comparison of the reactivity of *cis*-$[RuCl_2(DMSO)_4]$ and *trans*-$[RuCl_2(DMSO)_4]$ towards nucleosides, **281**, 10
linkage isomers of alizarin-bis(bipyridine)ruthenium(II), **281**, 126
substitution reaction of aqua(ethylenediaminetetraacetato) ruthenium(III) with 2-mercaptobenzoic acid, **282**, 38
pentamethylcyclopentadienyl ruthenium(II) complexes containing chiral diphosphines: synthesis, characterisation and electrochemical behaviour. X-ray structure of $(\eta^5\text{-}C_5Me_5)Ru\{(S,S)\text{-}Ph_2PCH(CH_3)CH(CH_3)PPh_2\}Cl$, **282**, 163
the effect of second-sphere coordination. 7. Isolation of 18-crown-6 ether adducts of ruthenium–ammine complexes, **282**, 247
intercalation into calf thymus DNA of 2-(4-arylphenyl)imidazo-[4,5-f]-[1,10]phenanthroline (aryl = -OMe, $-NMe_2$ or $-NO_2$) in its bipyridyl ruthenium(II) complex, **283**, 17
preparation and properties of a series of (μ-oxo)di(μ-acetato) diruthenium(III) complexes, $[Ru_2(\mu\text{-}O)(\mu\text{-}CH_3COO)_2(pyridine)_4(L)_2]^{2+}$ and $[Ru_2(\mu\text{-}O)(\mu\text{-}CH_3COO)_2(AA)_2(L)_2]^{2+}$, where AA is 2,2′-bipyridine or 1,10-phenanthroline, and L is a monodentate ligand, **283**, 151

Ruthenium(II) complexes
the reactivity of five-coordinate Ru(II)(1,4-bis(diphenylphosphino)-butane) complexes with the N-donor ligands: ammonia, pyridine, 4-substituted pyridines, 2,2′-bipyridine, bis(*o*-pyridyl)amine, 1,10-phenanthroline, 4,7-diphenylphenanthroline and ethylenediamine, **267**, 209

Salicylaldehyde-2-mercaptoanil complexes
salicylaldehyde-2-mercaptoanilato(2−)-2-hydroxyphenyl-benzthiazolato-nitrosyl-molybdenum(II) — a chelate which simultaneously contains an azomethine ligand and its cyclic oxidation product, **269**, 350

Salicylaldiminate complexes
planar or tetrahedral coordination in copper(II) complexes induced by bromo substitution, **267**, 151

Salicylamide derivative complexes
crystal structure and characterization of binuclear copper(II) complex with *N*-(2-hydroxyphenyl)-3-methoxycarbonyl-4,6-dimethylsalicylamide, **277**, 134

Samarium complexes
syntheses, properties and structural characteristics of several new (polypyrazolylborato)samarium(III) complexes: agostic interaction of the BH_2 proton with samarium, **277**, 8

Schiff base complexes
alkylcobalt chelates with Schiff bases derived from a β-diketone bearing both alkyl and aryl groups, **270**, 169

3-ferrocene-substituted 3-mercaptopropenales and their Schiff bases as ligands for transition metal complexes, **271**, 40
synthesis, spectroscopic and electrochemical study of nickel(II) complexes with tetradentate asymmetric Schiff bases derived from salicylaldehyde and methyl-2-amino-1-cyclopentenedithiocarboxylate, **271**, 83
cyclopalladated complexes of Schiff bases of homoveratrylamine and tryptamine. Synthesis and CO insertion, **272**, 18
functionalized acyclic Schiff bases and related complexes with d- and f-metal ions, **272**, 235
Ni(II) and Cu(II) Schiff base complexes with an extended H-bond network, **272**, 283
copper(II) complexes with aminosugar derived Schiff bases as ligands, **273**, 191
general synthesis of 'salicylaldehyde half-unit complexes': structural determination and use as synthon for the synthesis of dimetallic or trimetallic complexes and of 'self-assembling ligand complexes', **274**, 73
photosubstitution reactivity, crystal structures, and electrochemistry of ruthenium(II)(III) complexes containing tetradentate (O_2N_2, S_2N_2, and P_2N_2) Schiff base ligands, **274**, 184
the Co–CH_3 bond in Schiff base B_{12} models: influence of the *trans* and equatorial ligands as assessed by Fourier transform Raman spectroscopy, **275–276**, 90
characterisation of covalent copper and manganese organometallic complexes with Schiff bases by ionspray mass spectrometry, **275–276**, 462
synthesis and X-ray crystal structure of an oxorhenium(V) complex with a double Schiff base, **275–276**, 551
five-coordinate Salen(^{t}Bu) complexes of zinc, **277**, 157
preparation and electrochemistry of six-coordinate monooxo molybdenum(VI) complexes containing bidentate catecholate and tridentate NOS-donor Schiff base ligands, **278**, 1
spectroscopic and magnetic properties and structure of a five-coordinate, O_2-binding cobalt(II) Schiff base complex and of the copper(II) analogue, **278**, 127
synthesis and characterization of a neutral, low spin iron(III) complex of a hexadentate tripodal ligand containing three imidazolate arms. Use as a dinucleating agent, **278**, 197
synthesis and structural characterization of metal complexes of an unsymmetric N_3O tetradentate Schiff base ligand, **279**, 111
electrophoretic study of cationic cobalt(III) complexes with [N_2O] and [N_2O_2] Schiff base ligands, **280**, 295
mononuclear, dinuclear, and polynuclear dioxomolybdenum(VI)–tridentate Schiff base complexes: crystal structures and properties, **282**, 209

Schiff-base macrocycle complexes
complexes of lead(II) and lanthanide(III) ions with two novel 26-membered-imine and -amine macrocycles derived from 2,6-bis(2-formylphenoxymethyl)pyridine, **267**, 55

Second sphere coordination
second sphere coordination behavior of aquo and amine ligands bound to a η^6-benzeneruthenium(II) cation, **270**, 87

Selenaporphyrin complexes
formation and reactivity of paramagnetic organometallic nickel complexes of 21-oxa- and 21-selenaporphyrins — ^{1}H NMR and EPR investigations, **273**, 184

Selenium complexes
synthesis and crystal structure of [Cd{(c-C_6H_{11})$_2$P(Se)C(S)NPh}$_2$], **269**, 317
polynuclear silver compound formed from aggregation of Se^{2-} and Ag(I)-thiolate complex. Synthesis, structure and spectroscopic characterization of $Ag_{11}(\mu_5\text{-Se})(\mu_4\text{-Et}_2NCS_2)_3(\mu_3\text{-Et}_2NCS_2)_6$, **277**, 171

Seleno-bridged complexes
new efficient synthesis of the triangular cluster $[W_3Se_4(H_2O)_9]^{4+}$ and X-ray structure of the first mixed-metal seleno-bridged cubane-type tungsten–tin cluster $(Me_2NH_2)_6[(SCN)_9W_3Se_4SnCl_3]\cdot 0.5H_2O$, **269**, 292

Selenomolybdate complexes
syntheses and spectroscopic characterizations of the compounds containing univalent coinage metal and tetraselenomolybdate. Crystal structure of $[MoSe_4(AuPPh_3)_2]$, **271**, 93

Selenooxalate complexes
bis(1,2-diselenooxalato)nickelate(II) and -cuprate(II) — the first diselenooxalate complexes, **278**, 118

Self-assembly
dinuclear metal complexes of Cd(II), Zn(II) and Fe(II) with triple-helical structure and predetermined chirality, **271**, 36

Semiconducting polymers
new type of polymer containing 1,1′-ferrocenediyl unit in π-conjugated main chain. Preparation and redox behavior, **269**, 310

Semiquinone-bridged complexes
effects of ligand topology on the properties of dinuclear ruthenium complexes of bis-semiquinone bridging ligands, **267**, 1

Semiquinonediiminato complexes
kinetics and mechanism of the axial substitution reactions of (ligand)bis(4,5-dichloro-1,2-benzosemiquinonediiminato)-cobalt(III) complexes, **270**, 326

Sequestering agents
a novel iron sequestering agent: synthesis and iron-chelating properties of 1,1,1-tris(3-hydroxy-2-oxo-1,2-dihydro-1-pyridylpropoxymethyl)ethane, **281**, 60

Seven-coordinate complexes
synthesis and crystal structure of a seven-coordinate dinuclear iron(III) complex containing only bidentate ligands, **274**, 224

Seven-membered rings
tetrahedral coordination by a seven-membered chelate ring, **275–276**, 279

Siderophores
siderophore analogues. A new macrocyclic *bis*-(amine, amide, hydroxamate) ligand. Synthesis, solution chemistry, electrochemistry and molecular mechanics calculations for the iron complex, **278**, 51

Silanes
reactivity studies on cationic non-hydridic triruthenium carbonyl clusters. Reactions of $[Ru_3(\mu_3\text{-ampy})(CO)_{10}][BF_4]$ with hydrogen, triethylsilane and triphenylstannane, **268**, 131

Silica gels
a catalytic activity of glass beads, silica gels and anion-exchange resins modified with metal-porphines in oxidative reactions of ascorbic acid, **283**, 44

Silicon complexes
stereochemistry of six-coordinated silicon complexes. V. Racemization and geometrical isomerization of tris(4-isopropyltropolonato)-silicon(IV), **271**, 24

Silver clusters
photoreduction of methyl-viologen by zeolite-Y stabilized silver clusters, **269**, 253

Silver complexes
synthesis and structure of a novel bis(μ-η^2-thianthrene)disilver(I) bis(perchlorate), **271**, 145
2-(diphenylphosphino)pyridine (dppy) as monodentate or bridging ligand in homoleptic silver(I) complexes. Crystal structures of $[Ag(\eta^1\text{-dppy})_4][ClO_4]$ and $[Ag_2(\eta^1\text{-dppy})(\mu\text{-dppy})_2][ClO_4]_2$, **277**, 111
polynuclear silver compound formed from aggregation of Se^{2-} and Ag(I)-thiolate complex. Synthesis, structure and spectroscopic characterization of $Ag_{11}(\mu_5\text{-Se})(\mu_4\text{-Et}_2NCS_2)_3(\mu_3\text{-Et}_2NCS_2)_6$, **277**, 171
silver(I)–silver(I) interactions in the tetrameric silver thiolate phosphine complex $[Ag_4(SPh)_4(PPh_3)_4]$, **278**, 229
silver(I) complexes with heterocyclic thiones and tertiary phosphines as ligands. Part 3. Solution behaviour of mononuclear complexes

of silver(I) nitrate. The crystal structure of bis(triphenylphosphino)bis(benzoxazoline-2-thione)silver(I) nitrate, **281**, 121
coordination chemistry of the sterically hindered N_3-donor hydrotris(3,5-diphenylpyrazol-1-yl)borate toward silver(I)triorganophosphino compounds. Synthesis, structural and spectroscopic characterization, **282**, 1
phosphinoamino complexes of copper(I) and silver(I): a combined NMR and IR spectroscopy, and electrospray ionization and fast atom bombardment mass spectrometry analysis, **282**, 119
Silver(I) complexes
a tetranuclear silver complex with dppm and i-mnt ligands. Synthesis and structural characterization of $[Ag_4(\mu\text{-dppm})_4(\mu_4\text{-i-mnt})_2]\cdot 0.5$ DMF (i-mnt = 2,2-dicyano-1,1-ethylenedithiolate; dppm = bis(diphenylphosphino)methane), **267**, 313
stability constants of copper(I) and silver(I) complexes with aza-crown ethers and cryptands in dimethylsulfoxide, **279**, 186
Silylated cyclopentadienyl complexes
synthesis, immobilization and catalytic activity of some silylated cyclopentadienyl rhodium(I) complexes, **280**, 21
Silyl-bridged cyclopentadienyl complexes
cis- and *trans*-titanium complexes with doubly silyl-bridged dicyclopentadienyl ligands: molecular structure of $[(TiCl)_2(\mu\text{-O})\{(SiMe_2)_2(\eta^5\text{-}C_5H_3)_2\}]_2(\mu\text{-O})_2$, **280**, 1
Silylene-bridged complexes
preparation of doubly-silylene-bridged zirconocene alkyl complexes, $(Me_2Si)_2\{\eta^5\text{-}C_5H_2\text{-}4\text{-}CHMe_2\}\{\eta^5\text{-}C_5H\text{-}3,5\text{-}(CHMe_2)_2\}ZrR_2$ (R = CH_3, CH_2Ph) and investigations of their activity in 1-pentene polymerization. Molecular structure of $(Me_2Si)_2\{\eta^5\text{-}C_5H_2\text{-}4\text{-}CHMe_2\}\{\eta^5\text{-}C_5H\text{-}3,5\text{-}(CHMe_2)_2\}Zr(CH_2Ph)_2$, **280**, 226
Silylformylation
on the mechanism of silylformylation catalyzed by Rh–Co mixed metal complexes, **270**, 279
Site-directed mutagenesis
roles of valine-98 and glutamic acid-72 of putidaredoxin in the electron-transfer complexes with NADH-putidaredoxin reductase and P450cam, **272**, 80
Six-coordinate complexes
a mechanistic study of thermal C–H reductive elimination from a six-coordinate d^6 iridium complex, **270**, 537
spin crossover in six-coordinate $[Fe(L)_2(NCX)_2]$ compounds with L = DPQ = 2,3-bis-(2′-pyridyl)-quinoxaline, ABPT = 4-amino-3,5-bis(pyridin-2-yl)-1,2,4-triazole and X = S, Se: synthesis, magnetic properties and single crystal studies, **274**, 1
S,N,S-ligand complexes
crystal structure and fluxional behavior of novel Pd(II) allyl complexes containing a potentially tridentate S,N,S ligand: an example of selective *syn–anti* isomerism, **277**, 247
Sodium complexes
synthesis, characterization and crystal structures of complexes of sodium hexafluorophosphate with dibenzo-18-crown-6 and dibenzo-24-crown-8 macrocycles, **282**, 76
Sol–gel matrices
synthesis, immobilization and catalytic activity of some silylated cyclopentadienyl rhodium(I) complexes, **280**, 21
Solid state
electronic properties of some inorganic solids, **270**, 252
Soluble domains
spectroscopic investigations on the water-soluble fragment of the Rieske [2Fe–2S] protein from *Paracoccus denitrificans*, **275–276**, 493
Solvent effects
solvation, preferential solvation and complexation by the solvent of peroxovanadium complexes studied by ^{51}V NMR spectroscopy. Correlations with the oxidative reactivity, **272**, 62
Solvent exchange
fast water exchange on a Pt(II) center: variable temperature and pressure ^{17}O NMR study at 14.1 tesla of the ionic arylplatinum species $[Pt\{C_6H_3(CH_2NMe_2)_2\text{-}2,6\}(OH_2)]^+(OSO_2CF_3)$, **269**, 322
Solvent extraction
role of the intramolecular hydrogen bond and ligand rigidity in the complexation of trifluoroacetylcycloalkanones with lanthanides: novel strategy for the design of organic ligands of high selectivity, **267**, 201
Solvent mediated assembly
synthesis and single crystal X-ray structure of a novel trinuclear copper(II) methoxide complex, **269**, 241
Solvent motion
^2E relaxation of Cr(III) complexes in rigid and fluid media, **277**, 211
Solvothermal reactions
solvothermal synthesis and crystal structure of $[La(ethylenediamine)_4Cl]In_2Te_4$: A 1-D indium telluride, **273**, 255
Speciation
potentiometric and computer studies of yttrium–EDTMP, **275–276**, 37
Spectroelectrochemistry
conductive polymeric films of *meso*-tetrakis[2,2]-*para*-cyclophanylmetalloporphyrins — spectroelectrochemistry and catalytic properties, **271**, 167
Spin crossover
spin crossover in six-coordinate $[Fe(L)_2(NCX)_2]$ compounds with L = DPQ = 2,3-bis-(2′-pyridyl)-quinoxaline, ABPT = 4-amino-3,5-bis(pyridin-2-yl)-1,2,4-triazole and X = S, Se: synthesis, magnetic properties and single crystal studies, **274**, 1
spin crossover in bis(2,6-bis(benzimidazol-2-yl)pyridine) iron(II) tetraphenylborate, **278**, 190
Spin Hamiltonian parameters
angular overlap calculations of the spin Hamiltonian parameters of transition metal ions in low symmetry environments. High spin iron(II), iron(III) and manganese(III), **274**, 90
Spin transition
structural and electronic studies of the coordination of 6-(thiazol-2-yl)-2,2′-bipyridine and related systems to Fe(II), Co(II) and Ni(II), **274**, 32
Squarate complexes
synthesis, crystal structures and magnetic properties of $[Fe_2(bpym)(C_5O_5)_2(H_2O)_4]\cdot 2H_2O$ and two polymorphs of $[Fe_2(bpym)(C_4O_4)_2(H_2O)_6]\cdot 2H_2O$ (bpym = 2,2′-bipyrimidine), **279**, 127
Square-pyramidal Cu(II)-edda-type complexes
square-pyramidal copper(II) complexes of linear tetradentate edda-type ligands forming six-membered rings. Molecular structures of $[Cu(1,3\text{-pdda})(H_2O)]$ and $[Cu(eddp)(H_2O)]\cdot 3.5H_2O$, **268**, 221
Stability constants
interaction of mono- and triphosphate anions with a protonated polyaza macrocycle and its Cu(II) complexes, **270**, 207
stability of binary and ternary copper(II) complexes of the diphosphate analogue, methylphosphonylphosphate, in aqueous solution, **273**, 101
macrocyclic ligand design: structure–function relationships involving the interaction of cobalt(II), nickel(II) and copper(II) with mixed donor macrocyclic ligands, **273**, 372
electrostatic ligand–ligand interactions in ternary copper(II) complexes with 3,5-diiodo-L-tyrosine and polar amino acids, **278**, 136
the effect of end-group substitutions and use of a mixed solvent system on β-diketones and their iron complexes, **278**, 170
stability constants of copper(I) and silver(I) complexes with aza-crown ethers and cryptands in dimethylsulfoxide, **279**, 186
metal ion-coordinating properties of imidazole and derivatives in aqueous solution: interrelation between complex stability and ligand basicity, **280**, 50
a novel iron sequestering agent: synthesis and iron-chelating properties of 1,1,1-tris(3-hydroxy-2-oxo-1,2-dihydro-1-pyridylpropoxymethyl)ethane, **281**, 60

ternary complexes in solution. Intramolecular stacking interactions in mixed ligand complexes formed by copper(II), 2,2′-bipyridyl or 1,10-phenanthroline and a pyrimidine-nucleoside 5′-diphosphate (CDP^{3-}, UDP^{3-}, $dTDP^{3-}$), **283**, 193

Stacking
effect of substitution on packing in *trans*-di(*Z*-chlorobenzoato-*O*)-di(1,3-diaminopropane-*N*,*N*′)nickel(II) (*Z* = 2, 3 or 4) complexes, **267**, 249

Stannanes
reactivity studies on cationic non-hydridic triruthenium carbonyl clusters. Reactions of $[Ru_3(\mu_3\text{-ampy})(CO)_{10}][BF_4]$ with hydrogen, triethylsilane and triphenylstannane, **268**, 131

Stannylene complexes
synthesis, structure, spectroelectrochemical and theoretical investigations of $[(\eta^5\text{-}Me_5Cp)\overline{Co(\eta^4\text{-}MeC{=}CMe{-}CMe{=}CMe)Sn}\{CH(SiMe_3)_2\}_2]$. Formation of a cobalt complexed stannacycle via a mixed alkyne–stannylene cycloaddition reaction, **281**, 53

Stellacyanin
probing the metal site in *Rhus vernicifera* stellacyanin by Ni(II) substitution and paramagnetic NMR spectroscopy, **273**, 367

Stereochemistry
synthesis and crystal structure of the *trans*-dichloro(*N*,*N*′-bis-[2(*S*)-pyrrolidylmethyl]-ethane-1,2-diamine)cobalt(III) tetrachlorocobaltate(II) complex, **267**, 127
stereochemistry of six-coordinated silicon complexes. V. Racemization and geometrical isomerization of tris(4-isopropyltropolonato)-silicon(IV), **271**, 24

Stereoselectivity
stereoselectivity in reactions of metal complexes. Part XXI. Kinetics and mechanism of electron transfer between *rac*-$[Co(L)H_2O]^+$ (L = *N*,*N*′-[(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]) and optically active iron(II) complexes, **275–276**, 1
stereoselectivity in reactions of metal complexes. Part XXII. Kinetics and mechanism of electron-transfer reactions between *rac*-$[Co(L)X]^+$ (L = [(pyridine-2,6-diyl)bis(methylene)]bis[amino acid]; X = pyridine, imidazole, 1-methylimidazole and pyrazine) and optically active iron(II) complexes, **275–276**, 9

Stereospecific oxidation
the stereochemistry of the oxidation of diamines and of the reduction of diimines in the coordination sphere of ruthenium(II), **272**, 153

Stereospecific reduction
the stereochemistry of the oxidation of diamines and of the reduction of diimines in the coordination sphere of ruthenium(II), **272**, 153

Stibine complexes
rhodium(III) ditertiary stibine complexes, **271**, 203
trans influence of triphenylstibine. Crystal and molecular structures of *cis*-$[PtCl_2(SbPh_3)_2]$ and *trans*-$[PtI_2(SbPh_3)_2]$, **277**, 236

Strong electron-acceptor ability
interaction between the NO^+ ligand and *trans* π-donor ligands in chloro(2-methyl-8-quinolinolato)nitrosyl(8-quinolinolato)-ruthenium(II) complexes, **267**, 147

Substituted cyclopentadienyl complexes
on the reduction behaviour of 2-methoxyethyl and 2-methylthioethyl substituted zirconocene dichlorides: crystal structure of $(\eta^5,\sigma\text{-}C_5Me_4CH_2CH_2S)_2Zr$, **280**, 257

Substitution kinetics
dissociative kinetics of 'lightly bonded ligands' *L* from the clusters $Rh_6(CO)_{15}L$ (*L* = DMSO, NCMe, cyclooctene, THF and EtOH), **268**, 189

Substitution reactions
kinetics and mechanism of the axial substitution reactions of (ligand)bis(4,5-dichloro-1,2-benzosemiquinonediiminato)-cobalt(III) complexes, **270**, 326
substitution reaction of aqua(ethylenediaminetetraacetato) ruthenium(III) with 2-mercaptobenzoic acid, **282**, 38

Sugar complexes
unusual coordination behavior of D-fructose towards dimethyltin(IV): metal-promoted deprotonation of alcoholic OH groups in aqueous solutions of low pH, **274**, 167

Sulfide complexes
complexes of niobium and tantalum with chloride, sulfide and oxide ligands: synthesis and crystal structure of the (tetrachloro)bis-(η^2-disulfido)(μ-oxo)(μ-η^2:η^2-disulfido) bismetallates $[Cl_2(S_2)Nb(\mu\text{-}O)(\mu\text{-}S_2)Nb(S_2)Cl_2]^{2-}$ and $[Cl_2(S_2)Ta(\mu\text{-}O)(\mu\text{-}S_2)Ta(S_2)Cl_2]^{2-}$ as their tetraphenylphosphonium salts, **274**, 251

Sulfide oxidation
desulfoviridin, the dissimilatory sulfite reductase from *Desulfovibrio desulfuricans* (Essex): new structural and functional aspects of the membranous enzyme, **275–276**, 52

Sulfido complexes
synthesis of the first sulfido-bridged octahedral rhenium(III) aqua ion $[Re_6S_8(H_2O)_6]$, **271**, 228
synthesis of double incomplete-cubane clusters $[PPh_4]_2[(\eta^5\text{-}C_5Me_5)MS_3Cu_3X_3]_2$ (M = Mo, W; X = Cl, Br) and reactions with PPh_3, dppm (1,2-bis(diphenylphosphino)methane), and $AsPh_3$, **283**, 136

Sulfite oxidation
metallodrug complexes that mediate DNA and lipid damage via sulfite autoxidation: copper(II) famotidine and iron(III) bis(salicylglycine), **275–276**, 314

Sulfonation
electrophilic substitution of metal-coordinated pyrazole: nitration, sulfonation and bromination of $[Co(NH_3)_5(pyzH)]^{3+}$ (pyzH = pyrazole), **277**, 89

Sulfoxide complexes
the first X-ray structures of chromium(III) sulfoxide complexes, **269**, 229

Sulfur complexes
transition metal complexes with sulfur ligands. Part CXXVII. Azido, halido and nitrido ruthenium complexes with sulfur-rich coordination spheres, **269**, 63
synthesis and structures of the cuboidal iron–sulfur–nitrosyl–phosphine clusters $[Fe_4S_3(NO)_4(PR_3)_3]^{0,1+}$ (R = Et, Pr^i, C_6H_{11}), **270**, 46
generation of mixed iron chalcogenide clusters $MFeX_n^+$ (M = V, Fe; X = O, S; *n* = 2, 3) in the gas phase, **273**, 111
solid-state synthesis of cluster compound $[MoCu_3OS_3(PPh_3)_3(CH_3COO)]$ with its structure, reactivity and optical non-linearities, **279**, 172
transition metal complexes with sulfur ligands. Part CXXIX. Retention and reactivity of the [Fe–NH–NH–Fe] chromophore in the iron sulfur diazene complex $[\mu\text{-}N_2H_2\{Fe(PPr_3)('S_4')\}_2]$ in exchange and oxidation processes. ('S_4'$^{2-}$ = 1,2-bis(2-mercaptophenylthio)ethane(2−)), **280**, 39
preparation and characterization of twin-cubane-type clusters with molybdenum–copper–sulfur cores: X-ray structure determination of $[\{(NH_3)_6(H_2O)(OH)Mo_3CuS_4(O)\}_2]Cl_4\cdot 8H_2O$, **282**, 50
folding properties of iron–sulfur proteins, **283**, 12
a synthetic iron-containing prismane complex: peculiar electron-delocalization and spin-coupling properties deduced from Mössbauer and magnetization studies, **283**, 111

Sulfur dioxide
palladium(II)-catalyzed alternating copolymerization of alkenes with sulfur dioxide, **270**, 8

Sulfurization
unprecedented facile reactions of propane and cycloalkanes with elemental sulfur leading to dialkylsulfides and dicycloalkylsulfides, **280**, 355

Sulfur ligand complexes
transition metal complexes with sulfur ligands. Part CXXV. Synthesis and characterization of hydrido and chloro complexes with rhodium sulfur cores, **272**, 211

synthesis and crystal structure of dinuclear palladium(I) dppm complex with 4-mercaptopyridine in the thione mode. $[Pd_2(\mu\text{-dppm})_2(HpyS)_2](ClO_4)_2 \cdot CH_3OH$, **281**, 116

Superoxide complexes
oxidations in the H_2O_2/V^V/AcOH system. Evidence of vanadium(V) complexes with superoxide anion and singlet dioxygen molecule as ligands, **275–276**, 453

Superoxide dismutase complexes
proton ENDOR spectroscopy of cyanide-inhibited copper–zinc superoxide dismutase in randomly oriented samples, **273**, 131
the structure of the azide coordinated superoxide dismutase of *Propionibacterium shermanii* investigated by X-ray structure analysis, extended X-ray absorption fine structure, Mössbauer and electron paramagnetic resonance spectroscopy, **275–276**, 65
dynamic properties of bovine Cu,Zn superoxide dismutase from crystallographic data, **275–276**, 349
crystal structure and properties of a new five-coordinate manganese superoxide dismutase mimic, **277**, 21

Superoxide radicals
the role of $O_2^{\bullet -}$ radicals in DNA–metal complexes, **273**, 201

Superoxide scavenging
sterically hindered iron(II) complex of a new tripodal polyimidazole ligand: structure and reactivity toward superoxide, **275–276**, 234

Supramolecular chemistry
a large cavity reinforced cryptand for the binding of metal cations and anions, **273**, 326

Supramolecules
artificial ion channels of amphiphilic ion pairs composed of oligoether–ammonium and hydrophobic carboxylate or phosphates, **283**, 167

Suprofen complexes
crystal structure, spectroscopic, magnetochemical, thermoanalytical and electrochemical properties of binuclear copper(II) complexes of Suprofen, **268**, 239

Synchrotron radiation
Mössbauer spectroscopy with synchrotron radiation: a new technique entering biological inorganic chemistry, **275–276**, 334

Synthesis
reinforced piperazine rings containing polyamines: metal complex equilibria and structural studies, **268**, 63
synthesis, electrical properties of $[Ni(dmit)_2]^-$ complexes and X-ray crystal structure of $[EDA][Ni(dmit)_2]$ (dmit = 1,3-dithiol-2-thione-4,5-dithiolate), **268**, 103
reactivity of molybdenum complexes containing both NHNPhR and NNPhR ligands (R = Ph, Me) toward hydridotris(pyrazolyl)-borate, Tp and hydridotris(3,5-dimethylpyrazolyl)borate, Tp*, ligands. X-ray crystal structure of $[TpMo(NNMePh)_2Cl]$, **268**, 199
4′-(4-fluorophenyl)-2,2′:6′,2″-terpyridine – an aryl-2,2′:6′,2″-terpyridine with an electron-releasing aryl substituent, **275–276**, 359
Boc-Glu-Thr-Ile-His-OMe/Zn^{2+} in terms of the zinc-binding sites of proteases, **283**, 243

Tantalum complexes
the synthesis and structures of tantalum complexes that contain a triamido or a diamidoamine ligand, **270**, 353
synthesis and properties of $[M_6X_{12}(C_3H_7NO)_6]X_2$ (M = Nb, Ta; X = Cl, Br), **271**, 239
complexes of niobium and tantalum with chloride, sulfide and oxide ligands: synthesis and crystal structure of the (tetrachloro)bis-(η^2-disulfido)(μ-oxo)(μ-η^2:η^2-disulfido) bismetallates $[Cl_2(S_2)Nb(\mu\text{-}O)(\mu\text{-}S_2)Nb(S_2)Cl_2]^{2-}$ and $[Cl_2(S_2)Ta(\mu\text{-}O)(\mu\text{-}S_2)Ta(S_2)Cl_2]^{2-}$ as their tetraphenylphosphonium salts, **274**, 251

Taranakite
taranakite – the mineral with the longest crystallographic axis, **269**, 47

TCNQ anion complexes
synthesis and crystal structure of a new salt of the complex anion $[(TCNQ)_3]^{2-}$ $[Mo_2(\mu_2\text{-}S_2)_2(S_2CNMe_2)_4](TCNQ)_3$, **278**, 108

Technetium complexes
the synthesis and characterization of $[Tc(SPh)_2(N{=}NC_5H_4N)(HN{=}NC_5H_4N)]_2$, a paramagnetic dimer with bridging phenylthiolate ligands. The X-ray structure of $[Tc(SPh)_2(N{=}NC_5H_4N)(HN{=}NC_5H_4N)]_2$, **267**, 165
the synthesis and characterization of a technetium nitrosyl complex with *cis*-{2-pyridyl,diphenylphosphine} coligands. The X-ray crystal structure of $[TcCl_2(NO)(pyPPh_2\text{-}P,N)(pyPPh_2\text{-}P)]$, **267**, 319
metal–metal multiply-bonded complexes of technetium. 7. Oxidative decomposition of tetrachlorotetrakis(dimethylphenylphosphino)-ditechnetium(II) by an aminium hexachloroantimonate(V), **271**, 187
the synthesis and characterization of a technetium(III) isodiazene complex. The X-ray crystal structure of $[TcCl_3(N{=}NPh_2)(PPh_3)_2]$, **271**, 191
technetium and rhenium carbonyl complexes with 1,4-benzodiazepine derivatives: X-ray crystal structure, **272**, 267
the reaction chemistry of a technetium(I) nitrosyl complex with potentially chelating organohydrazines: the X-ray crystal structure of $[TcCl_2(NO)(HN{=}NC_5H_4N)(PPh_3)]$, **279**, 206
interaction of $[ReO(SNS)(S)]$ and $[^{99m}TcO(SNS)(S)]$ mixed ligand complexes with glutathione: isolation and characterization of the product, **281**, 148

Telluride complexes
solvothermal synthesis and crystal structure of $[La(ethylenediamine)_4Cl]In_2Te_4$: A 1-D indium telluride, **273**, 255
synthesis, structure characterization and magnetic properties of tellurostannates $[M(en)_3]_2(Sn_2Te_6)$ (M = Mn, Zn), **273**, 310

Tellurium complexes
triphenyltellurium(IV) dichalcogenoimidodiphosphinates containing novel six-membered TeXYP$_2$N chelate rings. X-ray structures of $Ph_3Te[(SPPh_2)(OPPh_2)N]$ and $Ph_3Te[(OPPh_2)_2N]$, **271**, 75

Template synthesis
complexes of lead(II) and lanthanide(III) ions with two novel 26-membered-imine and -amine macrocycles derived from 2,6-bis(2-formylphenoxymethyl)pyridine, **267**, 55
alkylcobalt chelates with Schiff bases derived from a β-diketone bearing both alkyl and aryl groups, **270**, 169
$(C_5H_7N_2O)Cu_3Br_4$: template synthesis of a two-dimensional copper(I) bromide lattice, **279**, 1

N,N,C-Terdentates
synthesis and properties of 3,2′-polymethylene-6-(2″-pyridyl)-2-phenylpyridines and their cycloplatinated complexes, **267**, 265

Terimine complexes
structural and electronic studies of the coordination of 6-(thiazol-2-yl)-2,2′-bipyridine and related systems to Fe(II), Co(II) and Ni(II), **274**, 32

Terminal alkynes
mechanisms of ≡C–H bond activation in oxidative carbonylations of alkynes catalyzed by palladium complexes, **280**, 202

Ternary complexes
Pd(gly-L-his-L-lys)Cl: solution structure and ternary complex formation with mono- and tetranucleotides, **273**, 31
stability of binary and ternary copper(II) complexes of the diphosphate analogue, methylphosphonylphosphate, in aqueous solution, **273**, 101
copper complexes modelling the interaction between benzolamide and Cu-substituted carbonic anhydrase. Crystal structure of Cu(bz)-$(NH_3)_4$ complex, **273**, 334
electrostatic ligand–ligand interactions in ternary copper(II) complexes with 3,5-diiodo-L-tyrosine and polar amino acids, **278**, 136
ternary complexes in solution. Intramolecular stacking interactions in mixed ligand complexes formed by copper(II), 2,2′-bipyridyl or

1,10-phenanthroline and a pyrimidine-nucleoside 5′-diphosphate (CDP^{3-}, UDP^{3-}, $dTDP^{3-}$), **283**, 193
the formation of ternary complexes between VO(maltolate)$_2$ and small bioligands, **283**, 202
Ternary copper(II) complexes
structures and properties of ternary copper(II) complexes with 4,5-dihydroxyphenylalanine or 2,4,5-trihydroxyphenylalanine and aromatic diamines, **283**, 260
Terpolymerization
palladium(II)-catalyzed alternating copolymerization of alkenes with sulfur dioxide, **270**, 8
Terpyridine complexes
remarkable substituent effects on the photophysics of Pt(4′-X-trpy)Cl^+ systems (trpy = 2,2′; 6′,2″-terpyridine), **273**, 346
interaction of organometallic cationic complex ions containing terpyridine ligands with nucleic acids: an investigation on aggregative phenomena, **275–276**, 242
photoinduced processes in 4′-(9-anthryl)-2,2′:6′,2″-terpyridine, its protonated forms and Zn(II), Ru(II) and Os(II) complexes, **277**, 225
Terpyridyl complexes
electron donating substituent effects on redox and spin state properties of iron(II) bis-terpyridyl complexes, **282**, 131
Tertiary phosphine complexes
syntheses, spectroscopic characterization and X-ray crystal structures of [HP-i-Pr_3]$_2$[$IrCl_6$] (**1**), [HP-i-Pr_3]$_2$[Ir_2Cl_8(P-i-Pr_3)] (**2**) and [HP-i-Pr_3][Ir_2Cl_7(P-i-Pr_3)$_2$] (**3**): three possible precursors of the complex Ir(H)(Cl)$_2$(P-i-Pr_3)$_2$ (**4**), **282**, 17
Tetradentate ligand complexes
complexes of thallium(I) and lead(II) with the potentially tetradentate ligand bis[3-(2-pyridyl)-pyrazolyl]dihydroborate, **267**, 323
Tetrahedral complexes
tetrahedral coordination by a seven-membered chelate ring, **275–276**, 279
Tetrakis(pyrazolyl)borate
synthesis, characterization and study of the redox properties of rhenium(V) diolates, **271**, 65
Tetranuclear complexes
a tetranuclear silver complex with dppm and i-mnt ligands. Synthesis and structural characterization of [Ag_4(μ-dppm)$_4$(μ_4-i-mnt)$_2$] · 0.5 DMF (i-mnt = 2,2-dicyano-1,1-ethylenedithiolate; dppm = bis(diphenylphosphino)methane), **267**, 313
structure and magnetic properties of a linear oximato-bridged tetranuclear copper(II) complex, **268**, 263
[cyclo{tetrakis-μ_3(η^2-S,η^1-N)-(1,3-thiazolidine-2-thionato)-bis-(η^1-S)-(1,3-thiazolidine-2-thione)tetracopper(I)}]: electrochemical synthesis, characterisation and crystal structure at 160 K, **271**, 57
pentamethylphenylcopper(I): a square-planar tetranuclear cluster, **277**, 232
silver(I)–silver(I) interactions in the tetrameric silver thiolate phosphine complex [Ag_4(SPh)$_4$(PPh_3)$_4$], **278**, 229
Tetrapeptide complexes
interaction of Hg(II) with tetrapeptides containing cysteinyl and histidinyl residues, **272**, 7
Tetrathiolate metalloproteins
the use of ^{113}Cd NMR chemical shifts as a structural probe in tetrathiolate metalloproteins, **273**, 279
Thallium complexes
complexes of thallium(I) and lead(II) with the potentially tetradentate ligand bis[3-(2-pyridyl)-pyrazolyl]dihydroborate, **267**, 323
electrochemical synthesis and characterisation of gallium, indium and thallium compounds with 1-hydroxopyridine-2-thione. The crystal structures of tris(1-oxopyridine-2-thionato)indium(III) and tris(1-oxopyridine-2-thionato)thallium(III), **281**, 70
Theoretical study
comparative DFT and HF-MP2 study on structure, relative stabilities and rotational barriers of end-on (η^1) and side-on (η^2) (PH_3)$_2$Pd HPPH model complexes, **273**, 72
Thermal quenching
^{2}E relaxation of Cr(III) complexes in rigid and fluid media, **277**, 211
Thermodynamics
cyanide binding to phthalocyaninatoiron(II) in dimethyl sulfoxide in the presence of carbon monoxide: kinetic and equilibrium study, **267**, 109
thermodynamics of the formation of *cis*-bis(dialkylsulfide)dimethylplatinum(II) complexes from bis(μ-dialkylsulfide)bisdimethylplatinum(II) and free dialkylsulfide, **268**, 311
thermodynamics of the binding of nitrogenous ligands to β-trifluoromethylcobinamide and β-cyanomethylcobinamide, **279**, 178
Thermolysis
intramolecularly alkylated Cosalen complexes: thermolysis and photolysis of coenzyme B_{12} models with a cobalt-to-ligand carbon bridge, **269**, 203
a mechanistic study of thermal C–H reductive elimination from a six-coordinate d^6 iridium complex, **270**, 537
Thianthrene complexes
synthesis and structure of a novel bis(μ-η^2-thianthrene)disilver(I) bis(perchlorate), **271**, 145
Thiazolidine-thionato complexes
[cyclo{tetrakis-μ_3(η^2-S,η^1-N)-(1,3-thiazolidine-2-thionato)-bis-(η^1-S)-(1,3-thiazolidine-2-thione)tetracopper(I)}]: electrochemical synthesis, characterisation and crystal structure at 160 K, **271**, 57
Thioamide complexes
arm-off reactions of the tripodal polyphosphane MeC(CH_2PPh_2)$_3$ in octahedral rhodium(III) complexes of heterocyclic thioamides, **267**, 39
Thiocarboxylate complexes
synthesis and characterization of molybdenum(VI)-dioxo complexes containing both coordinated thiolate and carboxylate groups. Reactions with their own free ligands, **268**, 145
modeling for the active site nitrate reductase. Oxidation of the complex [$Mo^VO(O_2CC(S)CH_3Ph)_2]^-$ by nitrate and nitrite in methanol, **271**, 174
Thiocyanate complexes
three reactions of Ru_2^{5+} compounds of the paddlewheel type that lead to cleavage of the Ru–Ru bond, **275–276**, 541
Thiocyanato-bridged complexes
{*catena*-($\mu_{N,S}$-NCS)[Ni(Medien)(NCS)]}: a new 1D nickel(II) complex with bridging thiocyanato ligands in *cis* position. Crystal structure and magnetic properties, **269**, 313
Thioester
bioinorganic chemistry of nickel and carbon dioxide: an Ni complex behaving as a model system for carbon monoxide dehydrogenase enzyme, **272**, 38
Thioether complexes
tripod ligands of constitutional C_1 symmetry containing a deliberate set of mixed P/S donor groups: synthesis and coordination chemistry, **269**, 143
synthesis and X-ray structure of the thioethereal mercury(II) complex [$N(CH_2CH_2SCHMe_2)_3HgCl]_2(Hg_2Cl_6)$, **269**, 274
Thioformamide complexes
synthesis and crystal structure of [Cd{(c-C_6H_{11})$_2$P(Se)C(S)NPh}$_2$], **269**, 317
binuclear diorganophosphinothioformamido complexes of gold(I): synthesis and crystal structure of [Au($Ph_2PC(S)NPh$)]$_2$ and related systems, **281**, 64
Thiolate complexes
synthesis and characterization of uranyl thiolate complexes. Crystal and molecular structures of [Me_4N][$UO_2(NO_3)_2(C_5H_4NS)$] and [$C_{16}H_{25}N_2S_2Si_2$][$UO_2(NO_3)_2(C_8H_{12}NSSi)$], **268**, 163

misdirected π-donor ligands: $(\eta^5\text{-}C_5H_5)_2Zr(Cl)(SR)$ with sterically more demanding R groups have lower rotational barriers, **270**, 227
silver(I)–silver(I) interactions in the tetrameric silver thiolate phosphine complex $[Ag_4(SPh)_4(PPh_3)_4]$, **278**, 229
intramolecular protonation and other mechanisms for substitution reactions of hydrido(thiolato)– and di(thiolato)–ruthenium(II) phosphine complexes, **280**, 75
binuclear diorganophosphinothioformamido complexes of gold(I): synthesis and crystal structure of $[Au(Ph_2PC(S)NPh)]_2$ and related systems, **281**, 64
zinc thiolate complexes with chelating nitrogen ligands, **282**, 193

Thiolato-bridged complexes
the synthesis and characterization of $[Tc(SPh)_2(N{=}NC_5H_4N)(HN{=}NC_5H_4N)]_2$, a paramagnetic dimer with bridging phenylthiolate ligands. The X-ray structure of $[Tc(SPh)_2(N{=}NC_5H_4N)(HN{=}NC_5H_4N)]_2$, **267**, 165

Thiomolybdates
synthesis and characterization of tetraalkylammonium thiomolybdates and thiotungstates in aqueous solution, **274**, 108

Thiomorphiline complexes
synthesis and metal complexes of a chelating ligand containing thiomorpholine and pyridine: 2,6-bis(thiomorpholinomethyl)pyridine, **279**, 221

Thionate complexes
electrochemical synthesis and characterisation of gallium, indium and thallium compounds with 1-hydroxopyridine-2-thione. The crystal structures of tris(1-oxopyridine-2-thionato)indium(III) and tris(1-oxopyridine-2-thionato)thallium(III), **281**, 70

Thione complexes
reactivity of $[NBu_4]_2[\{(C_6X_5)_2M(\mu\text{-}OH)\}_2]$ (X = F or Cl; M = Pd or Pt) towards heterocyclic thiones: crystal structure of $[NBu_4]_2[\{(C_6F_5)_2Pd(\mu\text{-}\eta^2\text{-}LL')\}_2]$ (LL′ = methylimidazole-2-thiolate), **281**, 165

Thiophene complexes
synthesis and characterization of heterobimetallic complexes containing C–S cleaved thiophenes, **272**, 55

Thio-pyrimidine
electrochemistry of 2-thio- and 2-oxo-pyrimidines in dimethyl formamide in the presence of dioxygen, **270**, 145

Thiosemicarbazone complexes
synthesis and NMR spectroscopy of dichlorobis(acetophenonethiosemicarbazone)mercury(II) formed from phenylmercury(II) chloride and acetophenonethiosemicarbazone: the first example of symmetrisation in organomercury(II)-thiosemicarbazone chemistry, **267**, 169
synthetic, spectroscopic and X-ray crystallographic studies on copper(II) complexes with pyruvic acid and pyridoxal thiosemicarbazones, **269**, 297
EPR characterization of a series of mono- and bis-thiosemicarbazone copper(II) complexes, **275–276**, 557
spectroscopic and magnetic properties of the pyridine-2-carbaldehyde thiosemicarbazonateiron(III) complexes: $[Fe(C_7H_7N_4S)_2]X\cdot nH_2O$ (X = Cl, ClO_4, NO_3, PF_6). Crystal structure of the hexafluorophosphate compound, **278**, 150

Thiosulfinato complexes
oxidative addition of 2,3,4-benzotrithiepin 1-oxide to platinum(0) compounds: a convenient synthetic route to platinum(II) thiosulfinato complexes, **273**, 106

Thiosulfinic acid anions
oxidative addition of 2,3,4-benzotrithiepin 1-oxide to platinum(0) compounds: a convenient synthetic route to platinum(II) thiosulfinato complexes, **273**, 106

Thiotungstates
synthesis and characterization of tetraalkylammonium thiomolybdates and thiotungstates in aqueous solution, **274**, 108

Thiourea complexes
synthesis, X-ray and spectroscopic characterization of $[SnI_2(mbit)_2](I_3)_2\cdot 2/3I_2$ obtained through the one-step reaction of mbit·$2I_2$ with tin metal powder (mbit = 1,1′-bis(3-methyl-4-imidazoline-2-thione)methane), **267**, 115

Thioxane
the kinetics and mechanism of the oxidation of 1,4-thioxane by potassium ferrate, **267**, 159

Thorium complexes
trimethylsilyl iodide as a halide exchange reagent in inorganic chemistry: synthesis and structure of $Cp^*_2ThI_2$ ($Cp^* = \eta^5\text{-}C_5Me_5$), **274**, 232

Three-coordinate complexes
synthesis of *mer*-trichlorotris(tetrahydrofuran)tungsten(III) by intermetal chlorine atom transfer, **278**, 103

Three-dimensional compounds
hydrothermal synthesis and structural characterization of the three-dimensional framework solids $(NH_4)[V_2(OH)(PO_4)_2(H_2O)]\cdot H_2O$ and $Rb_6[(Mo_6V_3O_6)(PO_4)_{10}(H_2PO_4)_3(OH)_9]\cdot 8.5H_2O$, **275–276**, 122

Three-dimensional structure
refinement of the three-dimensional structures of cytochrome c_3 from *Desulfovibrio vulgaris* Hildenborough at 1.67 Å resolution and from *Desulfovibrio desulfuricans* ATCC 27774 at 1.6 Å resolution, **273**, 213

L-Threonine dehydrogenase
structural characterization of the zinc site in *Escherichia coli* L-threonine dehydrogenase using extended X-ray absorption fine structure spectroscopy, **275–276**, 215

Thymine recognition
a new macrocyclic tetraamine, 2,4-dinitrophenylcyclen (= 1-(2,4-dinitrophenyl)-1,4,7,10-tetraazacyclododecane): synthesis, cation reporter properties, and recognition of 1-methylthymine by its zinc(II) complex, **270**, 424

Thymulin
metal ion–FTS nonapeptide interactions. II. Copper(II) complex equilibria and structures under physiological conditions, **283**, 211

Tin complexes
synthesis, X-ray and spectroscopic characterization of $[SnI_2(mbit)_2](I_3)_2\cdot 2/3I_2$ obtained through the one-step reaction of mbit·$2I_2$ with tin metal powder (mbit = 1,1′-bis(3-methyl-4-imidazoline-2-thione)methane), **267**, 115
new efficient synthesis of the triangular cluster $[W_3Se_4(H_2O)_9]^{4+}$ and X-ray structure of the first mixed-metal seleno-bridged cubane-type tungsten–tin cluster $(Me_2NH_2)_6[(SCN)_9W_3Se_4SnCl_3]\cdot 0.5H_2O$, **269**, 292
mechanistic details for metathetical exchange between XCO (X = O and RN) and the tin(II) dimer, $\{Sn[N(SiMe_3)_2](\mu\text{-}OBu^t)\}_2$, **270**, 118
crystal and molecular structures of some tri-aryltin isothiocyanates (Ar_3SnNCS; Ar = C_6H_5, *p*-$CH_3C_6H_4$, *o*-$CH_3C_6H_4$, *o*-$CH_3OC_6H_4$) and of the adduct hexamethylphosphoramide(*N*-thiocyanato)-tris(*o*-tolyl)tin, **282**, 30

Titanium complexes
an infrequent geometry for an element of the first transition series: structural characterization of a distorted eight-coordinate compound $TiCl_4[C_2H_4(PMe_2)_2]_2$, **267**, 173
partial hydrolysis of Ti(III) and Ti(IV) chlorides in the presence of $[PPh_4]Cl$, **267**, 271
high pressure liquid chromatography resolution of racemic mixtures of chiral tricarbonyl arene chromium(0) and cyclopentadienyl-titanium(IV) complexes, **268**, 323
titanium–carbon functionalities on an oxo surface defined by a calix[4]arene moiety and its redox chemistry, **270**, 298
extracting absolute titanium–alkyl and –hydride bond enthalpies from relative $D(TiR(H))$ in $(silox)_2(^tBu_3SiNH)TiR$: electronegativity and *ECT* models, **270**, 414

trimethylphosphine halide complexes of titanium, zirconium and hafnium in oxidation states IV, III and II. Crystal structure of $ML_3(PMe_3)_n$ (M = Zr, n = 3; M = Hf, n = 2.5), $M_2X_6(PMe_3)_4$ (M = Hf, Ti, X = I; M = Zr, X = Br) and (μ-η^6,η^6-C_6H_6)-$Hf_2I_4(PMe_3)_4 \cdot C_6H_6$, **271**, 180

synthesis of *mer*-trichlorotris(tetrahydrofuran)tungsten(III) by intermetal chlorine atom transfer, **278**, 103

reactions of Ti(O–i-Pr)Cl_3 with 2-aminophenol and the crystal structure of the zwitterionic complex [Ti(O–i-Pr)Cl_3(THF)-($^-OC_6H_4NH_3^+$)] · THF, **278**, 232

synthesis through an in situ esterification process and characterization of oxo isopropoxo titanium clusters, **279**, 144

cis- and *trans*-titanium complexes with doubly silyl-bridged dicyclopentadienyl ligands: molecular structure of [$(TiCl)_2$(μ-O)-{$(SiMe_2)_2(\eta^5$-$C_5H_3)_2$}]$_2$(μ-O)$_2$, **280**, 1

Titanium compounds

transformations of molecular nitrogen into aromatic amines under the action of titanium compounds, **280**, 264

T_1 measurements

in situ IR and NMR study of the interactions between proton donors and the Re(I) hydride complex [{$MeC(CH_2PPh_2)_3$}Re$(CO)_2$H]. ReH···H bonding and proton-transfer pathways, **280**, 302

Topology

chemical applications of topology and group theory. 30. Distortions from idealized symmetrical coordination polyhedra by lifting molecular orbital degeneracies, **270**, 68

Trans influence

trans influence of triphenylstibine. Crystal and molecular structures of *cis*-[$PtCl_2(SbPh_3)_2$] and *trans*-[$PtI_2(SbPh_3)_2$], **277**, 236

the *trans* influence of F, Cl, Br and I ligands in a series of square-planar Pd(II) complexes: relative affinities of halide anions for the metal centre in *trans*-[$(Ph_3P)_2Pd(Ph)X]^-$, **280**, 87

Transition metal π-complexes

ricochet inter-ring haptotropic rearrangement of σ-methyl-(η^5-indenyl)chromium tricarbonyls. Experimental kinetic and theoretical DFT study, **280**, 328

Transition metal complexes

the effect of net charge and π-backbonding contribution of ML_5 fragments containing Fe(II), Ru(II), Os(II), Co(III) and Rh(III) on the ^{1}H NMR shifts of coordinated N-heterocycles, **267**, 11

synthesis and spectroscopic characterization of copper, nickel and cobalt complexes with *N*-(*N*″,*N*″-disubstituted-aminothiocarbonyl)-*N*′-substituted benzamidines, **268**, 13

synthesis and coordination properties of amphiphilic 3-oxodithiocarboxylic esters, **269**, 83

ferrocenyl-derived face-capping ligands: syntheses and molecular structures of [(FcTt)Cu]$_4$ and [FcTt]$_2$M (M = Fe, Co, Ni; FcTt$^-$ = ferrocenyltris((methylthio)methyl)borate), **270**, 543

syntheses and properties of bivalent metal complexes with *N*-(2-hydroxyethyl)ethylenediamine-*N*,*N*′,*N*′-triacetic acid and the crystal structure of *N*-(2-hydroxyethyl)ethylenediamine-*N*,*N*′,*N*′-triacetato-copper(II) monohydrate, **271**, 13

dinuclear complexes of first transition series metals with 4,4′-dipyrazolylmethane: characterisation, DNA binding and anticancer properties, **271**, 151

macrocyclic ligand design: structure–function relationships involving the interaction of cobalt(II), nickel(II) and copper(II) with mixed donor macrocyclic ligands, **273**, 372

tetrahedral coordination by a seven-membered chelate ring, **275–276**, 279

potentiometric and spectroscopic studies of transition metal complexes of bis(imidazolyl) and bis(pyridyl) derivatives of amino acids, **283**, 233

Transmetallation

multinuclear NMR study of the interactions between the La(III) complex of DTPA-bis(glucamide) and Zn(II) or borate, **268**, 249

Triamido ligand complexes

the synthesis and structures of tantalum complexes that contain a triamido or a diamidoamine ligand, **270**, 353

Triamine complexes

hydrolysis of phosphate esters catalyzed by copper(II)-triamine complexes. The effect of triamine ligands on the reactivity of the copper(II) catalysts, **283**, 51

Triarylphosphine complexes

unexpected bell-shaped effect of the ligand on the rate of the oxidative addition to palladium(0) complexes generated in situ from mixtures of Pd(dba)$_2$ and *para*-substituted triarylphosphines, **273**, 76

Triazaacetate macrocycle complexes

NMR conformational study of diamagnetic complexes of some triazatriacetate macrocycles, **273**, 288

Triazenido complexes

molecular dynamics of *N*,*N*′-di-*p*-fluorophenyltriazenido complex of platinum(II), **280**, 238

Triazole-bridged complexes

synthesis, crystal structure and magnetic properties of bis[μ-bis(3-(pyridin-2-yl)-1,2,4-triazole-*N*′,*N*1,*N*2)] bis[triaqua nickel(II)] tetranitrate, **273**, 54

Triazole complexes

synthesis, X-ray structure, spectroscopic and magnetic properties of [bis(4-amino-3,5-bis(pyridin-2-yl)-1,2,4-triazole-*N*′,*N*1)-(aqua)copper(II)]bis(hydrogensulfate), **268**, 37

synthesis, crystal structure and magnetic properties of the one-dimensional chain compound *catena*-poly[aquabis(4,4′-bis-1,2,4-triazole-*N*1)(thiocyanato-*N*)-manganese-μ-[4,4′-bis-1,2,4-triazole-*N*1,*N*1′]thiocyanate], **282**, 257

Triazolopyrimidine complexes

synthesis, characterization and crystal structure of *trans*-aquatrichlorobis(5,7-dimethyl[1,2,4]triazolo[1,5-*a*]pyrimidine-N^3)-ruthenium(III) monohydrate, **273**, 259

ternary copper(II) complexes with the versatile 4,7-dihydro-5-methyl-7-oxo-[1,2,4]triazolo[1,5-*a*]pyrimidine ligand, **274**, 53

Tridentate amine complexes

a series of [$Cr_2(\mu$-OH)$_n$(μ-RCO$_2$)$_{3-n}$(bispma)$_2$] type chromium(III) dinuclear complexes (n = 1, 2) with tridentate bis(2-pyridylmethyl)amine (bispma): synthesis, crystal structure, and magnetic properties, **278**, 15

Tridentate ligand complexes

copper(II) complexes with tridentate O,N,O-chelate dianions and additive monodentate donor ligands. I. Crystal structures and EPR results of urea adducts of *N*-salicylidene-α-aminoacidato-copper(II) compared with *N*-salicylideneglycinato-thiourea-copper(II), **268**, 117

structural and electronic studies of the coordination of 6-(thiazol-2-yl)-2,2′-bipyridine and related systems to Fe(II), Co(II) and Ni(II), **274**, 32

Tridentate ONS ligand complexes

ligand exchange reactions of $ReCl_4(PPh_3)_2$ with tridentate diacidic ligands with the donor set O–N–O(S). Molecular structures of the resulting oxo-rhenium(V) complexes, **268**, 109

Tridentate phosphine complexes

quadruply bonded complexes containing η^2- or η^3-tridentate phosphine ligands; syntheses, structures and ^{31}P{^{1}H} NMR spectra of $Mo_2Cl_4(\eta^3$-etp)(L) and $Mo_2Cl_4(\eta^2$-etp)$_2$ (L = CH_3OH, PMe_3 and PEt_3; etp = $Ph_2PCH_2CH_2P(Ph)CH_2CH_2PPh_2$), **274**, 7

Triflate complexes

preparation of tetrabutylammonium octa-μ_3-bromohexa(trifluoromethanesulfonato)octahedro-hexamolybdate(2−), $(Bu_4N)_2$-[$Mo_6Br_8^i(CF_3SO_3)_6^a$] and other derivatives containing the $Mo_6Br_8^{4+}$ core, **278**, 241

Trifluoromethyl complexes

the crystal structure of trifluoromethylcobalamin, a coenzyme B_{12} analog, **267**, 305

synthesis and properties of the dichloro[tetramethyl(trifluoromethyl)-cyclopentadienyl]ruthenium dimer: X-ray structure of $[M_2(\eta^5\text{-}C_5Me_4CF_3)_2Cl_2(\mu\text{-}Cl)_2]$ (M = Ru, Rh), **280**, 249

Trifluorophosphine complexes
highly reduced organometallics. 42. A new method for the syntheses of $[V(CO)_6]^-$ and $[V(PF_3)_6]^-$ involving anthracenide mediated reductions of $VCl_3(THF)_3$, **269**, 58

Trigonal bipyramidal complexes
syntheses and structure of dinuclear cobalt(II) complexes of the polypyridine ligand 6,6′-bis[bis(2-pyridylmethyl)aminomethyl]-2,2′-bipyridine, **268**, 159
novel trigonal-bipyramidal copper(II) complexes containing one discrete amino acid chelate: crystal structures of [Cu(L-val)(pmdt)]ClO_4 and [Cu(aiba)(pmdt)]$ClO_4 \cdot H_2O$ (L-val = L-valinate ion, aiba = α-aminoisobutyrate ion, and pmdt = *N,N,N′,N″,N″*-pentamethyldiethylenetriamine), **274**, 247

Trinuclear complexes
DNA nicking by a trinuclear chromium complex, **268**, 211
some oxidative addition reactions with a diphosphadibora[1.1.0]bicyclobutane: formation of cage compounds and of an unusual trinuclear coordination compound of palladium, **269**, 101
synthesis and single crystal X-ray structure of a novel trinuclear copper(II) methoxide complex, **269**, 241
3-ferrocene-substituted 3-mercaptopropenales and their Schiff bases as ligands for transition metal complexes, **271**, 40

Trioxide complexes
selective C–H bond activation of arenes catalyzed by methylrhenium trioxide, **270**, 55

Tripeptide complexes
solution behaviour and zinc complexation of tripeptides with cysteine and/or histidine at both termini, **272**, 115
Pd(gly-L-his-L-lys)Cl: solution structure and ternary complex formation with mono- and tetranucleotides, **273**, 31
potentiometric and spectroscopic studies on the copper(II) and nickel(II) complexes of tripeptides of methionine, **275–276**, 440

Triphos complexes
arm-off reactions of the tripodal polyphosphane $MeC(CH_2PPh_2)_3$ in octahedral rhodium(III) complexes of heterocyclic thioamides, **267**, 39

Triphosphate complexes
interaction of mono- and triphosphate anions with a protonated polyaza macrocycle and its Cu(II) complexes, **270**, 207

Triple-decker complexes
heteroleptic triple-decker (phthalocyaninato)(porphyrinato)-europium(III) complexes: synthesis and electrochemical study, **268**, 49
the role of cyclo-triphosphorus (P_3) in-plane σ orbitals in coordinating transition metal fragments, **275–276**, 366

Triply-bridged complexes
triply-bridged diruthenium(II) 1,4-bis(diphenylphosphino)butane (dppb) and (*R*)-2,2′-bis(diphenylphosphino)-1,1′-binaphthyl (binap) complexes, including structural characterisation of [(dppb)ClRu(μ-D_2O)(μ-Cl)$_2$RuCl(dppb)], [(η^2-H_2)-(dppb)Ru(μ-Cl)$_3$RuCl(dppb)] and the [(dppb)ClRu(μ-Cl)$_3$RuCl(dppb)]$^-$ anion, **270**, 130

Tripodal ligand complexes
sterically hindered iron(II) complex of a new tripodal polyimidazole ligand: structure and reactivity toward superoxide, **275–276**, 234
synthesis and characterization of a neutral, low spin iron(III) complex of a hexadentate tripodal ligand containing three imidazolate arms. Use as a dinucleating agent, **278**, 197
a novel iron sequestering agent: synthesis and iron-chelating properties of 1,1,1-tris(3-hydroxy-2-oxo-1,2-dihydro-1-pyridylpropoxymethyl)ethane, **281**, 60
structures and electrochemical properties of the Co(III) ternary complexes containing NO_3-type tripodal tetradentate ligands and amino acids: effect of the outer coordination sphere on the electrochemical properties, **283**, 160

Tripodal phosphine complexes
the synthesis of the tripodal phosphine $CH_3C\{CH_2P(m\text{-}CF_3C_6H_4)_2\}_3$ and its rhodium coordination chemistry, **270**, 499

Tripodal P-ligand complexes
in situ IR and NMR study of the interactions between proton donors and the Re(I) hydride complex [{$MeC(CH_2PPh_2)_3$}Re$(CO)_2$H]. ReH···H bonding and proton-transfer pathways, **280**, 302

Tripod complexes
a tripod gadolinium(III) poly(amino carboxylate) relevant to magnetic resonance imaging: structural and dynamical ^{17}O NMR and ^1H NMRD studies, **275–276**, 15

Tripod ligand complexes
tripod ligands of constitutional C_1 symmetry containing a deliberate set of mixed P/S donor groups: synthesis and coordination chemistry, **269**, 143

Trisimidazole phosphine complexes
the synthesis, NMR and X-ray analysis of tris(1-ethyl-2-isopropylimidazol-5-yl)phosphine gold(I)-2,3,4,6-tetra-O-acetyl-1-thio-β-D-glucopyranoside, a bifunctional auranofin analogue, **274**, 137

Trithiepin 1-oxide
oxidative addition of 2,3,4-benzotrithiepin 1-oxide to platinum(0) compounds: a convenient synthetic route to platinum(II) thiosulfinato complexes, **273**, 106

Tropical diseases
toward a novel metal based chemotherapy against tropical diseases. 4. Synthesis and characterization of new metal-clotrimazole complexes and evaluation of their activity against *Trypanosoma cruzi*, **275–276**, 528

Tropolonato complexes
stereochemistry of six-coordinated silicon complexes. V. Racemization and geometrical isomerization of tris(4-isopropyltropolonato)-silicon(IV), **271**, 24

Trypanosoma cruzi
toward a novel metal based chemotherapy against tropical diseases. 4. Synthesis and characterization of new metal-clotrimazole complexes and evaluation of their activity against *Trypanosoma cruzi*, **275–276**, 528

Tumour-inhibiting complexes
imidazole release from the antitumor-active ruthenium complex imidazolium *trans*-tetrachlorobis(imidazole)ruthenate(III) by biologically occurring nucleophiles, **267**, 137

Tungstate anions
preparation and characterization of two novel heteropoly-fluorooxotungstate anions $[NaCoW_{11}O_{43}FH_{12}]^{6-}$ and $[NaNiW_{11}O_{43}FH_{12}]^{6-}$, **282**, 136

Tungsten complexes
new efficient synthesis of the triangular cluster $[W_3Se_4(H_2O)_9]^{4+}$ and X-ray structure of the first mixed-metal seleno-bridged cubane-type tungsten–tin cluster $(Me_2NH_2)_6[(SCN)_9W_3Se_4SnCl_3] \cdot 0.5H_2O$, **269**, 292
elimination of benzene from a benzyloxy ligand to form a carbonyl group on a tungsten-phosphine center, **270**, 202
the propensity of alkoxide and aryloxide derivatives of tungsten carbonyls to aggregate in solution. Synthesis and X-ray structures of dinuclear, trinuclear and tetranuclear complexes derived from the $MeOW(CO)_5^-$ anion, **270**, 405
first- and second-order mechanisms for oxidative addition for bound methyl disulfide in the complex W(CO)$_3$(phen)(MeSSMe), **270**, 488
the synthesis and characterization of α-Re_2Cl_4(depe)$_2$ and α-W_2Cl_4(depe)$_2$, **271**, 49
synthesis of chloro and methyl imido cyclopentadienyl molybdenum and tungsten complexes. X-ray molecular structures of [WCp^*Cl_3(N^{t}Bu)], [$MoCp^*ClMe_2$(N^{t}Bu)] and [WCp^*ClMe_2(N^{t}Bu)], **273**, 270

synthesis and characterization of the tungsten–gallium heterometallic cluster $Na_2[W_3GaO_4(O_2CEt)_8]_2$, **277**, 130
protonation of Fischer-type alkylidyne carbonyltungsten complexes. Structural comparison of alkylidyne and alkylidene metal complexes, including a neutron diffraction study of $[W(CHCH_3)Cl_2(CO)(PMe_3)_2]$, **279**, 7
the synthesis, molecular structure and fluxional properties of the 1-phenylpropyne complex $[WI_2(CO)\{Ph_2P(CH_2)PPh_2\}(\eta^2\text{-}MeC_2Ph)]$, **279**, 65
reactivities of the coordinated organonitriles in molybdenum(0) and tungsten(0) phosphine complexes: protonation of the nitrile carbon and cleavage of the C≡N triple bond, **280**, 163
synthesis of double incomplete-cubane clusters $[PPh_4]_2[(\eta^5\text{-}C_5Me_5)MS_3Cu_3X_3]_2$ (M = Mo, W; X = Cl, Br) and reactions with PPh_3, dppm (1,2-bis(diphenylphosphino)methane), and $AsPh_3$, **283**, 136

Two-dimensional complexes
$(C_5H_7N_2O)Cu_3Br_4$: template synthesis of a two-dimensional copper(I) bromide lattice, **279**, 1

Tyrosine derivatives
structures and properties of ternary copper(II) complexes with 4,5-dihydroxyphenylalanine or 2,4,5-trihydroxyphenylalanine and aromatic diamines, **283**, 260

Tyrosine radical
characterization of the active site of galactose oxidase and its active site mutational variants Y495F/H/K and W290H by circular dichroism spectroscopy, **275–276**, 175

Uranium complexes
synthetic, spectroscopic, and X-ray crystallographic studies on the uranyl(VI) complexes of the β-diketophenol ligands 1-(2-hydroxyphenyl)-1,3-butanedione and 1-(2-hydroxyphenyl)-3-phenyl-1,3-propanedione, **277**, 37
chloride ion transfer during the U(IV)/U(III) reduction of UCl_4 in tetrahydrofuran (THF), studied by cyclic voltammetry; synthesis and molecular structure of $[NBu_4][buCl_5(THF)]$, **279**, 210

Uranyl complexes
synthesis and characterization of uranyl thiolate complexes. Crystal and molecular structures of $[Me_4N][UO_2(NO_3)_2(C_5H_4NS)]$ and $[C_{16}H_{25}N_2S_2Si_2][UO_2(NO_3)_2(C_8H_{12}NSSi)]$, **268**, 163
electron spectroscopy for chemical analysis investigation of the interaction of uranyl and calcium ions with humic acids, **273**, 234

Urease model complexes
synthesis, structure and characterization of dinuclear nickel(II) complexes containing the derivatives of *N,N,N′,N′*-tetrakis[(2-benzimidazolyl)methyl]-2-hydroxy-1,3-diaminopropane and an acetate or a propionate ion, **283**, 118

Vanadate complexes
solvent effects on ^{51}V NMR chemical shifts: characterization of vanadate and peroxovanadate complexes in mixed water/acetonitrile solvent, **283**, 37

Vanadium-51 NMR
solvation, preferential solvation and complexation by the solvent of peroxovanadium complexes studied by ^{51}V NMR spectroscopy. Correlations with the oxidative reactivity, **272**, 62
solvent effects on ^{51}V NMR chemical shifts: characterization of vanadate and peroxovanadate complexes in mixed water/acetonitrile solvent, **283**, 37

Vanadium chloroperoxidase
X-ray structures of apo and tungstate derivatives of vanadium chloroperoxidase from the fungus *Curvularia inaequalis*, **273**, 160

Vanadium complexes
synthesis and characterization of new hydrotri(3,5-dimethyl-1-pyrazolyl)borate vanadium halfsandwich complexes, **267**, 19
the preparation and synthetic potential of chlorovanadium(V and IV) complexes supported by enamines and bis(enamines), **267**, 229
hydrothermal synthesis and structural characterization of a three-dimensional organically-templated vanadium phosphate $[H_3NCH_2CH_2NH_3][(VO)_3(H_2O)_2(PO_4)_2(HPO_4)]$, **268**, 257
highly reduced organometallics. 42. A new method for the syntheses of $[V(CO)_6]^-$ and $[V(PF_3)_6]^-$ involving anthracenide mediated reductions of $VCl_3(THF)_3$, **269**, 58
oxidations in the H_2O_2/V^V/AcOH system. Evidence of vanadium(V) complexes with superoxide anion and singlet dioxygen molecule as ligands, **275–276**, 453
synthesis and characterization of two novel monooxovanadium(V) complexes with bidentate benzohydroxamate ligand, **282**, 149

Vanadium compounds
a new form of decavanadate: hydrothermal synthesis and structural characterization of $Ba_{5.33}(H_3O)_{0.33}[V_{10}O_{30}]$, **277**, 263

Vanadyl complexes
insulin-mimetic vanadyl–dithiocarbamate complexes, **283**, 175

Variable pressure
fast water exchange on a Pt(II) center: variable temperature and pressure ^{17}O NMR study at 14.1 tesla of the ionic arylplatinum species $[Pt\{C_6H_3(CH_2NMe_2)_2\text{-}2,6\}(OH_2)]^+(OSO_2CF_3)$, **269**, 322

Vinylidene complexes
concomitant coordination of vinylidene and carbene ligands made possible at ruthenium(II), **272**, 1
study of the relative π-acceptor ability of unsaturated ligands $:C(=C)_nPh_2$ ($n = 0, 1, 2$ and 4) based on cyclic voltammetry of complexes *trans*-$[RhCl(L)(P^iPr_3)_2]$ (L = $:C(=C)_nPh_2$, C_2H_4 and CO), **275–276**, 435

Voltammetry
the 1,8-bis(2′-pyridyl)-3,6-dithiaoctane complex of nickel(II): X-ray crystal structure and borohydride adduct formation, **278**, 217
chloride ion transfer during the U(IV)/U(III) reduction of UCl_4 in tetrahydrofuran (THF), studied by cyclic voltammetry; synthesis and molecular structure of $[NBu_4][buCl_5(THF)]$, **279**, 210

Water soluble bisviologen complexes
preparation and characterization of water-soluble viologen-linked zinc porphyrin and bisviologen-linked zinc porphyrin, **267**, 257

Water-soluble complexes
new non-ionic water-soluble porphyrins: evaluation of manganese(III) polyhydroxylamide porphyrins as MRI contrast agents, **275–276**, 106

Water-soluble porphyrin complexes
sulfonated and acetamidosulfonylated tetraarylporphyrins as biomimetic oxidation catalysts under aqueous conditions, **272**, 228

Work function
electronic properties of some inorganic solids, **270**, 252

XANES
X-ray absorption spectroscopic studies of the binding of ligands to FeMoco of nitrogenase from *Klebsiella pneumoniae*, **275–276**, 150

XANES–EXAFS spectroscopy
tuning up the T_C in Mn(II)Cu(II) bimetallic planes and design of molecular-based magnets, **278**, 159

X-ray analysis
preparation and molecular structure of (*N*-benzyl-1,2-didehydroglycinato) bis(1,10-phenanthroline)ruthenium(II) chloride: [Ru(*N*-Bn-gly-H_2)(phen)$_2$]Cl, **267**, 329
chain-like crystal structure and magnetic properties of [catena-bis(1,2-diaminoethane)copper(II)-μ-dicyano-argentate]-dicyanoargentate, **281**, 134

X-ray crystallography
synthesis and structural characterization of the [2,9-dimethyl-4,7-diaza-2,9-decanedithiolato]oxorhenium(V) complex, **268**, 169

X-ray crystal structures

structures and properties of ternary copper(II) complexes with 4,5-dihydroxyphenylalanine or 2,4,5-trihydroxyphenylalanine and aromatic diamines, **283**, 260

X-ray photoelectron spectroscopy
preparation and characterization of twin-cubane-type clusters with molybdenum–copper–sulfur cores: X-ray structure determination of $[\{(NH_3)_6(H_2O)(OH)Mo_3CuS_4(O)\}_2]Cl_4\cdot 8H_2O$, **282**, 50

X-ray structure
NMR spectra of dinuclear manganese and iron compounds containing 1,4,7-triazacyclononane and 1,4,7-trimethyl-1,4,7-triazacyclononane, **268**, 43

o-Xylylene complexes
mononuclear and binuclear *o*-xylylene derivatives of Pd(II), **269**, 191

Ytterbium complexes
C_{60}-fullerenides of ytterbium and lutetium, **280**, 138

Yttrium complexes
syntheses and structure of dimeric yttrium tris[bis(trimethylsilyl)-phosphanide]; X-ray structures of $[(thf)LiCl\cdot(thf)_2LiP^tBu_2]_2$ and $Y[CH(SiMe_3)_2]_3$, **269**, 91
coordination of a disubstituted alkene in a chelated d^0-metal–alkyl–alkene complex. Evidence for equilibrium between complexes with coordinated and free alkenes, **270**, 5
potentiometric and computer studies of yttrium–EDTMP, **275–276**, 37

Zeolite-Y
photoreduction of methyl-viologen by zeolite-Y stabilized silver clusters, **269**, 253

Zero-dimensional complexes
X-ray diffraction and EPR studies of binuclear complex $[CuCl(C_6H_9N_3O_3)_2\cdot H_2O]_2Cl_2$ crystals, **269**, 326

Ziegler–Natta polymerization
preparation of doubly-silylene-bridged zirconocene alkyl complexes, $(Me_2Si)_2\{\eta^5\text{-}C_5H_2\text{-}4\text{-}CHMe_2\}\{\eta^5\text{-}C_5H\text{-}3,5\text{-}(CHMe_2)_2\}ZrR_2$ ($R = CH_3$, CH_2Ph) and investigations of their activity in 1-pentene polymerization. Molecular structure of $(Me_2Si)_2\{\eta^5\text{-}C_5H_2\text{-}4\text{-}CHMe_2\}\{\eta^5\text{-}C_5H\text{-}3,5\text{-}(CHMe_2)_2\}Zr(CH_2Ph)_2$, **280**, 226

Zinc complexes
coordination compounds of the rigid rod ligand 1-(2-pyridyl)-2-(2-thienyl)ethyne (pythio). Single crystal X-ray structures of $[Cu(pythio)_2(NO_3)_2]$ and $[Zn(pythio)_2Cl_2]$, **267**, 193
complexes of the terdentate N-donor ligand 6-(2-aminophenyl)-2,2′-bipyridine (L): crystal structures of mononuclear $[ZnL_2][PF_6]_2$ and tetranuclear $[\{CuL(MeCN)\}_4(\mu^4\text{-}PO_4)][PF_6]_5$ containing an unusual μ^4-bridging phosphate ion, **267**, 239
preparation and characterization of water-soluble viologen-linked zinc porphyrin and bisviologen-linked zinc porphyrin, **267**, 257
synthesis, spectral and structural characterization of the zinc carboxylate $[Zn(2\text{-quinolinecarboxylato})_2(1\text{-methylimidazole})_2]$, **269**, 283
a new macrocyclic tetraamine, 2,4-dinitrophenylcyclen (=1-(2,4-dinitrophenyl)-1,4,7,10-tetraazacyclododecane): synthesis, cation reporter properties, and recognition of 1-methylthymine by its zinc(II) complex, **270**, 424
dinuclear metal complexes of Cd(II), Zn(II) and Fe(II) with triple-helical structure and predetermined chirality, **271**, 36
solution behaviour and zinc complexation of tripeptides with cysteine and/or histidine at both termini, **272**, 115
the synthesis and spectroscopic characterization of a chiral zinc(II) porphyrin dimer containing an amino acid bridge: model complexes for the induced circular dichroism of porphyrin protein supramolecules, **272**, 295
synthesis and coordinating properties of the functionalized macrocyclic ligand 1,4-bis(1-methylimidazol-2-ylmethyl)-7-carboxymethyl-1,4,7-triazacyclononane. Crystal structures of cobalt(III), nickel(II) and zinc(II) complexes, **273**, 151
a reversible Cu^{II}/Cu^{I} electron transfer centre based on a flexible di-imidazole ligand, **273**, 225
syntheses of ruthenium hydridotris(1-pyrazoryl)borate complexes having sulfur-donor coligands, **273**, 238
synthesis, structure characterization and magnetic properties of tellurostannates $[M(en)_3]_2(Sn_2Te_6)$ (M = Mn, Zn), **273**, 310
crystal structures of aryloxide complexes of zinc embracing sodium aryl group interactions: $Na[Zn(2,6\text{-diphenylphenoxide})_3(H_2O)]$ and $Na[Zn_2(2,6\text{-diisopropylphenoxide})_4Cl]\cdot 3THF$, **274**, 115
some novel carbamato chemistry of cadmium and zinc, **274**, 239
a ternary complex of carbonic anhydrase: X-ray crystallographic structure of the adduct of human carbonic anhydrase II with the activator phenylalanine and the inhibitor azide, **275–276**, 295
dynamic properties of bovine Cu,Zn superoxide dismutase from crystallographic data, **275–276**, 349
zinc as a structural and folding element of proteins which interact with DNA, **275–276**, 481
five-coordinate Salen(tBu) complexes of zinc, **277**, 157
photoinduced processes in 4′-(9-anthryl)-2,2′:6′,2″-terpyridine, its protonated forms and Zn(II), Ru(II) and Os(II) complexes, **277**, 225
preparation and some properties of perfluoroalkoxy-substituted phthalocyanine complexes of iron(III), nickel(II) and zinc(II), **278**, 253
molecular structure of a five-coordinate binuclear zinc complex, **279**, 249
synthesis, structure and characterization of zinc(II), copper(II), zinc(II)barium(II) and copper(II)barium(II) complexes of macrocyclic heteronucleating ligands based on isothiosemicarbazide, **282**, 61
zinc thiolate complexes with chelating nitrogen ligands, **282**, 193

Zinc peptides
Boc-Glu-Thr-Ile-His-OMe/Zn^{2+} in terms of the zinc-binding sites of proteases, **283**, 243

Zinc proteases
Boc-Glu-Thr-Ile-His-OMe/Zn^{2+} in terms of the zinc-binding sites of proteases, **283**, 243

Zinc site
structural characterization of the zinc site in *Escherichia coli* L-threonine dehydrogenase using extended X-ray absorption fine structure spectroscopy, **275–276**, 215

Zirconium-catalyzed
zirconium-catalyzed and zirconium-promoted cyclization reactions of non-conjugated dienes with alkylmagnesium halides to give cycloalkylmethylmagnesium derivatives, **280**, 8

Zirconium complexes
eight-coordinate chelate complexes of zirconium(IV) and niobium(IV): X-ray diffractometric and EPR investigations, **270**, 177
preparation of trimethylsilyl alkyne complexes of bis(pentamethylcyclopentadienyl)zirconium, $(\eta^5\text{-}C_5Me_5)_2Zr(RC{\equiv}CSiMe_3)$, and their regioselective reactions with nitrous oxide, **270**, 399
synthesis, characterization, and reactivity of organometallic Zr(IV) carboxylate complexes, **270**, 527
bis(η^3-allyl)(η^8-pentalene)zirconium: preparation, structure and structural dynamics, **270**, 555
trimethylphosphine halide complexes of titanium, zirconium and hafnium in oxidation states IV, III and II. Crystal structure of $MI_4(PMe_3)_n$ (M = Zr, $n = 3$; M = Hf, $n = 2.5$), $M_2X_6(PMe_3)_4$ (M = Hf, Ti, X = I; M = Zr, X = Br) and $(\mu\text{-}\eta^6,\eta^6\text{-}C_6H_6)\text{-}Hf_2I_4(PMe_3)_4\cdot C_6H_6$, **271**, 180

Zirconocene complexes
zirconocene–alumoxane (1:1) – a catalyst for the selective dimerization of α-olefins, **270**, 20

misdirected π-donor ligands: $(\eta^5\text{-}C_5H_5)_2Zr(Cl)(SR)$ with sterically more demanding R groups have lower rotational barriers, **270**, 227
the synthesis of substituted bis[(diarylphosphinomethyl)cyclopentadienyl]zirconocene dichloride complexes for the preparation of heterodimetallic complexes containing early/late transition metal combinations, **270**, 446
preparation of doubly-silylene-bridged zirconocene alkyl complexes, $(Me_2Si)_2\{\eta^5\text{-}C_5H_2\text{-}4\text{-}CHMe_2\}\{\eta^5\text{-}C_5H\text{-}3,5\text{-}(CHMe_2)_2\}ZrR_2$ ($R = CH_3$, CH_2Ph) and investigations of their activity in 1-pentene polymerization. Molecular structure of $(Me_2Si)_2\{\eta^5\text{-}C_5H_2\text{-}4\text{-}CHMe_2\}\{\eta^5\text{-}C_5H\text{-}3,5\text{-}(CHMe_2)_2\}Zr(CH_2Ph)_2$, **280**, 226
on the reduction behaviour of 2-methoxyethyl and 2-methylthioethyl substituted zirconocene dichlorides: crystal structure of $(\eta^5,\sigma\text{-}C_5Me_4CH_2CH_2S)_2Zr$, **280**, 257
Zirconocenes
alkyl exchange between aluminum trialkyls and zirconocene dichloride complexes — a measure of electron densities at the Zr center, **270**, 376
Zwitterionic complexes
reactions of $Ti(O\text{–}i\text{-}Pr)Cl_3$ with 2-aminophenol and the crystal structure of the zwitterionic complex $[Ti(O\text{–}i\text{-}Pr)Cl_3(THF)(^-OC_6H_4NH_3^+)]\cdot THF$, **278**, 232

ELSEVIER

www.elsevier.nl/locate/ica

Inorganica Chimica Acta 296 (1999) 281–290

Inorganica Chimica Acta

Cumulative Author Index of Volumes 284–296

Abbott, B., **289**, 198
Abedin, Z., **285**, 97
Abruña, H.D., **288**, 189
Adeyemo, A.O., **290**, 145
Aeby, A., **296**, 45
Aguilar-Cornejo, A.F., **285**, 142
Ainscough, E.W., **290**, 228
Akbar Ali, M., **287**, 8
Akkawi, M., **286**, 229
Albano, V.G., **285**, 70; **291**, 372
Albertini, R., **286**, 134
Albert, K., **286**, 30
Albinati, A., **284**, 191; **296**, 246
Albrecht, M., **287**, 204
Alderighi, L., **285**, 39
Aldridge, S., **289**, 85
Alexander, J.J., **288**, 159; **290**, 159
Aliffi, A., **286**, 121
Ali, S., **296**, 26
Al-Jeboori, M.J., **285**, 76
Almeida, V.R., **291**, 212
Amadei, G.A., **288**, 40; **293**, 245
Amani, S., **286**, 24
Amezcua, C.A., **290**, 80
Amorium, M.T., **284**, 20
Anderson, O.P., **289**, 76
Anderson, R.J., **284**, 273
Andreasen, L.V., **295**, 153
Angelici, R.J., **286**, 207
Angus, P.M., **287**, 1
An, J., **287**, 82
Anjali, K.S., **295**, 9
Antunes, O.A.C., **292**, 1
Aoki, K., **290**, 145; **295**, 71
Araki, K., **285**, 197
Arce, A.J., **285**, 277
Arena, C.G., **292**, 84
Arif, A.M., **287**, 109
Arisawa, M., **296**, 67
Arrieta, J.M., **295**, 106
Arulsamy, N., **284**, 158
Atkinson, A., **285**, 60
Auer, E., **287**, 167
Auffrant, A., **296**, 139
Ault, B.S., **286**, 1
Averill, B.A., **285**, 152
Azuma, N., **288**, 90
Babich, J.W., **284**, 252
Bacchi, A., **295**, 171
Bachmann, C., **290**, 167
Bae, J.-Y., **289**, 141
Baggio, R., **284**, 49; **286**, 74
Bai, J.-F., **292**, 117
Baik, M.-H., **294**, 140
Bajaj, H.C., **285**, 89
Baker, A.T., **284**, 266
Baker, L.J., **292**, 260
Bakos, J., **296**, 222
Balazs, G., **287**, 61
Baldacchini, T., **295**, 200
Baldoli, C., **296**, 236
Bandoli, G., **294**, 114
Baraldo, L.M., **285**, 129
Barclay, J.E., **291**, 101
Barranco, E.M., **291**, 60
Barton, L., **285**, 290; **289**, 95; **289**, 134
Basallote, M.G., **287**, 134
Basato, M., **285**, 18; **290**, 14
Batista, A.A., **290**, 207
Batten, S.R., **292**, 231
Baumann, F., **285**, 129
Bauminger, E.R., **286**, 229
Baxter, J.T., **296**, 52
Bayón, J.C., **295**, 64
Beall, H., **289**, 1
Beattie, J.K., **294**, 99
Beauchamp, A.L., **295**, 209
Beckel, C.L., **289**, 198
Beck, K., **288**, 159
Beck, W., **290**, 237
Beguin, C., **291**, 148
Belay, Y.F., **285**, 318
Belicchi Ferrari, M., **286**, 134
Bellacchio, E., **286**, 121
Bellachioma, G., **288**, 200
Bell, K.E., **290**, 80
Belsky, V.K., **284**, 180
Belyansky, M., **289**, 191
Benetollo, F., **291**, 39
Benito, M., **291**, 247
Bérézovsky, F., **284**, 8
Bergter, R., **285**, 1
Bermejo, E., **294**, 79
Bermejo, M.R., **293**, 210
Bernal, I., **292**, 149; **295**, 91
Bernal, M., **295**, 39
Bevilacqua, V.L.H., **285**, 217
Bian, G., **285**, 332
Bian, J., **287**, 82
Bianchi, A., **288**, 244
Bill, E., **291**, 258
Billodeaux, D.R., **295**, 91
Birdsall, D.J., **290**, 1
Bird, S.J., **295**, 56
Bismondo, A., **286**, 103
Blackman, A.G., **294**, 14
Blauer, G., **286**, 229
Blomgren, P.A., **296**, 164
Blower, P.J., **294**, 170
Bobrowski, A., **293**, 223
Boersma, J., **291**, 326
Boever, C., **284**, 284
Böhm, M.C., **291**, 82
Bollinger, J.C., **292**, 220
Bombieri, G., **291**, 39
Bonavia, G.H., **284**, 252
Bond, A.M., **291**, 21; **293**, 223
Bondarev, O.G., **295**, 164
Bonilla, B.M., **292**, 137
Bonnington, L.S., **290**, 213
Bordignon Luiz, M.T., **287**, 134
Bordoni, S., **291**, 333
Boubekeur, K., **284**, 139
Bould, J., **285**, 290; **289**, 95
Bouwman, E., **287**, 105
Bouziotis, P., **290**, 247
Boyajian, Y.D., **285**, 134
Brandts, J.A.M., **291**, 326
Bray, P.J., **289**, 158
Bressan, M., **288**, 122
Breuer, J., **291**, 438
Brianese, N., **293**, 178
Bridgeman, A.J., **288**, 17
Brinn, I.M., **292**, 1
Brito, F., **291**, 158
Brodie, A.M., **290**, 228
Brook, M.A., **296**, 208
Browning, J., **288**, 181; **292**, 73; **292**, 198
Brownless, N.J., **287**, 89
Brown, T.M., **288**, 112
Brozik, J.A., **294**, 275
Bruckmann, J., **288**, 118
Brüdgam, I., **291**, 266
Bruenger, F.W., **287**, 109
Brüggeller, P., **290**, 167
Brunet, J.-J., **291**, 300
Bruno, G., **292**, 84
Bryan, J.C., **287**, 8; **290**, 86
Bu, W.-M., **287**, 52
Bucher, U.E., **284**, 191
Buckingham, D.A., **294**, 14
Bullita, E., **287**, 117
Bünzli, J.-C.G., **293**, 53
Burckhardt, U., **296**, 183
Buretea, M., **287**, 152
Burriel, R., **291**, 279
Buschmann, W.E., **284**, 175
Busetto, L., **291**, 333
Butcher, R.J., **287**, 8
Butenschön, H., **296**, 150
Byriel, K.A., **290**, 133
Cabaleiro, S., **294**, 87
Calabi, L., **288**, 244
Calamai, P., **285**, 309
Calatayud, M.L., **287**, 173
Calligaris, M., **285**, 116; **292**, 144
Cambi, R., **288**, 200
Campazzi, E., **286**, 46
Camps, X., **296**, 94
Camus, A., **290**, 180
Candini, M., **292**, 189
Canovese, L., **293**, 44
Canty, A.J., **287**, 27
Capacchi, S., **286**, 134
Capparelli, M.V., **285**, 277
Carballo, R., **295**, 106
Carcelli, M., **285**, 138; **292**, 121; **295**, 171
Cardaci, G., **288**, 200
Carlson-Day, K.M., **284**, 300
Carpenter, J.D., **286**, 1
Carrano, C.J., **286**, 80
Carty, A.J., **291**, 178
Caruso, U., **292**, 163
Carvalho, C.C., **292**, 249
Carvalho, C.E.M., **292**, 1
Casellato, U., **287**, 117; **293**, 178
Cassani, M.C., **291**, 333
Castellano, E.E., **285**, 142
Castellari, C., **291**, 372
Castiñeiras, A., **294**, 47, 79; **295**, 106
Castro, I., **287**, 173
Castro, J., **294**, 87
Castro, M., **291**, 279
Catalano, V.J., **286**, 181
Cattabriga, M., **286**, 46
Caubel, Y., **288**, 233
Caulton, K.G., **291**, 226

Elsevier Science S.A.
PII: S0020-1693(99)00485-5

Cavaco, I., **293**, 1
Cecconi, F., **285**, 39
Cervera, B., **288**, 57
Cevik, S., **292**, 112
Cha, J., **288**, 189
Cha, S.K., **288**, 189
Chadwick, A.V., **294**, 170
Chan, W.H., **290**, 28
Chandrasekhar, V., **286**, 127
Chanon, M., **293**, 80
Chantrapromma, S., **292**, 57
Chatterjee, D., **285**, 170
Chauvin, R., **291**, 300
Che, C.-M., **292**, 117
Chen, C., **286**, 7; **293**, 95
Chen, D., **293**, 206
Chen, H., **285**, 55
Chen, J.-D., **288**, 82; **292**, 182
Chen, J.-T., **285**, 241; **293**, 89; **293**, 129
Chen, X., **292**, 157
Chen, X., **285**, 326
Chen, Y., **285**, 157; **293**, 123
Chen, Z.-D., **287**, 82
Cheng, B., **291**, 49
Chesnut, D.J., **292**, 41
Chessa, G., **293**, 44
Cheung, K.-K., **284**, 205; **285**, 223
Chi, T.-Y., **285**, 107
Chiffre, J., **291**, 300
Chim, J.L.C., **290**, 28
Chinea, E., **285**, 39
Chin, M., **292**, 254
Ching, S., **294**, 123
Chinnakali, K., **292**, 57, 117; **293**, 106
Chiotellis, E., **285**, 97; **287**, 142; **290**, 247; **295**, 1
Chiumia, G.C., **285**, 297
Cho, S.-I., **289**, 141
Choi, K.-Y., **293**, 100
Choppin, G.R., **293**, 167
Chou, S.-Y., **286**, 155
Choudhary, N.F., **293**, 147
Chung, C.-S., **285**, 107
Chung, W.P., **291**, 388
Chung, Y.-H., **293**, 30
Chung, Y.K., **286**, 215; **295**, 214
Cindrić, M., **284**, 223
Ciringh, Y., **294**, 133
Clark, C.R., **294**, 14
Clark, J.R., **288**, 35
Clark, R.W., **293**, 61
Claver, C., **295**, 64
Clegg, W., **289**, 95
Colacio, E., **284**, 292
Colbran, S.B., **284**, 76
Coll, R.K., **290**, 213
Commenges, G., **285**, 49
Comuzzi, C., **286**, 103
Connac, F., **295**, 209
Consiglio, G., **296**, 45
Cooke, P.A., **285**, 290
Cooksey, C.J., **288**, 112
Copper, C.L., **288**, 229
Cornman, C.R., **285**, 134
Correia, I., **293**, 1
Costes, J.-P., **285**, 49; **294**, 8
Cotton, F.A., **284**, 304; **285**, 149; **287**, 159; **288**, 69
Coulais, Y., **295**, 209
Cowie, J., **289**, 125
Cox, B.G., **285**, 1; **286**, 160
Cozzini, P., **285**, 138
Craig, D.C., **284**, 266; **285**, 297
Crawford, M.-J., **294**, 68
Csákai, Z., **286**, 93
Cserépi-Szűcs, S., **296**, 222
Cui, C.-P., **293**, 89
Cui, Y., **293**, 129
Cullison, S., **292**, 220
Cunningham, D., **291**, 252
Cupertino, D., **290**, 1
Czarnecki, K., **287**, 21
Czernuscewicz, R.S., **286**, 80

Dahan, F., **294**, 8
Daiguebonne, C., **284**, 139
Dai, J., **285**, 332
Dai, W.-N., **290**, 127
Dakternieks, D., **285**, 39
Dal, A., **287**, 167, 199
d'Alessandro, N., **288**, 122
Dalley, N.K., **284**, 85
Daniels, L.M., **285**, 149; **287**, 159; **288**, 69
Danil de Namor, A.F., **285**, 142
Danzeisen, O.F., **287**, 218
Dartiguenave, M., **295**, 209
Das, A., **285**, 89
DasGupta, B., **292**, 172
da Silva, A.S., **292**, 1
Davankov, V.A., **295**, 164
David, M.C., **292**, 137
Davidović, N., **294**, 35
Davies, P.J., **294**, 103
Davies, S.C., **291**, 101
Davis, C.E., **292**, 220
Díaz, C., **294**, 20
de Brito, M.A., **290**, 207
Decréau, R., **293**, 80
Decurtins, S., **286**, 108
Dedieu, A., **287**, 27
Deeming, A.J., **285**, 277
Delgado, E., **284**, 14
del Río, I., **287**, 113
Delville, M.-H., **291**, 1
Deng, Y., **286**, 7; **293**, 95
Dennison, K.J., **285**, 217
Derwahl, A., **285**, 313
des Abbayes, H., **291**, 166
De Sanctis, Y., **285**, 277
De Silva, C., **287**, 21
de Silva, M.A.A., **292**, 1
Desurmont, G., **290**, 159
Detomi, N., **285**, 18
Devereux, M., **293**, 160
Dewan, J.C., **291**, 388
Diaz, C., **286**, 169
Di Bernardo, P., **286**, 103
Dickman, M.H., **288**, 40
Diéguez, M., **295**, 64
Dietrich, J., **288**, 206
Dikarev, E.V., **284**, 304
Dilworth, J.R., **285**, 76
Dimmock, P., **288**, 40
Dixon, D.A., **290**, 197; **294**, 193
Doedens, R.J., **292**, 112
Doherty, S., **291**, 178
Dolmella, A., **294**, 114
Domasevich, K.V., **284**, 85
Domasevitch, K.V., **293**, 234
Domínguez, S., **285**, 39; **291**, 158; **294**, 47
Donaldson, W.A., **296**, 261
Dong, G., **284**, 266
Donnadieu, B., **291**, 300
Donohoe, D.J., **289**, 125
Don Wang, J., **294**, 214
Dörfler, U., **285**, 290
Drago, D., **292**, 84
Drake, J.E., **287**, 61
Drew, M.G.B., **285**, 31
Drexler, H.-J., **285**, 305
Driessen, W.L., **288**, 239
Drillon, M., **294**, 8
Drommi, D., **296**, 183
Dronsfield, A.T., **288**, 112
Duan, C., **294**, 1; **295**, 48, 229
Duarte, M.T., **293**, 1
Dubey, S.N., **293**, 61
Duda, M., **291**, 289
du Preez, J.G.H., **294**, 114
Dupuis, A., **285**, 49
Durán, M.L., **284**, 133; **294**, 79; **295**, 39
Duthie, A., **285**, 39

Earley, J.E., **288**, 40; **293**, 245
Easton Lawrence, K., **296**, 52
Ebihara, M., **292**, 244
Echegoyen, L., **292**, 34
Edwards, C., **294**, 153
Edwards, D.A., **287**, 89
Edwards, R.A., **285**, 313
Eglin, J.L., **284**, 300
Eilmes, J., **290**, 14
El-Ahl, A.-A.S., **296**, 261
El Kasmi, A., **292**, 204
Ellermann, J., **284**, 288
Ellis, S.R., **293**, 223
Englert, U., **289**, 58
Enright, G.D., **291**, 178
Ernst, K.A., **296**, 170
Ernst, R.D., **288**, 154; **296**, 170
Escuer, A., **286**, 62, 189
Esparza, P., **291**, 158
Evans, C., **294**, 183
Evans, D.J., **284**, 296; **291**, 39; **291**, 101

Facchinetti, A., **290**, 180
Fairlie, D.P., **290**, 133
Fanwick, P.E., **288**, 35

Faraone, F., **292**, 84
Farmer, J.M., **285**, 318; **292**, 137
Fatin-Rouge, N., **293**, 53
Faus, J., **288**, 57
Fawcett, J., **295**, 146
Fawzi, R., **296**, 103
Fehlhammer, W.P., **291**, 266
Fehlner, T.P., **289**, 85
Femoni, C., **291**, 372
Fenton, R.R., **288**, 29
Ferencic, M., **286**, 114
Férey, G., **291**, 317
Ferlay, S., **286**, 108
Fernando, Q., **295**, 115, 125
Ferrara, M.L., **285**, 70
Fettinger, J.C., **286**, 67
Fiedler, J., **285**, 129; **291**, 66
Finke, R.G., **291**, 426
Fiore, C., **285**, 249
Fischer, J., **292**, 96
Flett, J.I., **290**, 213
Flörke, U., **287**, 167, 199
Flor, M.T., **295**, 64
Fogagnolo, M., **286**, 46
Fondo, M., **293**, 210
Font-Bardía, M., **286**, 62, 169, 189
Foresti, M., **288**, 244
Fossheim, R., **284**, 167
Fowler, J.H., **288**, 112
Foxman, B.M., **288**, 215
Francese, G., **286**, 108
Frank, W., **296**, 127
Fraústo da Silva, J.J.R., **291**, 39
Freitas Nunes, C.C., **292**, 249
Friebe, M., **295**, 1
Fronczek, F.R., **292**, 149; **295**, 91
Frühauf, H.-W., **291**, 438
Fu, D.-G., **287**, 52
Fu, Y.-J., **290**, 127
Fuess, H., **287**, 181
Fujihara, T., **295**, 84
Fukuda, Y., **287**, 95; **292**, 7
Fukuoka, A., **294**, 266, 281
Fukushima, Y., **294**, 281
Fukuta, Y., **296**, 72
Fun, H.-K., **292**, 57, 117; **293**, 106; **294**, 1
Furubayashi, Y., **295**, 234

Gaines, D.F., **289**, 1
Galanski, M., **292**, 127
Gałdecka, E., **293**, 110
Gałdecki, Z., **293**, 110
Gang, L.-L., **290**, 51
Ganguly, D., **288**, 189
García-Deibe, A., **293**, 210
García, G., **284**, 296
García, J., **287**, 37
García-Vázquez, J.A., **284**, 133; **294**, 87; **295**, 39
Garcia Diaz, J., **287**, 186
Garcia Gomez, G., **296**, 94
Gardberg, A., **294**, 153
Garland, M.T., **284**, 49; **286**, 74
Garst, J.F., **296**, 52
Gaspar, M., **284**, 20
Gasque, L., **288**, 106
Gau, H.-M., **285**, 254; **293**, 12
Gavrilov, K.N., **295**, 164
Gemel, C., **286**, 114
Genge, A.R.J., **288**, 142
Geraghty, M., **293**, 160
Gerasimchuk, N.N., **284**, 85
Gerasimchuk, O.A., **284**, 85
Gérault, Y., **284**, 139
Gerber, T.I.A., **294**, 114
Geremia, S., **292**, 144
Germain, G., **295**, 106
Ghedini, M., **292**, 163
Giannandrea, R., **284**, 116
Giester, G., **292**, 127
Gili, P., **295**, 106
Gillard, R.D., **293**, 1
Gimeno, M.C., **291**, 60
Gipson, S.L., **285**, 318
Girerd, J.-J., **290**, 139
Gislason, J., **284**, 284
Gleizes, A., **288**, 233
Goanta, M., **287**, 218
Godfrey, S.M., **292**, 213
Gómez, C., **295**, 39
Gong, Y., **287**, 14
González, A.M., **293**, 210
González-Platas, J., **284**, 258
Goodwin, N.J., **295**, 18
Gopal Dongol, K., **296**, 150
Görls, H., **286**, 14
Gormal, C.A., **291**, 212
Gorrichon, J.-P., **288**, 233
Gossage, R.A., **287**, 113
Goto, M., **294**, 56
Gou, S., **287**, 14, 95; **294**, 1
Graiff, C., **296**, 236
Gramlich, V., **284**, 191; **291**, 355
Granifo, J., **295**, 56
Grätzel, M., **296**, 250
Gray, E.J., **290**, 213
Griesar, K., **287**, 181
Grijalva, H., **295**, 125
Grohmann, A., **286**, 98
Grönberg, K.L.C., **291**, 212
Grotjahn, M., **285**, 305
Guerchais, V., **291**, 449
Guerri, A., **285**, 309
Guilloa, O., **284**, 139
Guo, D., **292**, 157
Gurrieri, S., **286**, 121
Gust, R., **285**, 184
Gutiérrez-Zorrilla, J.M., **290**, 57
Guzei, I.A., **286**, 207

Haar, C.M., **291**, 32
Haase, W., **287**, 181; **291**, 82
Haasnoot, J.G., **286**, 24
Haefner, S.C., **285**, 149; **287**, 159; **288**, 69
Hagback, P.H., **284**, 273
Hagihara, Y., **292**, 64
Hahn, F., **287**, 101
Hahn, F.E., **288**, 47
Haid, R., **290**, 167
Haiduc, I., **287**, 61; **290**, 256
Hajem, A.A., **284**, 8
Hamor, T.A., **285**, 55
Hampel, F., **291**, 91
Han, J., **294**, 95
Hao, Z., **284**, 112
Harada, M., **294**, 119
Harada, Y., **285**, 336
Harnisch, J.A., **286**, 207
Hartmann, S., **287**, 199
Hasebe, K., **296**, 19
Hashimoto, M., **284**, 237
Hasić, J., **285**, 208
Hasselgren, C., **292**, 266
Hattersley, A.D., **289**, 149
Haupt, H.-J., **287**, 167, 199
Haushalter, R.C., **288**, 1; **292**, 41
Hay, B.P., **290**, 86
Hayashi, H., **296**, 254
Hayashi, K., **294**, 56
Hayashi, T., **296**, 19
Hayashi, Y., **291**, 426
He, L., **287**, 14
He, Q., **295**, 244
He, X., **285**, 326
Heck, R.A., **286**, 181
Hegedüs, C., **296**, 222
Heidari, N., **286**, 160
Heinemann, F.W., **284**, 288; **286**, 98; **288**, 206
Heinrich, H., **285**, 184
Hellemons, J.C., **295**, 189
Henderson, R.A., **285**, 178; **291**, 39; **291**, 212
Henderson, R.K., **287**, 105
Henderson, W., **285**, 145; **290**, 213; **292**, 260; **294**, 183; **295**, 18
Henriques, R.T., **293**, 1
Herber, R.H., **287**, 152; **291**, 74
Heřmánek, S., **289**, 20
Hernández, E., **284**, 14
Hernández-Molina, M., **284**, 258
Herndon, J.W., **296**, 273
Herradora, R., **285**, 269
Herrera, J.O.M., **292**, 1
Hidai, M., **285**, 336
Higes, F.J., **293**, 1
Higgs, T.C., **286**, 80
Hill, M.G., **286**, 181
Hiller, W., **285**, 76
Himeno, S., **284**, 237
Hinton, H.A., **285**, 55
Hirano, M., **291**, 341; **294**, 266
Hitchcock, P.B., **293**, 147
Hiyama, T., **296**, 231
Hodgson, D.J., **284**, 158
Hofmann, M., **289**, 58
Hogarth, G., **291**, 178
Hogerheide, M.P., **291**, 326
Hoggard, P.E., **291**, 136

Holdt, H.-J., **285**, 305
Holloway, G., **292**, 220
Holwerda, R.A., **284**, 124; **288**, 211
Homrighausen, C.A., **290**, 159
Horner, O., **290**, 139
Horwitz, C.P., **294**, 133
Hoshino, Y., **292**, 64
House, D.A., **285**, 313; **288**, 53, 181; **292**, 73; **292**, 198
Housecroft, C.E., **289**, 149
Howell, F.S., **295**, 121
Hriljac, J.A., **294**, 179
Hsu, M.-A., **294**, 232
Hu, H.-M., **293**, 89
Hu, H.-W., **290**, 127; **295**, 48
Hu, N.-H., **290**, 145; **295**, 71
Huang, J.-S., **285**, 241; **293**, 89, 129; **294**, 73
Huang, P., **287**, 52
Huang, S.D., **294**, 95
Huang, T.-Y., **293**, 12
Huang, W., **291**, 380
Huang, X.-Y., **285**, 152; **290**, 121
Huang, Z., **294**, 109
Hübler, K., **287**, 204
Huffman, J.C., **291**, 226; **292**, 220
Hughes, D.L., **291**, 101
Hughes, R.S., **295**, 146
Huguet, S., **291**, 300
Huhmann-Vincent, J., **294**, 240
Hummel, H., **291**, 258
Humphry-Baker, R., **296**, 250
Hunter, A.P., **290**, 189
Hutchison, J.C., **294**, 153
Hutchison, J.E., **293**, 195
Huttner, G., **296**, 222
Hvastijová, M., **287**, 186
Hwu, J.R., **285**, 107

Iacopino, D., **292**, 189
Ianelli, S., **292**, 121
Iapalucci, M.C., **291**, 372
Ichikawa, M., **294**, 281
Ide, K., **285**, 156
Igoshi, T., **286**, 233
Ikeda, Y., **294**, 119
Ikeno, T., **296**, 86
Inada, K., **294**, 56
Inagaki, S., **294**, 281
Inamasu, N., **296**, 231
Inomata, S., **291**, 395
Inomata, Y., **295**, 121
Inoue, M., **295**, 115, 125
Inoue, M.B., **295**, 115, 125
Ishihara, K., **290**, 222
Ishii, H., **296**, 176
Ishikawa, Y., **294**, 56
Israel, T., **292**, 172
Ito, K., **290**, 113
Ito, M., **284**, 149
Itoh, K., **296**, 72
Ivanova, S.M., **289**, 76
Ivanov, S.V., **289**, 76
Iwasawa, K., **285**, 156

Izquierdo, I., **294**, 20
Jaballas, J., **289**, 11
Jackson, W.G., **284**, 37 **287**, 1; **290**, 133
Jacobson, R.A., **284**, 284
Jäger, E.-G., **291**, 190
Jäger, L., **287**, 186
Jagirdar, B.R., **292**, 105
Jagner, S., **292**, 266
Jain, V.K., **285**, 190
James, B.D., **284**, 215; **285**, 122
James, M.A., **284**, 215
Jameson, G.B., **294**, 14
Janas, Z., **285**, 178
Janczak, J., **287**, 101; **288**, 174; **293**, 155
Janiak, C., **295**, 249
Janssen, M.D., **291**, 326
Jaumà Cayuela, A., **296**, 94
Jeffery, J.C., **292**, 231; **294**, 170
Jensen, W.P., **284**, 284
Jeon, H.-J., **286**, 142
Jerzykiewicz, L.B., **285**, 178
Ji, D., **286**, 80
Ji, L., **285**, 326
Ji, W., **294**, 73
Ji, Z.-P., **293**, 89
Jiang, D., **287**, 212
Jiang, L.-J., **295**, 48
Jiang, Q., **290**, 64
Jin, H., **287**, 27
Jin, X.-L., **287**, 82
Johannsen, B., **295**, 1
Johnson, M.D., **293**, 229
Jolk, I., **285**, 262
Jones, C.J., **285**, 55, 283
Jones, F.E., **294**, 207
Joó, F., **293**, 218
Journaux, Y., **288**, 57
Julliard, M., **293**, 80
Julve, M., **284**, 258; **287**, 173; **288**, 57
Jung, S.-J., **286**, 155
Jung, S.-K., **293**, 140
Jung, T., **287**, 167
Jurek, P., **287**, 47
Justin Thomas, K.R., **286**, 127

Kahn, O., **291**, 279
Kaim, W., **285**, 129; **287**, 204; **291**, 66
Kaizaki, S., **292**, 16; **295**, 84
Kajikawa, Y., **288**, 90
Kakiuchi, K., **296**, 37
Kamenar, B., **284**, 223
Kane, J.C., **287**, 209
Kaneko, T., **296**, 195
Kanematsu, N., **292**, 244
Kang, B.-S., **290**, 121
Kang, S.-G., **293**, 140
Kang, S.O., **289**, 141
Kaplan, S.F., **285**, 116
Käpplinger, I., **291**, 190
Karam, A., **285**, 277
Kashino, S., **293**, 20
Katada, M., **291**, 127
Katsukawa, Y., **294**, 255

Katz, C., **292**, 172
Kautz, J.A., **285**, 318; **292**, 137
Kawai, N.T., **285**, 217
Kawamura, T., **292**, 244
Kayran, C., **284**, 229
Kelly, H.C., **290**, 80
Keng, T.-C., **288**, 82; **292**, 182
Kennedy, B.J., **288**, 29
Kennedy, J.D., **285**, 290; **289**, 95
Keppler, B.K., **292**, 127
Kerber, R.C., **291**, 109
Kernbach, U., **291**, 266
Ketchum, D.R., **294**, 200
Ketterle, M., **291**, 66
Keutel, H., **291**, 190
Khan, M.I., **292**, 112
Khan, T.A., **294**, 179
Kickelbick, G., **284**, 1; **290**, 100
Kim, D.W., **293**, 100
Kim, J., **293**, 140
Kim, J.C., **286**, 67
Kim, J.-S., **294**, 119
Kim, K.M., **292**, 52, 238
Kim, S.-Y., **294**, 119
Kim, Y.I., **286**, 67
Kimura, E., **294**, 56
Kingsley, S., **286**, 127
Kirchner, K., **286**, 114
Kiseleva, N.P., **285**, 203
Kitagawa, S., **293**, 20
Kitamori, Y., **290**, 251
Kivekas, R., **284**, 292
Kivlighon, L.M., **284**, 215
Klapötke, T.M., **294**, 68
Klein, H.-F., **287**, 167, 199
Klein, M., **293**, 115
Kleinpeter, E., **285**, 305
Kleverlaan, C.J., **284**, 61
Kloo, L.A., **292**, 108
Klooster, W.T., **285**, 10
Knör, G., **288**, 150
Ko, J., **289**, 141
Kobayashi, K., **296**, 67
Kobayashi, S., **296**, 158
Kodama, K.-y., **294**, 266
Koek, J.H., **295**, 189
Koelle, U., **290**, 44
Kofod, P., **286**, 98
Kohlen, E.W.M.J., **295**, 189
Kohout, J., **287**, 186
Koikawa, M., **285**, 156
Kokozay, V.N., **293**, 234
Kolis, J.W., **294**, 200
Kollár, L., **286**, 93
Komaei, S.A., **286**, 24; **287**, 226
Komatsu, T., **295**, 234
Komiya, S., **291**, 341; **294**, 266
Kondo, Y., **290**, 222
Konno, T., **288**, 7
Kooijman, H., **286**, 24; **287**, 226
Kopacka, H., **290**, 167
Korostylev, A.V., **295**, 164
Kou, X., **284**, 85
Koubek, E., **288**, 229
Kovalevsky, A.Y., **295**, 164
Kozanoglu, F., **284**, 229

Kozelka, J., **284**, 103
Kožíšek, J., **287**, 186
Krause Bauer, J.A., **288**, 159; **290**, 159
Krause, N., **296**, 1
Krauser, R., **285**, 184
Krebs, B., **285**, 262; **291**, 289
Kreiter, C.G., **284**, 229; **296**, 127
Krejzler, J., **286**, 175
Kremer, C., **294**, 47
Kremer, E., **294**, 47
Kresinski, R.A., **285**, 283; **290**, 189
Kruger, C., **288**, 118
Krüger, J., **296**, 150
Kruger, J.S., **285**, 217
Krukowska, K.S., **294**, 123
Krylov, V.K., **285**, 116
Kuang, S.-M., **284**, 119, 278; **293**, 106
Kubas, G.J., **294**, 240
Kubiak, R., **287**, 101; **288**, 174; **293**, 155
Kuge, K., **291**, 395
Kühn, F.E., **287**, 159
Kukushkin, V.Y., **285**, 116, 203; **292**, 272
Kukushkin, Y.N., **285**, 116, 203; **292**, 272
Kulsomphob, V., **296**, 170
Kumar Saha, M., **288**, 118
Kumar, S.B., **286**, 62
Kumeta, K., **291**, 231
Kundu, P., **288**, 118
Kuo, C.-N., **285**, 254; **293**, 12
Kuroboshi, M., **296**, 204
Kuroda-Sowa, T., **290**, 153, 251
Kurosaki, H., **294**, 56
Kurosawa, H., **296**, 37
Kurtaran, R., **286**, 181
Kwak, B., **290**, 21
Kwiatkowski, E., **293**, 115

LaDuca, R.L., Jr., **292**, 131
Lagrave, S., **285**, 49
Laguna, A., **290**, 44; **291**, 60
Lahoz, L., **285**, 25
Lai, W., **290**, 28
Lai, Y.-H., **285**, 31
Lam, K.-C., **296**, 170
Lam, L., **290**, 28
Lamprecht, A., **291**, 21
Lanfaloni, C., **288**, 200
Langer, V., **294**, 83
Langes, C., **290**, 167
Lang, H., **284**, 99
Lang, R., **294**, 62
Langsdorf, B.L., **294**, 207
Lapinte, C., **291**, 403
Lappin, A.G., **290**, 197
LaRonde, F.J., **296**, 208
Larsen, E., **293**, 239
Larsen, R.W., **288**, 74
Lau, K.Y., **285**, 223
Lauceri, R., **286**, 121
Laurent, J.-P., **285**, 49; **294**, 8
Lawrence, R.G., **285**, 283
Le, X., **285**, 326
Lebuis, A.-M., **294**, 224
Lecerf, A., **284**, 139
Lee, C.-L., **292**, 182
Lee, C.-S., **285**, 254
Lee, G.-H., **286**, 155; **293**, 30; **294**, 232
Lee, H., **289**, 11
Lee, S.-G., **286**, 215
Lee, S.S., **286**, 215; **292**, 52
Lee, Y.-A., **292**, 52; **295**, 214
le Gall, N., **291**, 166
Leglaye, P., **291**, 300
Leigh, G.J., **293**, 147
Leirer, M., **288**, 150
Lemke, M., **287**, 167
Lemke, U., **287**, 167
Leoni, P., **284**, 246
Le Stang, S., **291**, 403
Létard, J.-F., **291**, 279
Létard, S., **291**, 279
Leung, P.-H., **284**, 99
Leung, W.-H., **290**, 28; **294**, 73
Levason, W., **288**, 142
Levitin, I., **285**, 155
Lexa, D., **292**, 204
Li, A.-R., **290**, 51
Li, H., **285**, 25
Li, J., **294**, 109
Li, Y., **285**, 31
Liable-Sands, L., **284**, 175
Liao, F.-L., **295**, 229
Liaw, W.-F., **286**, 155
Licandro, E., **296**, 236
Lilie, J., **285**, 322
Lin, W., **288**, 215
Linden, A., **284**, 215; **285**, 122
Lindner, E., **296**, 103
Linert, W., **287**, 95
Linn, D.E., Jr., **291**, 142
Lipshutz, B.H., **296**, 164
Lisowski, J., **285**, 233
Liu, G., **287**, 109
Liu, H.-K., **295**, 129
Liu, J.-C., **295**, 229
Liu, Q., **286**, 7; **293**, 95
Liu, Q.-T., **290**, 121
Liu, S.-X., **294**, 163
Liu, Y.-J., **295**, 48
Lloret, F., **284**, 258; **287**, 173; **288**, 57
Lloris, M.J., **292**, 28
Lobbia, G.G., **285**, 81
Lohr, G.J., **296**, 13
Lonergan, M.C., **294**, 207
Long, D.-L., **285**, 241; **293**, 89; **294**, 73
Long, L., **285**, 326
Longhurst, S., **291**, 101
Longoni, G., **291**, 372
López, G., **287**, 37; **295**, 136
Lorenz, I.-P., **288**, 101
Lorenzo-Luis, P.A., **295**, 106
Losi, P., **288**, 244
Lotz, S., **284**, 127
Louattani, E., **291**, 207
Lu, J., **295**, 222
Lu, N., **289**, 198
Lu, Z.-l., **284**, 205
Luart, D., **291**, 166
Lucchese, Y., **295**, 209
Lukov, V., **293**, 210
Lumetta, G.J., **293**, 195
Lunghi, P., **286**, 134
Luo, Q.-H., **295**, 48
Luque, A., **290**, 57
Lutz, M., **287**, 113
Lyssenko, K.A., **295**, 164

Ma, C., **287**, 212
Ma, J.-G., **291**, 49
Ma, R., **291**, 238
Ma, S., **294**, 109
Macartney, D.H., **291**, 365
Macías, B., **288**, 127
Macchioni, A., **288**, 200
Macdonald, C.J., **292**, 198
MacDonnell, F.M., **293**, 37
Macleman, S., **294**, 99
Madeira, C., **293**, 1
Maekawa, M., **290**, 153, 251; **293**, 20
Mahmoudkhani, A.H., **294**, 83
Mahon, M.F., **287**, 89
Maina, T., **285**, 97; **295**, 1
Maiorana, S., **296**, 236
Makarov, S.V., **286**, 149
Mak, T.C.W., **284**, 119, 278; **290**, 127; **293**, 106
Maletin, Y.A., **284**, 180
Malet, P., **288**, 127
Malkov, A.V., **296**, 139
Maloney, D.J., **293**, 37
Manabe, K., **296**, 158
Mandel, F.S., **294**, 214
Mangold, A., **291**, 91
Mangrich, A., **290**, 207
Mangrich, A.S., **292**, 1
Manners, I., **287**, 152
Manotti Lanfredi, A.M., **290**, 180
Mansilla, N., **284**, 14
Manzoni de Oliveira, G., **288**, 101
Manzur, J., **286**, 74
Maran, F., **288**, 122
Marchetti, F., **294**, 28
Marchi, A., **286**, 46
Marek, J., **290**, 256
Maresca, K.P., **284**, 252; **295**, 39
Maresca, L., **285**, 301
Mariezcurrena, R., **294**, 47
Marken, F., **291**, 21
Marroni, R., **285**, 138
Marsich, N., **290**, 180
Martell, A.E., **284**, 30; **286**, 55; **287**, 47, 134, 193; **291**, 238; **293**, 206
Martins, L.M.D.R.S., **291**, 39
Martínez-Cruz, L.A., **284**, 14
Martínez-Máñez, R., **292**, 28
Martínez-Ripoll, M., **290**, 57
Marty, W., **288**, 181
Marvelli, L., **286**, 46

Masai, H., **296**, 231
Mascharak, P.K., **285**, 269
Massoud, S.S., **295**, 91
Masters, A.F., **294**, 99
Mastrorilli, P., **284**, 116
Masuda, H., **286**, 37
Matasi, J.J., **296**, 273
Matković-Čalogović, D., **284**, 223; **285**, 208; **294**, 35
Matsubayashi, G., **284**, 55; **295**, 239
Matsushima, H., **285**, 156
Matsuda, F., **295**, 239
Matsuda, I., **296**, 72
Matsuda, T., **294**, 119
Matsumoto, Y., **296**, 267
Matsuo, H., **294**, 56
Mattausch, H., **289**, 174
Matthews, J.P., **284**, 266
Mayr, A., **284**, 205; **285**, 223
McArdle, P., **291**, 252
McAuliffe, C.A., **292**, 213; **293**, 210
McCaffrey, L.J., **285**, 145
McCann, M., **293**, 160
McFarlane, W., **284**, 99
McKee, V., **293**, 160
McKeon, J.A., **290**, 133
McLemore, D.K., **294**, 193
McManus, D., **293**, 206
McNamara, B.K., **293**, 195
McQuillan, F.S., **285**, 55
Mederos, A., **291**, 158; **294**, 47; **295**, 106
Medina, G., **288**, 106
Meehan, P.R., **284**, 71
Menabue, L., **292**, 189
Mendizábal, F., **294**, 20
Meneghetti, M., **285**, 18
Meng, Q., **292**, 157
Mennekes, T., **289**, 58
Mereiter, K., **286**, 114
Messori, L., **285**, 309
Mestroni, S., **292**, 144
Meyer, R., **284**, 127
Meyer-Klaucke, W., **291**, 289
Michelin, R.A., **291**, 39
Midollini, S., **285**, 39; **294**, 47
Mikami, K., **296**, 267
Mikhel, I.S., **295**, 164
Miki, E., **290**, 36
Mikuriya, M., **290**, 228
Milet, A., **287**, 27
Miller, B., **290**, 237
Miller, J.S., **284**, 175
Miller, S.C., **287**, 109
Miller, S.M., **289**, 76
Millikan, M.B., **284**, 215
Mintert, M., **287**, 72
Mitra, S., **288**, 118
Mitsui, T., **291**, 395
Mizobe, Y., **285**, 336
Mizuno, K., **296**, 231
Möckel, H., **285**, 322
Mok, K.F., **285**, 31
Mokhir, A.A., **284**, 85
Moliner, N., **291**, 279
Molinié, P., **284**, 8
Molins, E., **287**, 37; **295**, 136
Moll, M., **284**, 288
Mombrú, A.W., **294**, 47
Monacelli, F., **295**, 200
Monari, M., **285**, 70; **291**, 372
Montet, Y., **284**, 103
Moreno-Esparza, R., **288**, 106
Moretó, J.M., **296**, 94
Mori, W., **293**, 20
Mori, Y., **296**, 158
Morikita, T., **291**, 341; **296**, 254
Morris, R.J., **292**, 220
Morrone, S., **292**, 163
Morson, P.A., **285**, 217
Morvillo, A., **288**, 122
Moss, J.R., **286**, 221
Motekaitis, R.J., **286**, 55; **291**, 238; **293**, 206
Mothes, E., **291**, 300
Moussa, S.M., **288**, 29
Muir, K.W., **284**, 107
Mukaida, M., **290**, 113
Mukaiyama, T., **296**, 86
Mullen, G.E.D., **294**, 170
Mullica, D.F., **285**, 318; **292**, 137
Munakata, M., **285**, 332; **290**, 153, 251; **293**, 20
Mundoma, C., **286**, 149
Muñoz, M.C., **288**, 57; **291**, 279
Murugavel, R., **292**, 105
Myakishev, K.G., **289**, 51

Nachtigal, C., **296**, 103
Nádasdi, L., **293**, 218
Nagao, H., **290**, 113
Nagao, N., **290**, 113
Nagasawa, A., **292**, 7
Nagata, T., **296**, 86
Nagayama, S., **296**, 158
Nagendran, S., **286**, 127
Naggar, E., **288**, 215
Nakagawa, M., **295**, 121
Nakahama, A., **284**, 55
Nakahata, S., **296**, 204
Nakamoto, T., **291**, 127
Nakamura, Y., **296**, 267
Nakanishi, S., **291**, 231
Nakano, M., **284**, 55
Nakao, Y., **293**, 20
Nakatsui, K., **296**, 33
Narbutt, J., **286**, 175
Nathan, L.C., **292**, 254
Natile, G., **285**, 301
Nauss, J.L., **288**, 159
Navarrete, A., **284**, 292
Navarro, R.E., **295**, 115
Nazeeruddin, M.K., **296**, 250
Neal, T.J., **291**, 49
Necas, M., **290**, 256
Nelson, J.H., **292**, 96
Neu, A., **289**, 58
Neumann, B., **291**, 311
Neumann, J., **293**, 239
Neves, A., **290**, 207
Ng, A.C.H., **288**, 226
Ng, D.K.P., **288**, 226
Ng, S.W., **293**, 147
Ngandu, L., **292**, 204
Nguyen, Q., **292**, 254
Nguyen, T.H., **291**, 136
Nicholson, B.K., **285**, 145; **295**, 18
Niero, A., **293**, 44
Nishide, H., **295**, 234
Nishiguchi, S., **296**, 37
Nishimoto, K., **295**, 180
Niu, S., **293**, 206
Nobile, C.F., **284**, 116
Nock, B., **285**, 97; **295**, 1
Noftle, R.E., **287**, 232
Nolan, S.P., **291**, 32
Noll, B.C., **285**, 134
Noveron, J.C., **285**, 269
Novosad, J., **290**, 256
Nugent, K.W., **285**, 122
Nukada, R., **290**, 228
Nunes, R., **284**, 30

Oberhauser, W., **290**, 167
Odashima, K., **296**, 158
Oeckler, O., **289**, 174
Ogasawara, M., **291**, 226
Ogino, H., **291**, 395
Oglieve, K.E., **291**, 212
Ogoshi, S., **296**, 37
Ohata, N., **286**, 37
Öhman, A., **286**, 181
Ohno, Y., **285**, 332
Öhrström, L., **292**, 266
Ohtsuka, Y., **296**, 86
Okamoto, K., **288**, 7
Okazaki, M., **291**, 395
Olabe, J.A., **285**, 129; **291**, 66
Oliva, G., **290**, 207
Oliver, A.G., **292**, 260
Olmstead, M.M., **285**, 134, 269
Onaka, S., **294**, 255
Onak, T., **289**, 11
O'Neill, L., **291**, 252
Ooyama, D., **290**, 113
Orioli, P., **285**, 309
Orrell, K.G., **295**, 56
Osada, M., **294**, 281
Osakada, K., **296**, 176
Osborne, A.G., **295**, 56
Oskui, B., **287**, 72
Ossola, F., **287**, 117
Ossowski, T., **285**, 1
Otto, J., **285**, 262
Owen, D.A., **296**, 139
Ozawa, F., **296**, 19
Özkar, S., **284**, 229

Packheiser, M., **296**, 127
Padilla-Tosta, M.E., **292**, 28
Padmavathy, D., **284**, 158
Paetzold, P., **289**, 58
Pajeau, T., **291**, 238
Pajunen, A., **284**, 292
Pakhomova, T.B., **292**, 272
Pala, J.S., **284**, 8

Palanca, P., **285**, 25
Palazzi, A., **291**, 333
Paleari, L., **288**, 244
Pang, Z., **284**, 124
Panunzi, A., **285**, 70
Panzalorto, M., **292**, 84
Paolo Speroni, G., **285**, 309
Papadimitriou, C., **290**, 256
Papadopoulos, M., **285**, 97; **295**, 1
Papadopoulos, M.S., **287**, 142; **290**, 247
Papagni, A., **296**, 236
Papucci, S., **284**, 246
Paradinas, R., **288**, 127
Pardo, T., **292**, 28
Parise, A.R., **291**, 66
Pariya, C., **285**, 107
Parkinson, J.A., **289**, 125
Park, K.-I., **289**, 141
Park, Y.-I., **289**, 141
Park, Y.-Y., **294**, 119
Pascaly, M., **291**, 289
Pasini, A., **285**, 249
Pasquali, M., **284**, 246
Paul, F., **291**, 403
Pavlović, G., **294**, 35
Pelagatti, P., **285**, 138; **292**, 121; **295**, 171
Pelecanou, M., **287**, 142
Pelizzi, C., **285**, 138; **295**, 171
Pelizzi, G., **292**, 121; **295**, 171
Pellei, M., **285**, 81
Pelosi, G., **286**, 134
Peñalba, E., **286**, 189
Peña, O., **286**, 74
Peng, S.-M., **286**, 155; **294**, 232
Pennington, W.T., **285**, 217; **294**, 200
Penn, J.H., **286**, 149
Perdicchia, D., **296**, 236
Perec, M., **284**, 49; **286**, 74
Pereira, M.M., **295**, 64
Pérez, J., **287**, 37; **295**, 136
Perez, W.J., **296**, 114
Perils, J., **294**, 114
Perkinson, J.L., **294**, 140
Pertici, P., **294**, 28
Pessoa, J.C., **293**, 1
Petersen, J.L., **286**, 149
Peters, G., **289**, 70
Pétillon, F.Y., **284**, 107
Petridis, A., **284**, 215; **285**, 122
Petrovskii, P.V., **295**, 164
Petrukhina, M.A., **284**, 304
Pettinari, C., **285**, 81
Pfab, H., **291**, 91
Phillips, D.J., **285**, 297
Philouze, C., **290**, 139
Pieraguoli, M.R., **286**, 46
Pierre, J.-L., **291**, 148
Pietzsch, H.-J., **295**, 1
Piltz, R.O., **288**, 29
Pinelli, S., **286**, 134
Pipal, J.R., **292**, 73
Pirmettis, I.C., **285**, 97; **287**, 142; **290**, 247; **295**, 1
Piro, O.E., **285**, 142
Platt, A.W.G., **290**, 189; **295**, 146
Plešek, J., **289**, 45
Polborn, K., **294**, 62
Pombeiro, A.J.L., **285**, 116; **291**, 39; **292**, 272
Poopasit, K., **292**, 57
Popović, Z., **285**, 208; **294**, 35
Potts, K.T., **288**, 189
Powell, A.K., **294**, 170
Preetz, W., **289**, 70
Pregosin, P.S., **290**, 95; **296**, 246
Pritchard, R.G., **292**, 213; **293**, 210
Prokopuk, N., **286**, 142
Pruchnik, F.P., **293**, 110
Purrello, R., **286**, 121
Pyo, M., **290**, 21
Pyo, S., **292**, 34

Qian, M., **294**, 1

Radanović, D.J., **292**, 16
Rae, A.D., **285**, 297
Rajasekharan, M.V., **284**, 158
Ramalakshmi, D., **284**, 158
Ramírez de Arellano, M.C., **295**, 136
Rapko, B.M., **293**, 195
Raptopoulou, C.P., **285**, 97; **287**, 142; **290**, 247; **295**, 1
Rarig, R.S., Jr., **292**, 131
Rath, N.P., **285**, 290; **289**, 95
Rauccio, M., **291**, 372
Razik, M., **293**, 155
Real, J.A., **291**, 279
Reddy, K.R., **284**, 158
Reed, D., **289**, 125
Reedijk, J., **286**, 24; **287**, 105, 226; **288**, 220, 239; **290**, 105
Rees, N.H., **284**, 99
Reffo, G., **286**, 134
Reichenbach, G., **288**, 200
Reid, G., **288**, 142
Reiland, V., **296**, 127
Renard, C., **296**, 139
Renkema, K.B., **291**, 226
Renn, O., **284**, 191
Renz, F., **287**, 95
Rheingold, A.L., **284**, 175; **287**, 209, 223; **289**, 149; **296**, 170
Ribas, J., **286**, 169
Ribou, A.-C., **288**, 134
Ricci, R., **288**, 200
Richards, R.L., **285**, 178
Richard Stephenson, G., **296**, 139
Rickard, C.E.F., **292**, 260
Rieder, A., **290**, 167
Riley, P.N., **288**, 35
Rillema, D.P., **296**, 114
Riou-Cavellec, M., **291**, 317
Riou, D., **291**, 317
Rippstein, R., **290**, 100
Ristanović, V.M., **292**, 16
Ritter, S.K., **287**, 232
Rivera, J., **292**, 34
Rivero, M., **294**, 47
Rives, V., **288**, 127
Rizzoto, M., **287**, 134
Robin, D., **292**, 204
Robinson, W.T., **292**, 73
Rocha Gonsalves, A.M.A., **295**, 64
Rocha, R.C., **285**, 197
Rochon, F.D., **295**, 25
Rodriguez, A., **284**, 133; **288**, 244
Rodriguez, M.D., **285**, 60
Rodriguez, V., **290**, 57
Röhr, C., **287**, 167
Romanowski, G., **293**, 115
Román, P., **290**, 57
Romero, J., **284**, 133; **294**, 79, 87; **295**, 39
Rompel, A., **291**, 289
Ronald Boone, J., **296**, 52
Roper, G.A., **294**, 224
Rösch, N., **286**, 30
Rose, D.J., **295**, 39
Rose, E., **296**, 139
Rose-Munch, F., **296**, 139
Rosini, C., **294**, 28
Ross, E., **286**, 55
Rossell, O., **291**, 247
Rossi, R., **286**, 46
Rosso, N.D., **284**, 30; **287**, 193
Roszak, A.W., **291**, 365
Rothery, J., **288**, 17
Rothwell, I.P., **288**, 35
Rotter, H.W., **287**, 218
Roviello, A., **292**, 163
Rüegger, H., **284**, 191; **290**, 64
Ruffo, F., **285**, 70
Ruiz, A., **295**, 64
Ruiz-Pérez, C., **284**, 258
Ruiz, R., **288**, 57
Ruiz-Ramírez, L., **288**, 106
Rusanova, J.A., **293**, 234
Russell, D.R., **295**, 146
Russell, S.W., **295**, 189
Russo, U., **287**, 117
Ryabov, A.D., **292**, 272
Ryan, M.D., **287**, 21
Ryu, K., **293**, 140

Saad, Y., **293**, 37
Saal, C., **291**, 82
Sachleben, R.A., **290**, 86
Sagara, A., **296**, 86
Sakagami, N., **288**, 7; **292**, 16
Sakai, K., **284**, 149
Sakamoto, S., **293**, 20
Sakano, T., **296**, 176
Saladini, M., **292**, 189
Salati, C., **295**, 171
Salaün, J.-Y., **291**, 166

Saldamlı, S., **284**, 229
Salek, S.N., **294**, 170
Sampanthar, J.T., **295**, 9
Sánchez, G., **287**, 37; **295**, 136
Sanchiz, J., **291**, 158
Sandoe, E.J., **296**, 139
Sanmartín, J., **293**, 210
Sano, H., **291**, 127
Santana, M.D., **284**, 296
Santini, C., **285**, 81
Santos, M.A., **284**, 20
Sanz, N., **286**, 169
Sanz, V., **285**, 25
Sapunov, V.N., **286**, 114
Sasabe, H., **288**, 134
Sasaki, A., **291**, 341
Sato, A., **294**, 266
Sattelberger, A.P., **288**, 69
Schauer, C.K., **294**, 140
Scheffer, M.H., **291**, 311
Scheidt, W.R., **291**, 49
Scheiring, T., **287**, 204
Schimek, G.L., **285**, 217; **294**, 200
Schindler, S., **288**, 206
Schmalle, H.W., **286**, 108
Schmid, M., **296**, 103
Schmid, R., **286**, 114
Schmidt, H.-G., **292**, 105
Schnebel, M., **296**, 150
Schneider, H., **285**, 1; **286**, 160
Schobert, R., **291**, 91
Schollhammer, P., **284**, 107
Schönenberger, H., **285**, 184
Schönherr, T., **295**, 84
Schrodt, A., **288**, 206
Schubert, U., **284**, 1; **290**, 100
Schulz, C., **291**, 449
Schulz, C.E., **291**, 49
Schweder, B., **286**, 14
Sclafani, J.A., **296**, 164
Scott, B.L., **294**, 240, 275
Scott Howell, F., **290**, 113
Seco, M., **291**, 247
Sedlák, P., **285**, 322
Segalés, G., **291**, 247
Seiffert, M., **288**, 101
Sen, A.K., **293**, 61
Senocq, F., **288**, 233
Sen, S., **288**, 118
Senthil Andavan, G.T., **286**, 127
Serra, M.E.S., **295**, 64
Serrano, J.L., **287**, 37; **295**, 136
Serratrice, G., **291**, 148
Severin, K., **294**, 62
Severin, T., **294**, 62
Sgarabotto, P., **285**, 138; **295**, 171
Shakadou, M., **296**, 80
Shang, M., **290**, 197
Shannon, P.J., **291**, 136
Shao, M.-Y., **285**, 254; **293**, 12
Sharma, K.D., **293**, 229
Shaw, S.L., **292**, 220
Sheffield, J.M., **292**, 213
Sheldrick, W.S., **287**, 72
Shelnutt, J.A., **291**, 49
Shen, M.-C., **295**, 48
Shen, Z., **292**, 117
Shepherd, R.E., **285**, 157, 170; **286**, 197; **292**, 225; **293**, 123
Shewmaker, T.E., **285**, 60
Shi, J.-C., **290**, 121
Shi, Q., **284**, 112; **294**, 109
Shibahara, T., **295**, 180
Shibata, T., **296**, 33
Shido, T., **294**, 281
Shimizu, M., **296**, 231
Shimizu, T., **296**, 278
Shionoya, M., **294**, 56
Shongwe, M.S., **290**, 228
Shriver, D.F., **286**, 142
Shu, L., **292**, 34
Shu, M.-H., **290**, 127
Shukla, A.D., **285**, 89
Shukri, K., **291**, 178
Shyu, H.-L., **290**, 8
Sieler, J., **293**, 234
Sift, B.H., **291**, 289
Šik, V., **295**, 56
Silvestru, C., **287**, 61
Simon, A., **289**, 174
Simonsen, O., **295**, 153
Simoyi, R.H., **286**, 149
Sinbandhit, S., **291**, 449
Singhal, A., **285**, 190
Skelton, B.W., **287**, 27
Skidd, G.M., **291**, 142
Skoda-Földes, R., **286**, 93
Slawin, A.M.Z., **290**, 1
Slep, L.D., **285**, 129
Sletten, J., **287**, 173
Smeets, W.J.J., **288**, 220; **290**, 105; **291**, 438
Smith, B.E., **291**, 212
Smith, K.C., **291**, 365
Smith, R.J., **294**, 170
Smoleński, P., **293**, 110
Soai, K., **296**, 33
Sobota, P., **285**, 178
Sohn, Y.S., **292**, 52, 238; **295**, 214
Solans, X., **286**, 62, 169, 189; **291**, 279
Soldin, Ž., **294**, 35
Solntsev, K.A., **289**, 76
Solujic, L., **292**, 96
Song, R., **292**, 238
Soo Lah, M., **290**, 21
Soto, J., **292**, 28
Sousa, A., **284**, 133; **293**, 210; **294**, 79, 87; **295**, 39
Sousa-Pedrares, A., **284**, 133; **295**, 39
Souza, D.D.H.F., **290**, 207
Spagna, R., **285**, 81
Spek, A.L., **286**, 24; **287**, 105, 113, 226; **288**, 220, 239; **290**, 105; **291**, 326, 438
Spencer, P.B., **284**, 37
Speroni, G.P., **285**, 70, 309
Spies, H., **295**, 1
Spodine, E., **286**, 74
Squattrito, P.J., **293**, 61
Srebnik, M., **290**, 159
Stahl, L., **288**, 154
Stammler, H.-G., **291**, 311
Stampfl, T., **290**, 167
Stash, A.I., **284**, 180
Stassinopoulou, C.I., **287**, 142
Steel, P.J., **284**, 273; **288**, 53
Steimann, M., **296**, 103
Stein, B.K., **290**, 189
Stener, M., **286**, 30
Stenhagen, G., **292**, 266
Stern, C., **286**, 142
Steuer, B., **289**, 70
Storhoff, J.J., **292**, 220
Storrier, G.D., **284**, 76; **288**, 189
Strauss, S.H., **289**, 76; **294**, 193
Streib, W.E., **291**, 226
Stringfield, T.W., **285**, 157; **292**, 225
Strizhakova, N.G., **284**, 180
Stufkens, D.J., **284**, 61
Suades, J., **291**, 207
Suenaga, Y., **285**, 332; **290**, 153, 251
Suen, M.-C., **288**, 82
Sueros Velarde, F.J., **285**, 142
Suescun, L., **294**, 47
Sugawara, M., **296**, 19
Sugimoto, K., **290**, 153
Sugioka, T., **296**, 195
Sugiyama, M., **294**, 56
Suh, I.-H., **293**, 100
Suib, S.L., **294**, 123
Sülü, M., **293**, 70
Sumi, M., **294**, 56
Sun, W.-Y., **290**, 127; **295**, 129
Sun, Y., **291**, 238
Surdykowski, A., **293**, 239
Suto, A., **290**, 222
Suzuki, T., **292**, 7
Suzuki, Y., **290**, 36
Svarovsky, S.A., **286**, 149
Svoboda, I., **287**, 181, 186
Swanson, B.I., **294**, 275
Sztyk, E., **293**, 239
Szölösy, Á., **296**, 222
Szpoganicz, B., **284**, 30; **287**, 134, 193

Tabbi, G., **288**, 239
Tajima, K., **288**, 90
Takahashi, S., **296**, 195
Takao, N., **290**, 222
Takata, T., **291**, 231
Talarmin, J., **284**, 107
Tamburini, S., **293**, 178
Tamm, M., **288**, 47
Tamura, H., **295**, 239
Tan, W.-L., **294**, 73
Tanaka, H., **296**, 204
Tanaka, Y., **294**, 56
Tang, W.-X., **287**, 52; **290**, 127; **295**, 129
Tang, Z., **284**, 112; **294**, 109
Taniguchi, I., **294**, 56
Tanturli, R., **294**, 28
Tarasconi, P., **286**, 134

Tate, D., **293**, 37
Taylor, M.J., **292**, 108
Tcherkaev, G.V., **295**, 164
Tchertanov, L., **290**, 139
Teat, S.J., **289**, 95
Tedesco, V., **291**, 21
Tekkaya, A., **284**, 229
Temizdemir, S., **295**, 249
Temple, K., **287**, 152
Terada, M., **296**, 267
Terasaki, Y., **295**, 84
ter Steeg, P.F., **295**, 189
Terzis, A., **285**, 97; **287**, 142; **290**, 247
Tessier, C., **295**, 25
Thaler, A., **285**, 1; **286**, 160
Thalji, R.K., **295**, 91
Thiele, G., **287**, 218
Thomas, F., **291**, 148
Thomas, L.M., **286**, 207
Thomas, R.L., **289**, 134
Thorand, S., **296**, 1
Thornton-Pett, M., **285**, 290; **289**, 95
Tiekink, E.R.T., **285**, 190; **286**, 127
Tilley, T.D., **287**, 152
Tippmann, E.M., **291**, 142
Tiripicchio, A., **296**, 236
To, H.T., **289**, 11
Tobita, H., **291**, 395
Togni, A., **291**, 355; **296**, 183
Tohier, M., **291**, 449
Tokii, T., **285**, 156
Tokuno, T., **290**, 145; **295**, 71
Toma, H.E., **285**, 197
Tomasin, P., **287**, 117; **293**, 178
Tomaz, I., **293**, 1
Tomioka, T., **296**, 164
Tomiyasu, H., **294**, 119
Tomizawa, H., **290**, 36
Torii, S., **296**, 204
Torralba, M.C., **284**, 296
Torres, A.S., **293**, 37
Toupet, L., **291**, 166
Tran, U., **289**, 11
Trenary, M., **289**, 191
Triki, S., **284**, 8
Trofimenko, S., **287**, 27
Trullinger, G.E., **294**, 140
Truong, T.U., **289**, 11
Trylus, K.-H., **291**, 266
Tschoerner, M., **290**, 95
Tsoukalas, C., **285**, 97; **295**, 1
Tsubomura, T., **284**, 149
Tsuboya, N., **290**, 113
Tsuchida, E., **295**, 234
Tsuji, M., **286**, 233
Tsutsumi, K., **296**, 37
Tuckerman, M., **291**, 388
Turner, M., **290**, 133

Ueda, T., **284**, 237
Uemura, M., **296**, 80
Ugozzoli, F., **290**, 180
Uguagliati, P., **293**, 44
Ungváry, F., **296**, 52
Unoura, K., **292**, 7
Urrutigoïty, M., **288**, 233

Vacca, A., **285**, 39
Valente, E.J., **284**, 300
Valko, M., **287**, 186
Valle, A., **285**, 39
Valle, G., **285**, 18; **290**, 14
Vallorani, F., **285**, 81
Valtancoli, B., **288**, 244
van Albada, G.A., **286**, 24; **287**, 226; **288**, 220; **290**, 105
van de Burgt, L.J., **293**, 167
VanDerveer, D., **285**, 217
van der Wolf, L., **295**, 189
van Koten, G., **287**, 113; **291**, 326
van Rooyen, P.H., **284**, 127
Varghese, B., **285**, 190
Varzatskii, O.A., **284**, 180
Vázquez-López, E., **294**, 87
Vega, A., **286**, 74
Vega, D., **284**, 49
Veldman, N., **287**, 105; **288**, 239; **290**, 105
Veltsistas, P., **290**, 256
Venanzi, L.M., **284**, 191; **290**, 64; **293**, 70
Vencato, I., **290**, 207
Verani, C.N., **290**, 207
Veronese, A.C., **285**, 18
Vetralla, G., **288**, 200
Vicente, R., **286**, 62, 189
Vicentini, G., **292**, 249
Vigato, P.A., **287**, 117; **293**, 178
Vij, A., **292**, 157
Vikić-Topić, D., **285**, 208; **294**, 35
Viland, T., **285**, 262
Villa, M.V., **288**, 127
Visentin, F., **293**, 44
Vitoria, P., **290**, 57
Vittal, J.J., **295**, 9
Vivekanandan, K., **286**, 127
Vogler, A., **288**, 150
Voigt, B., **296**, 150
Volkov, V.V., **289**, 51
Vollrath, H., **296**, 26
Voloshin, Y.Z., **284**, 180
von Ragué Schleyer, P., **289**, 58
Voss, E.J., **285**, 10
Vrdoljak, V., **284**, 223

Wada, T., **288**, 134
Wagenknecht, P.S., **292**, 254
Wagner, G., **292**, 272
Wainwright, K.P., **294**, 103
Wakamatsu, K., **295**, 180
Waldbaum, B.R., **291**, 109
Walmsley, J.A., **285**, 60
Walsh, R.J., **284**, 37
Walters, M.A., **291**, 388
Walther, D., **286**, 14
Waltz, W.L., **285**, 322
Wang, C., **288**, 1
Wang, D.-F., **290**, 127
Wang, E., **295**, 244
Wang, J.-C., **288**, 82; **292**, 182
Wang, S.-L., **295**, 229
Wang, S.-X., **287**, 82
Wang, T.-G., **285**, 152
Wang, Y., **290**, 8; **293**, 30
Wang, Z., **288**, 215
Wang, Z.-M., **293**, 167
Ward, M.D., **285**, 89; **292**, 231
Ward, M.S., **285**, 170; **286**, 197
Wartchow, R., **296**, 150
Wasgestian, F., **285**, 313
Watanabe, H., **296**, 204
Watanabe, M., **290**, 256
Watanabe, T., **296**, 80
Waters, J.M., **290**, 228
Watkinson, M., **293**, 210
Watson, M., **292**, 172
Wazzeh, R.A., **288**, 40
Webb, L., **296**, 52
Weber, L., **291**, 311
Wedgwood, J.L., **290**, 189
Wegner, P., **296**, 103
Wei, H.-H., **290**, 8, 51; **293**, 30
Weiner, B.R., **294**, 95
Weiner, H., **291**, 426
Weintraub, S.T., **294**, 133
Welch, A.J., **289**, 125
Welch, M.J., **291**, 238
Welker, M.E., **296**, 13
Weller, A.S., **289**, 85
Wen, R., **292**, 149; **295**, 91
Wen, T., **286**, 7; **293**, 95
Wentland, M.P., **288**, 189
Went, M.J., **294**, 170
Wernberg, O., **295**, 153
Wessels, P.L., **284**, 127
West, D.X., **294**, 79
Weyhermüller, T., **291**, 258
Wharf, I., **294**, 224
White, A.H., **287**, 27
White, P.S., **294**, 140
Whitmire, K.H., **294**, 153
Whittle, B., **285**, 89
Whittlesey, B.R., **284**, 124
Widmaier, R., **290**, 159
Wiede, P., **284**, 1
Wieghardt, K., **291**, 258
Wild, S., **290**, 237
Willey, G.R., **284**, 71
Williams, D.J., **285**, 217
Williams, G.N., **290**, 145
Williams, I.D., **294**, 73
Windmiller, M.S., **285**, 318
Winkler, A., **284**, 288
Wittmer, F.-G., **296**, 127
Wojtczak, A., **293**, 239
Wone, D., **292**, 254
Wong, E.H., **287**, 209, 223
Wong, W.-T., **290**, 28
Wonnemann, R., **285**, 262
Woollins, J.D., **290**, 1, 256
Woong Cho, K., **290**, 21
Wrackmeyer, B., **296**, 26
Wu, B.-M., **284**, 278

Wu, C.-P., **292**, 182
Wu, D., **286**, 7; **293**, 95
Wu, D.-X., **290**, 121
Wu, J., **287**, 212
Wu, L.P., **290**, 251
Wu, Q., **294**, 109
Wu, Y.-Y., **288**, 82; **292**, 182

Xin, X., **294**, 73
Xiong, R.-G., **288**, 215; **294**, 95
Xu, D.-M., **294**, 163
Xu, G.-X., **287**, 82
Xu, J.-Q., **285**, 152
Xu, X., **285**, 322
Xu, Y., **295**, 222, 229
Xue, F., **284**, 119; **290**, 127
Xue, W.-M., **296**, 114

Yamada, T., **296**, 86
Yamada, Y., **288**, 7; **294**, 255
Yamaguchi, I., **296**, 176
Yamaguchi, M., **296**, 67
Yamamoto, T., **296**, 176, 254, 278
Yamamoto, Y., **286**, 233
Yamashita, M., **294**, 255
Yamauchi, O., **286**, 37
Yamazaki, A., **292**, 7
Yan, J., **296**, 273
Yan, W., **287**, 212
Yang, H., **287**, 223
Yang, S., **294**, 109
Yap, G.P.A., **287**, 209, 223
Yatani, A., **293**, 20
Yeamin Reza, M., **285**, 156
Yeh, W.-Y., **294**, 232
Yeung, C.-H., **290**, 28
Yeung, Y.-O., **288**, 226
Yin, X., **286**, 221
Yoneda, E., **296**, 195
You, X., **287**, 14; **294**, 1
You, X.-Z., **292**, 117; **295**, 229
Youngme, S., **292**, 57
Yousaf, M., **289**, 198
Yu, K.-B., **295**, 129
Yuki, M., **291**, 395
Yun, Y.K., **296**, 261
Yutronic, N., **294**, 20

Zacchini, S., **291**, 372
Zafar Kaifi, F.M., **285**, 297
Zahn, T., **288**, 154
Zakeeruddin, S.M., **296**, 250
Zamora, F., **284**, 14
Zangrando, E., **285**, 116, 203
Zanotti, V., **291**, 333
Zapf, P.J., **292**, 131
Zeng, Q., **287**, 14; **294**, 1
Zhan, S., **292**, 157
Zhang, F., **294**, 109
Zhang, L.-M., **284**, 278
Zhang, S.-W., **296**, 195
Zhang, W., **287**, 212
Zhang, Y.-D., **293**, 89
Zhang, Z.-Z., **284**, 119, 278; **293**, 106
Zhao, J., **296**, 204
Zheng, H., **294**, 73
Zheng, L.-M., **287**, 52
Zheng, Y., **290**, 121; **294**, 163
Zhou, L.-M., **285**, 152
Zhou, X., **294**, 207
Zhou, X.-H., **285**, 152
Zhou, Z., **285**, 326
Zhu, H., **286**, 7; **293**, 95
Zhu, H.-L., **287**, 52; **295**, 129
Zhuang, J.-Z., **295**, 229
Ziegler, M.L., **288**, 154
Zimmermann, W., **292**, 127
Zinner, K., **292**, 249
Zinner, L.B., **292**, 249
Zipper, L., **288**, 181
Zompa, L.J., **292**, 172
Zanonato, P.R., **286**, 103
Zorbas, H., **290**, 237
Zovinka, E.P., **285**, 134
Zsolnai, L., **296**, 222
Zubieta, J., **284**, 133, 252; **292**, 41, 131; **295**, 39
Zubkowski, J.D., **284**, 300
Zukerman-Schpector, J., **292**, 249
Zuo, J.-L., **292**, 117
Zürcher, S., **291**, 355
Zybill, C.E., **291**, 380

ELSEVIER

www.elsevier.nl/locate/ica

Inorganica Chimica Acta 296 (1999) 291–341

Inorganica Chimica Acta

Cumulative Subject Index of Volumes 284–296

Ab initio calculations
ab initio calculations and X-ray structure determinations of dichloro(*O*-*O*′-*O*″-1,3,5-cyclohexanetriol)copper(II) and bis(*O*-*O*′-*O*″-1,3,5-hexanetriol)nickel(II), **284**, 284
coordination geometries for palladium and platinum: theoretical studies and the synthesis and structure of tris(indazol-1-yl)-borate complexes $MMe_3\{(ind)_3BH\}$, **287**, 27
conversion of site-specific deuterated *closo*-1,2- and 1,7-$C_2B_{10}H_{12}$ to deuterated *nido*-7,8- and 7,9-$[C_2B_9H_{12}]^-$ ions, respectively; mechanistic inferences. Comparisons of GIAO-NMR generated $[C_2B_9H_{12}]^-$ chemical shifts with experimental values, and consideration of two principal isomers of the 7,8-$[C_2B_9H_{12}]^-$ ion by comparison of geometry optimization data, NMR chemical shift data, and NMR coupling data to respective experimental data, **289**, 11
a facile synthesis of 1,3-disilacyclohexanes, **296**, 231
Absolute stereochemistry
stereochemical investigation of a kinetically labile chiral organopalladium complex in solution, **284**, 99
π-Acceptors
Rh-complex promoted synthesis of poly(arylacetylene) with pendant TTF units and formation of CT complexes of the polymer with π-acceptors, **296**, 278
Acetalization
some new chain-like terdentate phosphines, their ruthenium(II) coordination chemistry and the activity of the cations $[Ru(MeCN)_3(PhP\{CH_2CH_2P(p\text{-}X\text{-}C_6H_4)_2\}_2)]^{2+}$ (X = H, F, Me and OMe) as acetalization catalysts, **290**, 64
Acetamide complexes
evidence for a new nitrosyl-to-nitro reaction and an acetamide formation in the $\{RuNO\}^6$-type complex, *cis*-$[Ru(NO)(CH_3CN)(bpy)_2]^{3+}$, **290**, 113
Acetylacetonate complexes
studies of the binding modes of carboxylate donors with manganese(III) complexes containing tetradentate Schiff base ligands. Crystal structures of the complexes $[Mn(3\text{-}CH_3Osalpn)(HO_2CC_6H_4CO_2)]_n$ and $[\{Mn(5\text{-}NO_2salpn)(OH)(H_2O)\}\{Mn(5\text{-}NO_2salpn)(HO_2CC_6H_4CO_2)(H_2O)\}]\cdot C_2H_5OH$, **293**, 210
Acetylphosphane complexes
reactions of $Cp'(CO)_2MnPPh_2H$ with CH_3COCl and $CH_3S(O)_2Cl$ in THF/triethylamine: evidence of the first complex stabilization of a phosphorus homolog of the sulfonamides, **288**, 101
Activation of heterocycles
C–S, C–H, and N–H bond cleavage of heterocycles by a zero-valent iron complex, $Fe(N_2)(depe)_2$ [depe = 1,2-bis(diethylphosphino)ethane], **291**, 341
Activation volume
high pressure NMR studies on ligand exchange reactions in the equatorial sites of $[VO(L)_5]^{2+}$ (L = acetonitrile, *N*,*N*-dimethylformamide, and *N*,*N*-dimethylacetamide), **294**, 119
Acyl complexes
synthesis and characterization of long chain acyl and alkyl complexes of the types $[(\eta^5\text{-}C_5R'_5)Fe(CO)_2\{C(O)R\}]$ (R′ = H, CH_3), $[CpW(CO)_3\{C(O)R\}]$ and $[CpW(CO)_3R]$ (R = *n*-alkyl), **286**, 221
synthesis of bis-(acyl, carbamoyl or alkoxycarbonyl) iron monomer or dimer complexes by reaction of oxalyl chloride with various tetracarbonyl ferrates: $\{Fe[C(O)R](CO)_4\}^-$, **291**, 166
Acyliron complexes
addition of aryl substituents to cyclohexadienyliron electrophiles in the development of routes to C_{12} central building blocks for alkaloid synthesis, **296**, 139
Adamantane
oxidation of adamantane by palladium acetate systems, **294**, 99
Adenine mononucleotides
synthesis and characterization of compounds of di- and tributyltin chloride with adenine and guanine mononucleotides, **285**, 60
Alditolate complexes
structures of 1,3-bis(diphenylphosphino)propane platinum(II) alditolate complexes, **285**, 10
Aldol reaction
synthetic reactions using organometallics in water. Aldol and allylation reactions catalyzed by Lewis acid–surfactant-combined catalysts/Brønsted acids systems, **296**, 158
Alkaloid synthesis
addition of aryl substituents to cyclohexadienyliron electrophiles in the development of routes to C_{12} central building blocks for alkaloid synthesis, **296**, 139
Alkane activation
oxidation of adamantane by palladium acetate systems, **294**, 99
Alkene–carbene complexes
chemoselective synthesis of (allyloxy)carbene chelate complexes of iron, **291**, 449
Alkene complexes
phase dependent η^2-alkene versus η^1-alkyl transformation in platinum(II) complexes, **285**, 301
synthesis of *N*-allyl-2,3-dihydrobenzoxazol-2-ylidene complexes of chromium(0) and tungsten(0), **288**, 47
Alkene hydrogenation
structural, spectroscopic and catalytic properties of water-soluble hydride rhodium complexes $[RhH(Rtpa^+I^-)_4]H_2O$ (R = Me, Et), **293**, 110
Alkenyl complexes
highly enantioselective catalytic isopropenylation of aldehydes using diisopropenylzinc, **296**, 33
reactions of σ-π alkenyl complexes $[Fe_2(CO)_6(\mu\text{-alkenyl})(\mu\text{-}PPh_2)]$ with bis(diphenylphosphino)methane (dppm): carbonyl substitution, migratory carbonyl insertion, phospho-

Elsevier Science S.A.
PII: S0020-1693(99)00486-7

rus–carbon bond formation and cleavage, proton shift and alkenyl isomerisation reactions, **291**, 178

Alkoxide complexes
reactions of $Ti(O\text{-}i\text{-}Pr)Cl_3$ with bidentate ligands containing hydroxy and methoxy groups. The crystal structure of $Ti(O\text{-}i\text{-}Pr)Cl_3(HOCH_2CH_2OCH_3)$ and the ^1H variable-temperature NMR studies of solution structures of $Ti(O\text{-}i\text{-}Pr)Cl_3(HOCH_2CH_2OCH_3)$ and $Ti(O\text{-}i\text{-}Pr)Cl_3(CH_3OCH_2CH_2OCH_3)$, **285**, 254

Alkoxycarbonyl complexes
synthesis of bis-(acyl, carbamoyl or alkoxycarbonyl) iron monomer or dimer complexes by reaction of oxalyl chloride with various tetracarbonyl ferrates: $\{Fe[C(O)R](CO)_4\}^-$, **291**, 166

Alkylation
chemistry of polyfunctional molecules : CXXXI. Crystal structure and preparation of [bis(diphenyl–methyl–phosphino)iminium] tetraphenyloborate, $[(C_6H_5)_2(CH_3)P{=}N{=}P(CH_3)(C_6H_5)_2]^+[B(C_6H_5)_4]^-$, **284**, 288
highly enantioselective catalytic isopropenylation of aldehydes using diisopropenylzinc, **296**, 33

Alkyl complexes
monoindenyl titanium alkyl halides. The synthesis and molecular structures of $(\eta^5\text{-}C_9H_7)TiBr_3$, $(\eta^5\text{-}C_9H_7)Ti(CH_3)Br_2$, and $(\eta^5\text{-}C_9H_7)Ti(CH_3)Cl_2$, **292**, 220
phase dependent η^2-alkene versus η^1-alkyl transformation in platinum(II) complexes, **285**, 301
syntheses, spectroscopic properties and photochemistry of $[Re(R)(CO)_3(4,4'\text{-}Me_2\text{-}bpy)]$ (R = alkyl) complexes, **284**, 61
synthesis and characterization of long chain acyl and alkyl complexes of the types $[(\eta^5\text{-}C_5R'_5)Fe(CO)_2\{C(O)R\}]$ (R′ = H, CH_3), $[CpW(CO)_3\{C(O)R\}]$ and $[CpW(CO)_3R]$ (R = *n*-alkyl), **286**, 221

Alkylidene complexes
the μ-sulfonium–methylidene diiron complexes $[Fe_2\{\mu\text{-}C(X)SMe_2\}(\mu\text{-}CO)(CO)_2(Cp)_2]SO_3CF_3$ (X = CN, H) as precursors of μ-alkylidene complexes, **291**, 333

Alkyl sulfonated phosphine complexes
water soluble platinum(II) and palladium(II) complexes of alkyl sulfonated phosphines, **290**, 189

Alkyne
$Ni(COD)_2$: an efficient promoter in the construction of 5-cyclopent-2-enone derivatives from allyl halides, alkynes and carbon monoxide, **296**, 94

Alkyne complexes
synthesis and formation of metal complexes of 4-alkynyl and 4-cyano-2,6-diisopropylphenylisocyanides, **284**, 205

Alkyne polymerization
organometallic electron reservoir sandwich iron complexes as potential agents for redox and electron transfer chain catalysis, **291**, 1

Alkynes
photochemical reactions of transition metal organyl complexes with olefins: XXII. Photoreactions of pentacarbonyl(μ-$\eta^{5:3}$-4*H*-indenyl)diiron(Fe–Fe) with alkynes—substituent controlled C–H transfer or C–C bond formation, **296**, 127
spirocyclic stannole derivatives by 1,1-organoboration of 2,2-bis(1-alkynyl)-1,3-di-tert-butyl-1,3,2-diazastannacyclohexanes, **296**, 26

Alkynyl complexes
redox active organoiron alkynyl centers with a coordination site: an approach toward molecular plugs. Complexation reactions with $(THF)W(CO)_5$ and $(PhCN)_2MCl_2$ (M = Pd, Pt), **291**, 403

Allene complexes
reactions of cobaloximes with vinyl allenes as a route to 1-cobaloxime substituted 1,3-dienyl complexes, **296**, 13

Allenes
copper-mediated 1,6-, 1,8-, 1,10- and 1,12-addition and 1,5-substitution reactions in organic synthesis, **296**, 1

Allenyl complexes
cross-coupling reactions proceeding through η^1- and η^3-propargyl/allenyl–palladium(II) intermediates, **296**, 37

Allylation
Barbier-type allylation of carbonyl derivatives by use of aluminum as an electron pool. Double allylation of carboxylic esters, **296**, 204

Allylbenzylmalonate ligand
synthesis, structures, and linkage isomerism of (allylbenzylmalonate)platinum(II) complexes, **295**, 214

Allyl complexes
new ferracyclic π-allyl and η^3-pentadienyl complexes by carbonylative lactonization of unsaturated diols, **291**, 91
palladium(II) allyl complexes with potentially terdentate ancillary ligands. Mechanism of allyl amination by piperidine, **293**, 44

Allyl derivatives
$Ni(COD)_2$: an efficient promoter in the construction of 5-cyclopent-2-enone derivatives from allyl halides, alkynes and carbon monoxide, **296**, 94

Allylic C–H activation
intramolecular coupling of conjugated enynes with cyclopropylcarbene–chromium complexes; a complex reaction pathway, **296**, 273

Allylsilanes
reactions of bis(silyl)palladium(II) complexes with allyl halides. Synthesis of mono(silyl)palladium(II) halides and X-ray structure of *trans*-$PdCl(SiF_2Ph)(PMe_2Ph)_2$, **296**, 19

Aluminophosphates
hydrothermal synthesis of microporous materials with the direct incorporation of porphyrin molecules, **294**, 179

Aluminum amine
synthesis of new Group III fluoride–ammonia adducts in supercritical ammonia: structures of $AlF_3(NH_3)_2$ and $InF_2(NH_2)(NH_3)$, **294**, 200

Aluminum metal
Barbier-type allylation of carbonyl derivatives by use of aluminum as an electron pool. Double allylation of carboxylic esters, **296**, 204

Aluminum nitride
synthesis of new Group III fluoride–ammonia adducts in supercritical ammonia: structures of $AlF_3(NH_3)_2$ and $InF_2(NH_2)(NH_3)$, **294**, 200

Amide
effect of N–H···S amide–thiolate hydrogen bonding on bond lengths in rubredoxin models $[(CH_3)_3NCH_2CONH_2]_2[M(S_2\text{-}o\text{-}xyl)_2]$, **291**, 388

Amide complexes
cadmium(II) complexes with [(4-methylphenyl)sulfonyl]-2-pyridyl-amine: crystal structures of bis{[(4-methylphenyl)-sulfonyl]-2-pyridyl-amide}cadmium(II); aquo-bis(pyridine) bis{[(4-methylphenyl)sulfonyl]-2-pyridyl-amide}cadmium(II) and 1,10-phenanthroline bis{[(4-methylphenyl)sulfonyl]-2-pyridyl-amide}cadmium(II), **294**, 87
carbon–oxygen bond cleavage: a new reaction mode for amides bonded through oxygen to cobalt(III), **287**, 1
complexation of uranyl ion by tetrahexylmalonamides: an equilibrium modeling and infrared spectroscopic study, **293**, 195
electrochemical synthesis of cobalt, nickel, copper, zinc and cadmium complexes with *N*[(2-hydroxyphenyl)methylidine]-*N*′-tosylbenzene-1,2-diamine. The crystal structures of {(1, 10-phenanthroline)[*N*-(2-oxophenyl)methylidine]-*N*-tosylbenzene-1,2-diaminato}nickel(II) and {(1,10-phenanthroline)[*N*-(2-oxophenyl)methylidine]-*N*′-tosylbenzene-1,2-diaminato}-copper(II), **295**, 39

synthesis, structure and EPR spectroscopy of a vanadium(IV)–amide metallacyclic complex, **285**, 134

Amidine complexes
formation of two stable seven-membered chelate rings bound to one metal center due to addition of bifunctional amine to coordinated nitriles, **285**, 203

Amido complexes
synthesis and structure of a novel terminal imido complex of titanium, **292**, 105

Amine complexes
anionopentaaminecobalt(III) complexes with polyamine ligands: Part 30. The base hydrolysis kinetics and structure determination of some $[CoCl(N)_5]^{2+}$ complexes containing pyridine ligands, **288**, 181
control of the coordination mode of NO_2^- by the nature of the amine ligand: the importance of H-bond. Crystal structures of the [*mer*-Ni(dpt)$(NO_2)_2$] (**I**) and [*trans*-Ni(333-tet)$(NO_2)_2$] (**II**), **292**, 149
crystal structure of (μ-oxo)bis{aquatris(2-pyridylmethyl)amine}-iron(III) perchlorate, **284**, 124
palladium(I) phosphido-bridged dinuclear derivatives with both phosphine and amine ligands, **284**, 246
synthesis and structures of (*SP*-4-2)-diiodobis(2-hydroxyethylamine)platinum(II), (*SP*-4-2)-dichlorobis(2-hydroxyethylamine)platinum(II) and (*OC*-6-22)-bis(2-hydroxyethylamine)tetrachloroplatinum(IV) in the crystal, **292**, 127
synthesis, crystal structure and magnetic properties of *catena*-poly[(2,2′-bipyridylcopper)-di-μ-thiocyanato] and bis(2,2′-bipyridyl)thiocyanatocopper(II) perchlorate, **286**, 169
the hydrothermal synthesis and structural characterization of a new class of compounds: nickel organoamine-halocadmates, **292**, 41

Amine-*N*-oxide complexes
neodymium and europium picrate complexes with 3-picoline-*N*-oxide (3-picNO), **292**, 249

Amino acid complexes
hydrogen bonding in *cis*-bis(L-alaninato)copper(II): a single crystal neutron diffraction study, **288**, 29
the interaction of BMXD and its copper(II) complexes with glycine, aspartic acid, and histidine, **286**, 55

Amino acids
stereoselective reduction of ketones using extracoordinate silicon: C_2-symmetric ligands, **296**, 208

Aminoalcohols
aminophosphanes derived from chiral β-aminoalcohols and carbonyliron complexes thereof, **291**, 300

4-Amino-3,5-dimethyl-1,2,4-triazole
synthesis, structure and magnetic properties of a linear trimanganese(II,III,II) complex with hydroxo and 1,2,4-triazole as bridging ligands, **295**, 229

Aminoiminomethanesulfonic acid
structure and stability of aminoiminomethanesulfonic acid, **286**, 149

Amino ligands
the preparation and coordination chemistry of $^iPr_2P(E)NHP(E')^iPr_2$ (E, E′ = Se; E = Se, E′ = S; E = S, E′ = O; E,E′ = O, **290**, 1

Amino–phenol
synthesis, molecular and crystal structure of the zwitterionic Re(V)–oxo complex $ReOCl_3(L)(PPh_3)$ with L = 2-(diethylaminomethyl)-4-methylphenol, **295**, 209

Aminophosphine complexes
aminophosphanes derived from chiral β-aminoalcohols and carbonyliron complexes thereof, **291**, 300

Aminopyridinate complexes
preparation, structure and electrochemical behavior of dinuclear cyclooctadiene-chelated Ir(I) complexes with 2-aminopyridinato bridges, **292**, 244

Ammonothermal
synthesis of new Group III fluoride–ammonia adducts in supercritical ammonia: structures of $AlF_3(NH_3)_2$ and $InF_2(NH_2)(NH_3)$, **294**, 200

Anderson structure
hydrothermal synthesis and crystal structure of a new molybdenum(VI) arsenate(III), $Co^{III}(en)_3H_3O[(Co^{II}O_6)Mo^{VI}{}_6O_{18}(As^{III}{}_3O_3)_2]\cdot 2H_2O$, **295**, 244

Antiferromagnetic interactions
syntheses, crystal structures and autoreduction behavior of antiferromagnetically coupled dicopper(II) oximato complexes, **293**, 20

Antimony complexes
synthesis and X-ray characterization of phthalocyaninato(2)-antimony(III) triiodide, **293**, 155

Antisepsis agent
a reversible NO complex of $[Fe^{II}(edampda)]$ and the $[Ni^{II}(edampda)]$ analogue (edampda^{2-} = *N*,*N*′-bis(pyridylmethyl)ethylenediamine-*N*,*N*′-diacetate), **286**, 197

Antitumor activity
[*meso*- and *rac*-1,2-bis(4-fluorophenyl)ethylenediamine]chloro-[sulfinylbis(methane)-S]platinum(II) chloride new water soluble platinum complexes with high anti-breast cancer activities, **285**, 184
synthesis and properties of (diamine)platinum(II) complexes of pyridine carboxylate isomers and their antitumor activity, **292**, 238

Aquation kinetics
synthesis, crystal structure and aquation kinetics of cobalt(III) complex of N-substituted tetra-aza macrocycle: C-*rac*-1,5,8,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane, **285**, 107

Arene complexes
π-arene complexes : 11. Site control affected by steric and electronic factors of ring substituents in σ,π-coordinated bimetallic complexes, **284**, 127

Arsaalkene complexes
on the contrasting reactivity of the phosphaalkene $[(\eta^5\text{-}C_5Me_5)(CO)_2FeP{=}C(NMe_2)_2]$ and the arsaalkene $[(\eta^5\text{-}C_5Me_5)(CO)_2FeAs{=}C(NMe_2)_2]$ towards $[(\eta^5\text{-}C_5H_5)Rh(CO)_2]$, **291**, 311

Arsine complexes
electrospray mass spectrometric investigation of the relative ligand properties of EPh_3 ligands (E = P, As, Sb or Bi) towards Ag^+ and Cu^+, **290**, 213

Aryl complexes
^{13}C *ipso* chemical shifts in the aryl phosphino-thioether complexes $PdX(p\text{-}RC_6H_4)(P,S)$. On resonance effects into the aryl ring, **290**, 95

Aryl halides
Grignard reagent formation from aryl halides. There is no aryl radical intermediate along the dominant reaction channel, **296**, 52

(Aryllithium)chromium complexes
addition of aryl substituents to cyclohexadienyliron electrophiles in the development of routes to C_{12} central building blocks for alkaloid synthesis, **296**, 139

Aryllithium nucleophiles
addition of aryl substituents to cyclohexadienyliron electrophiles in the development of routes to C_{12} central building blocks for alkaloid synthesis, **296**, 139

Aryloxide complexes
synthesis and structures of $TiCl_2(2\text{-}OC_6H_4OCH_3)_2$ and $[TiCl_2(2\text{-}OC_6H_4OCH_3)(\mu\text{-}Cl)]_2$ and stepwise reactions of $[TiCl_2(2\text{-}OC_6H_4OCH_3)(\mu\text{-}Cl)]_2$ with 2-propanol, **293**, 12

Aryl radical
Grignard reagent formation from aryl halides. There is no aryl radical intermediate along the dominant reaction channel, **296**, 52
Asymmetric catalysis
molecular assembly of BINOL–Ti complexes into an active μ_3-oxo titanium catalyst, **296**, 267
synthesis, structure, and reactivity of new chiral ferrocenyl pyrazole ligands, **296**, 183
Asymmetric ligands
new versatile phosphorus-containing ligands—asymmetric $(XPR_2)(YPR'_2)NH$ (X, Y = O, S; R, R′ = Ph, OEt) and their potassium salts: the crystal and molecular structures of $(SPPh_2)[OP(OEt)_2]NH$, $K[(SPPh_2)\{OP(OEt)_2\}N]\cdot H_2O$ and $(OPPh_2)[OP(OEt)_2]NH\cdot 1/2HCl\cdot 1/4H_2O$, **287**, 61
Asymmetric reaction
synthetic reactions using organometallics in water. Aldol and allylation reactions catalyzed by Lewis acid–surfactant-combined catalysts/Brønsted acids systems, **296**, 158
Asymmetric reduction
stereoselective reduction of ketones using extracoordinate silicon: C_2-symmetric ligands, **296**, 208
Axial reactivity
steric effect on the axial substitution of bis(dimethyl sulphoxide)phthalocyaninatoiron(II) by substituted imidazoles, **295**, 200
Aza-crown ether complexes
synthesis and silver(I) coordination of N-functionalized aza-crown ethers with pendant aromatic carbocyclic or heterocyclic side-arms, **285**, 1
Azacryptate complexes
crystal structure and characterization of an unexpected dinuclear Mn(II)Mn(III) azacryptate, **287**, 14
Azamacrocycle complexes
a new structurally constrained dioxocyclam ligand: synthesis, characterization and molecular structure comparison of its nickel(II) complex with the nickel(II) complex of dioxocyclam, **292**, 254
effects of electronic structure on chiral induction in outer-sphere electron transfer reactions of complexes with the C_3 symmetric ligand, 1,4,7-triazacyclononane-1,4,7-tris[2′(*R*)-2′-propionate](-3), **290**, 197
synthesis of 6-amino-6-(4-aminobenzyl)-1,4,8,11-tetra-azacyclotetradecane, a new pentadentate polyamine ligand, and crystal structure of its Co(III) complex, **288**, 40
synthesis, crystal structure and aquation kinetics of cobalt(III) complex of N-substituted tetra-aza macrocycle: C-*rac*-1,5,8,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane, **285**, 107
synthesis of the {Co[(6-amino-6-(4-aminobenzyl)-1,4,8,11-tetra-azacyclotetradecane)][PO_4]}, a potential phosphate-binding site for molecular recognition, **293**, 245
Azamacrocyclic complexes
3-hydroxypropylation of tetraazamacrocycles and its effect on the stability of divalent metal complexes, **294**, 103
syntheses and structures of nickel(II) and copper(II) complexes with a 14-membered tetraaza macrocyclic ligand, **286**, 67
Azauridine complexes
^{13}C NMR confirmation that $[Ru^{II}(hedta)(6\text{-azauridine})]^-$ has $\eta^2(5,6)$ and not σ(C-5) coordination, **293**, 123
Azides
the preparation and characterisation of a new interpseudohalogen, $ClCS_2N_3$, and $(CS_2N_3)SS(CS_2N_3)$, a novel disulfurdipseudohalide, **294**, 68
Azido carbondisulfide
the preparation and characterisation of a new interpseudohalogen, $ClCS_2N_3$, and $(CS_2N_3)SS(CS_2N_3)$, a novel disulfurdipseudohalide, **294**, 68
Azo polymers
synthesis and characterisation of main-chain oligomeric cyclopalladated azobenzene complexes, **292**, 163

Bactericidal activity
synthesis and properties of hydrophobic $[Mn^{IV}_2(\mu\text{-}O)_3(L)_2]^{2+}$ complexes, derived from alkyl substituted 1,4,7-triazacyclononane ligands, **295**, 189
Barbier-type allylation
Barbier-type allylation of carbonyl derivatives by use of aluminum as an electron pool. Double allylation of carboxylic esters, **296**, 204
Base hydrolysis
base hydrolysis of $[Co(triamine)(diamine)Cl]^{2+}$ and epimerization of $[Co(triamine)(diamine)OH]^{2+}$: studies on resolved cobalt(III) complexes of 3-aza-hexane-1,6-diamine (2,3-tri), **284**, 37
^9Be NMR
interaction of beryllium(II) in aqueous solution with bidentate ligands containing phosphonate groups, **285**, 39
Benzimidazole
preparation of poly(aryleneethynylene) type π-conjugated polymers constituted of 2-alkylbenzimidazole-4,7-diyl units by using palladium catalyzed cross coupling reaction, **296**, 254
Benzo[*b*]thiophene complexes
oxidative addition reactions of benzo[*b*]thiophene and dibenzothiophene with $[Os_3(CO)_{10}(MeCN)_2]$: room-temperature ring opening of benzo[*b*]thiophene at triosmium clusters, **285**, 277
Benzocyclobutenedione ligand
tricarbonylchromium complexes with methoxy substituted benzocyclobutenone and benzocyclobutenedione ligands, **296**, 150
Benzocyclobutenone ligand
tricarbonylchromium complexes with methoxy substituted benzocyclobutenone and benzocyclobutenedione ligands, **296**, 150
Benzotriazole complexes
intervalence transfer properties of the binuclear μ-benzotriazolate- and μ-benzimidazolate-bis{ruthenium(II)/(III)-edta} complexes, **285**, 197
Benzoyloxy spacer
5,10,15,20-tetrakis(α,α,α,α-*o*-pivalamidophenyl)porphinatoiron(II) bearing a covalently linked axial imidazole via *m*-aminobenzoic acid. Synthesis and influence of imidazole basicity on O_2-binding affinity, **295**, 234
Beryllium complexes
interaction of beryllium(II) in aqueous solution with bidentate ligands containing phosphonate groups, **285**, 39
Betaine
characterization and crystal structure of copper(II) complex with betaine having a water-wheel-like structure, **295**, 121
B-Hematin
Mössbauer studies of different types of hematin, **286**, 229
Bicine
solid and solution studies of ternary copper(II) complexes formed by heteroaromatic N bases and the anion of *N*,*N*-bis(2-hydroxyethyl)glycine (bicine), **285**, 326
Bidentate ligands complexes
reactions of $Ti(O\text{-}i\text{-}Pr)Cl_3$ with bidentate ligands containing hydroxy and methoxy groups. The crystal structure of $Ti(O\text{-}i\text{-}Pr)Cl_3(HOCH_2CH_2OCH_3)$ and the ^1H variable-temperature NMR studies of solution structures of $Ti(O\text{-}i\text{-}Pr)Cl_3(HOCH_2CH_2OCH_3)$ and $Ti(O\text{-}i\text{-}Pr)Cl_3(CH_3OCH_2CH_2OCH_3)$, **285**, 254

Bidentate nitrogen ligand complexes
fluctuations between square-planar and tetrahedral coordination geometry with bis(2-benzimidazolyl)alkane ligands. Synthesis, spectroscopic properties and X-ray crystal structure of four representative examples, **290**, 105
Bifunctional complexes
synthesis and characterization of five-coordinate rhenium(V) and technetium(V) mixed ligand bifunctional complexes carrying the SNS/S or the SNN/S donor atom set. Crystal structure of $ReO\{[(C_2H_5)_2NCH_2CH_2N(CH_2CH_2S)_2](p\text{-}H_2N\text{–}PhS)\}$ and $ReO\{[(CH_2)_4NCH_2CH_2NCH_2CH_2S](p\text{-}H_2N\text{–}PhS)\}$, **285**, 97
Bimetallic clusters
relativistic density functional study on the bimetallic cluster $[Pt_3Fe_3(CO)_{15}]^{n-}$ ($n = 0, 1, 2$), **286**, 30
Bimetallic compounds
π-arene complexes : 11. Site control affected by steric and electronic factors of ring substituents in σ,π-coordinated bimetallic complexes, **284**, 127
Binaphthol
molecular assembly of BINOL–Ti complexes into an active μ_3-oxo titanium catalyst, **296**, 267
Binuclear complexes
photochemical reactions of transition metal organyl complexes with olefins: XXII. Photoreactions of pentacarbonyl(μ-$\eta^{5:3}$-4*H*-indenyl)diiron(Fe–Fe) with alkynes — substituent controlled C–H transfer or C–C bond formation, **296**, 127
synthesis and structural characterization of binuclear silver(I) complexes bridged by three polydentate phosphine ligands, **284**, 278
the first binuclear Mo–S cluster compound containing citrate ligands, $K_5(NH_4)[Mo_2O_2(\mu\text{-}S)_2(C_6H_4O_7)_2]\cdot CH_3OH\cdot 5H_2O$, characterized by X-ray single crystal structure determination, **285**, 152
Binucleating ligand complexes
potentiometric and spectroscopic studies of copper(II) complexes of bis(1,4,7-triazacyclononane) ligands containing polymethylene and xylyl linker groups, **292**, 172
Bioctahedral complexes
synthesis and structural characterization of face-sharing bioctahedral complexes containing poly(pyrazolyl)borate ligands: $[HB(Me_2Pz)_3BH][X_3Mo(\mu\text{-}X)_2(\mu\text{-}H)MoTp^*]$ (X = Cl or Br; $Tp^* = HB(Me_2Pz)_3$; Pz = pyrazolyl), **292**, 182
Biological activity
syntheses and anti-*Candida* activity of cobalt(II) complexes of octanedioic acid ($odaH_2$) and nonanedioic acid ($ndaH_2$); X-ray crystal structures of $[Co(phen)_3]oda\cdot 14H_2O$ and $[Co(phen)_3]nda\cdot 11.5H_2O$ (phen = 1,10-phenanthroline), **293**, 160
synthesis and biological activity of *cis*-dichloro mono- and bis(platinum) complexes with *N*-alkyl-ethylenediamine ligands, **290**, 237
synthesis, structural characterization and biological activity of helicin thiosemicarbazone monohydrate and a copper(II) complex of salicylaldehyde thiosemicarbazone, **286**, 134
Bioresorbable materials
manufacturing of specialty materials in supercritical fluid carbon dioxide, **294**, 214
Biphenyl diamine complexes
synthesis and X-ray crystallographic structures of dichloronickel(II) and -palladium(II) complexes containing the biphenyl diamine ligand 6-(2-dimethylaminoethyl)-6,7-dihydro-5-H-dibenz[c,e]azepin, **294**, 28
Bipyridine complexes
coordination chemistry of 2,4′-bipyridine. Synthesis and structures of $Co(2,4'\text{-bipyridine})_3(NO_3)_2(H_2O)$ and $Cd(2,4'\text{-bipyridine})_2(NO_3)_2(H_2O)_2$, **288**, 215
crystal structure and magnetic properties of the single-μ-chloro copper(II) chain $[Cu(bipy)Cl_2]$ (bipy = 2,2′-bipyridine), **284**, 258
kinetics of the photooxidation of tris(bipyridine)iron(II) in chloroform, **291**, 136
Bipyridylamine complexes
the crystal structures and electronic properties of bis(di-2-pyridylamine)copper(II), bis(tetrafluoroborate) and bis[bis-aquabis(di-2-pyridylamine)copper(II)] sulfate heptahydrate, **292**, 57
Bipyridyl complexes
dinuclear complexes of a new bridging ligand containing 2,2′-bipyridyl and dioxolene binding sites: syntheses, electrochemical and electronic spectroscopic properties, **285**, 89
Bipyrimidine complexes
infinite two-dimensional sheet-like array of a 2,2′-bipyrimidine-bis(oxalato)chromate(III) complex: synthesis, crystal structure and magnetic properties of $[K(H_2O)Cr(C_2O_4)_2(bpym)]$, **284**, 8
Birnessite
a new synthetic route to todorokite-type manganese oxides, **294**, 123
Bisferrocenyl cations
effects of β-cyclodextrin inclusion on the kinetics of the oxidation of bisferrocenyl cations by bis(pyridine-2,6-dicarboxylato) cobaltate(III) in aqueous solution: crystal structure of dimethyldi-(1-methylferrocene)ammonium bromide, **291**, 365
Bis-iron tetraene complexes
synthesis and reactivity of acyclic (pentadienyl)iron(1 +) cations: model studies for the preparation of the 8*E*,10*Z*,16*E*,18*E*-tetraene segment of macrolactin A, **296**, 261
Bismuthines
a semi-empirical quantum chemical study (PM3) of structure–reactivity relationships for mono- and binuclear bismuthines, **284**, 167
Bleomycin complexes
synthesis, spectroscopic, redox properties, and DNA cleavage activity of low-spin iron(III) complexes of bleomycin analogues, **294**, 56
Borane anion reactions
mechanistic aspects of boron hydride reactions, **289**, 1
Borane complexes
B-frame-supported bimetallics. Isoelectronic *arachno*-structured $[(PMe_2Ph)_4Pd_2B_8H_{10}]$ and *closo*-structured $[(PMe_3)_4(CO)_2Ir_2B_8H_8]$, **285**, 290
synthesis and comparative reactivity of thiomorpholine–borane: aqueous hydrolysis and oxidation by hypochlorite, **290**, 80
Borane derivatives
mechanochemical reactions in the chemistry of boranes, **289**, 51
Boranes
early versus late transition metals. Electronic structure of *nido*-2-$CpML_nB_4H_8$, $CpML_n = CpTaCl_2$, $CpWH_3$ and CpCo, **289**, 85
NMR as a tool for elucidation of structures and estimation of electron distribution in boranes and their derivatives, **289**, 20
Boride carbides
B and B–C as interstitials in reduced rare earth halides, **289**, 174
Boride clusters
the versatile nature of [ClAu(L–L)AuCl] (L–L = bis-(diphenylphosphino)methane (dppm) or -butane (dppb)) in reactions with $[Rh_2Ru_4(CO)_{16}B]^-$: crystal structures of $[\{Rh_2Ru_4(CO)_{16}B\}_2\{\mu\text{-}Au(dppb)Au\}]$ and $[RhRu_4(CO)_{14}B\text{-}\{Au(dppm)Au\}]$, **289**, 149
Borides
B and B–C as interstitials in reduced rare earth halides, **289**, 174
comparison of the surface chemical reactivity of hafnium diboride and hafnium, **289**, 191

Borolanylbutadiene complexes
iron complexes of dioxoborolanylbutadienes, **290**, 159
Boron
alkyl derivatives of $[B_6H_6]^{2-}$: NMR and vibrational spectra and crystal structure of $(Ph_4P)[B_6H_6CH_2Ph]$, **289**, 70
an approach to *megalo*-boranes. Mixed and multiple cluster fusions involving iridaborane and platinaborane cluster compounds. Crystal structure determinations by conventional and synchrotron methods, **289**, 95
application of the NOE experiment to the analysis of boron hydride derivatives: confirming the assignments of the *pseudocloso*-complex $[1,2\text{-}Ph_2\text{-}3\text{-}\{Cp^*\}\text{-}3,1,2\text{-}IrC_2B_9H_9]$ ($Cp^* = \eta^5\text{-}C_5Me_5$) and the *closo*-compounds 1-Ph-1,2-$C_2B_{10}H_{11}$ and 1-Ph-2-Me-1,2-$C_2B_{10}H_{10}$, **289**, 125
comparison of the surface chemical reactivity of hafnium diboride and hafnium, **289**, 191
the age of chiral deltahedral borane derivatives, **289**, 45
Boron clusters
novel tetraalkyltetraboranes of the type B_4R_4, $B_4H_2R_4$ and $B_4H_4R_4$, **289**, 58
[4-{(*p*-cym)Ru}-5-{$(Ph_3P)_2(CO)Os$}B_4H_8]: the first *nido*-heterobimetallahexaborane cluster, **289**, 134
Boron hydride reactions
mechanistic aspects of boron hydride reactions, **289**, 1
Boronic acid
mechanistic study on the reaction of boronic acids with H-resorcinol, **290**, 222
Boron–nitrogen complexes
the bonding and structure of B–N sandwich complexes, **288**, 17
Brønsted acids
synthetic reactions using organometallics in water. Aldol and allylation reactions catalyzed by Lewis acid–surfactant-combined catalysts/Brønsted acids systems, **296**, 158
Bromide complexes
structural diversity in thallium chemistry: II. Potential pentabromothallate(III) derivatives: examples of incipient bromide ion coordination?, **284**, 215
Buserite
a new synthetic route to todorokite-type manganese oxides, **294**, 123
Butadiene complexes
iron complexes of dioxoborolanylbutadienes, **290**, 159
Butadiyne bridge complexes
synthesis and characterization of a metal coordination polymer consisting of ruthenium(III) β-diketone units linked by butadiyne bridges, **292**, 64

Cadmium complexes
adduct formation between zinc and cadmium oxydiacetates and imidazol. Crystal structures of $[Zn(Him)_2(oda)]_n$ and $[Cd_2(Him)_4(oda)_2(H_2O)]_2\cdot 6H_2O$ (Him = imidazol, oda = oxydiacetate), **284**, 49
cadmium(II) complexes with [(4-methylphenyl)sulfonyl]-2-pyridyl-amine: crystal structures of bis{[(4-methylphenyl)-sulfonyl]-2-pyridyl-amide}cadmium(II); aquo-bis(pyridine) bis{[(4-methylphenyl)sulfonyl]-2-pyridyl-amide}cadmium(II) and 1,10-phenanthroline bis{[(4-methylphenyl)sulfonyl]-2-pyridyl-amide}cadmium(II), **294**, 87
$[Cd_2(4,4'\text{-bipy})_2(H_2O)_3(SO_4)_2]\cdot 3H_2O$: a novel three-dimensional coordination polymer with nonlinear optical properties, **294**, 95
coordination chemistry of 2,4′-bipyridine. Synthesis and structures of Co(2,4′-bipyridine)$_2(NO_3)_2(H_2O)$ and Cd(2,4′-bipyridine)$_2(NO_3)_2(H_2O)_2$, **288**, 215
macrocyclic complexes derived from the cyclocondensation of sodium 2,6-diformyl-4-substituted phenolates with *N*,*N*-bis(2-aminoethyl)-2-hydroxybenzylamine and in situ transmetallation. An anticipated dinuclear Cd(II) pendant-arm macrocyclic complex and an unexpected mononuclear Fe(III) macrocyclic complex, **294**, 1
metal(II) complexes with monodentate S and N ligands as structural models for zinc–sulfur DNA-binding proteins, **285**, 262
synthesis and structure of a heptacoordinated cadmium(II) complex, **288**, 118
vibrational spectra of the series CdX_2(18-crown-6) and CdXY(18-crown-6) (X or Y = Cl, Br or I), **292**, 108
Cage complexes
a *cyclo*-tetraphosphoxane cage complex of molybdenum(0) and palladium(II), **287**, 223
new capping agents for oximehydrazonate clathrochelates: sterically controlled synthesis, structural characterization and intramolecular reactions, **284**, 180
Calcium
crystal structures of two new heptamolybdates and of a pyrazole incorporating a γ-octamolybdate anion, **295**, 106
Calix(4)arene complexes
the first crystal structure of a complex from a lower rim calix(4)arene alicyclic amino derivative: {tetrakis[1-(ethyl)-piperidinium]oxyl}*p*-tert-butylcalix(4)arene-bis-[tetrachloro-aurate(III)]dichloro dihydrate, **285**, 142
Calorimetry
pyridines from allyl cobaloximes: a new decomposition pathway, **288**, 112
Carbamoyl complexes
synthesis of bis-(acyl, carbamoyl or alkoxycarbonyl) iron monomer or dimer complexes by reaction of oxalyl chloride with various tetracarbonyl ferrates: $\{Fe[C(O)R](CO)_4\}^-$, **291**, 166
Carbazole complexes
synthesis, characterization and optical properties of novel carbazole–ruthenium compounds: carbazole as a new building block?, **288**, 134
Carbene complexes
synthesis of *N*-allyl-2,3-dihydrobenzoxazol-2-ylidene complexes of chromium(0) and tungsten(0), **288**, 47
synthesis of the first chiral phosphorous chelate aminophosphino–carbene complex and its applications in stereoselective addition reactions, **296**, 236
Carbogallation
β-silylethenylation of phenol with silylethyne using gallium trichloride, **296**, 67
Carbohydrate complexes
binding ability of aldaric acid toward metal(II). X-ray study and solution state investigation on Cu(II)-galactaric acid system and its 2,2′-bypiridine adduct, **292**, 189
Carbohydrate ligands
chiral phosphine ligands derived from sugars. 15. Synthesis, structure and spectroscopic studies of platinum(II) complexes with chiral phosphine substituted carbohydrate ligands, **290**, 121
Carbometallation
β-silylethenylation of phenol with silylethyne using gallium trichloride, **296**, 67
Carbonate complexes
a new carbonate bridged dinuclear zinc complex with tripodal amine ligands, **288**, 206
Carbon dioxide
manufacturing of specialty materials in supercritical fluid carbon dioxide, **294**, 214
Carbon monoxide
palladium(II) complexes with hemilabile etherdiphos ligands in the alternating copolymerization of carbon monoxide with olefins, **296**, 103

Carbon-13 study
^{13}C *ipso* chemical shifts in the aryl phosphino-thioether complexes $PdX(p\text{-}RC_6H_4)(P,S)$. On resonance effects into the aryl ring, **290**, 95
Carbonylation
$Ni(COD)_2$: an efficient promoter in the construction of 5-cyclopent-2-enone derivatives from allyl halides, alkynes and carbon monoxide, **296**, 94
P^N versus N^N ligands for the palladium-catalyzed alternating copolymerization of styrene and carbon monoxide, **296**, 45
Carbonylative lactonization
new ferracyclic π-allyl and η^3-pentadienyl complexes by carbonylative lactonization of unsaturated diols, **291**, 91
Carbonyl complexes
absence of 1,3-metallotropic shifts in $M(CO)_5$ complexes of pyrimidine and 1,3,5-triazine (M = W, Mo, Cr), **292**, 225
a large C_6D_6 solvent effect on the ^1H NMR shifts of pyrazines coordinated to the $M(CO)_5$ series (M = W, Mo, Cr), **285**, 157
a molybdenum(0) complex of bis(hydroxymethyl)phenylphosphine, **287**, 209
carbosilane dendrimers functionalized with $AuFe_3$ clusters, **291**, 247
comparing structures and reactivity in analogous Fe and Ru complexes. (^{i}Pr-DAB)$Fe(CO)_2I_2$ and (^{i}Pr-DAB)FeI_2: a perfectly reversible CO-carrier system. (R-DAB = *N*,*N*′-R_2-1,4-diaza-1,3-butadiene), **291**, 438
iron carbonyls with bulky thiolate ligands: crystal structures of $[Fe_2(CO)_6(\mu\text{-}SC_6H_2{}^iPr_3\text{-}2,4,6)_2]$ and $(C_6H_2{}^iPr_3\text{-}2,4,6)_2S_2$, **284**, 14
isocyanide derivatives of manganese cobalt nonacarbonyl: synthesis and characterization, **288**, 159
new synthesis of $[RuH(CO)(PPh_3)_2L_2][BF_4]$ [L_2 = 2py, 2(4-Mepy) or bpy] and X-ray crystal structure of $[RuH\text{-}(CO)(PPh_3)_2(4\text{-}Mepy)_2][BF_4]$, **285**, 318
nucleophilic substitutions of iodide with bromides in the $Fe(CO)_3(PMe_3)_2MeI$ complex, **288**, 200
organometallic radiopharmaceuticals: rhenium(I) carbonyl complexes of natural bile acids and derivatives, **286**, 46
oxidative addition reactions of benzo[*b*]thiophene and dibenzothiophene with $[Os_3(CO)_{10}(MeCN)_2]$: room-temperature ring opening of benzo[*b*]thiophene at triosmium clusters, **285**, 277
reactions of σ-π alkenyl complexes $[Fe_2(CO)_6(\mu\text{-alkenyl})(\mu\text{-}PPh_2)]$ with bis(diphenylphosphino)methane (dppm); carbonyl substitution, migratory carbonyl insertion, phosphorus–carbon bond formation and cleavage, proton shift and alkenyl isomerisation reactions, **291**, 178
reactions of $Cp'(CO)_2MnPPh_2H$ with CH_3COCl and $CH_3S(O)_2Cl$ in THF/triethylamine: evidence of the first complex stabilization of a phosphorus homolog of the sulfonamides, **288**, 101
'ship-in-bottle' synthesis of platinum carbonyl clusters in titania- or zirconia-modified mesoporous channels of FSM-16. Structural characterization and catalysis in CO hydrogenation, **294**, 281
solution thermochemical study of ligand substitution reactions of hybrid alkyl/fluoroalkoxy phosphorus ligands in the $L_2Fe(CO)_3$ system, **291**, 32
syntheses, spectroscopic properties and photochemistry of $[Re(R)(CO)_3(4,4'\text{-}Me_2\text{-}bpy)]$ (R = alkyl) complexes, **284**, 61
synthesis and crystal structure of $[Re(CO)_3(\mu\text{-}CH_3CO_2)(THF)]_4$, **284**, 304
synthesis of hexametallic $CH_2\{(\eta^5\text{-}C_5H_4)Fe(CO)_2SnPh_2\text{-}(\eta^5\text{-}C_5H_5)Fe(CO)_2\}_2$, **291**, 252
the relationship between surface derivatization and the physical properties of bisarylphosphinidenetriironnonacarbonyl clusters, **294**, 140
transformation of the carboxyl group adjacent to the $(\eta^4$-diene)$Fe(CO)_3$ moiety: a convenient route to $(\eta^4$-diene)$Fe(CO)_3$ complexes containing α,β- and β,γ-unsaturated carbonyl groups, and 1,3-dicarbonyl groups, **291**, 231
Carbonylnitronato complexes
metal(II) complexes of the carbonylnitronate ligands $O_2NCHC(O)R^-$ (R = Ph, OMe or OEt) and X-ray structure of $(DMANH)[Ni(O_2NCHC(O)Ph)_3]$ (DMAN = 1,8-bis(dimethylamino)naphthalene), **285**, 18
Carboranes
conversion of site-specific deuterated *closo*-1,2- and 1,7-$C_2B_{10}H_{12}$ to deuterated *nido*-7,8- and 7,9-$[C_2B_9H_{12}]^-$ ions, respectively; mechanistic inferences. Comparisons of GIAO-NMR generated $[C_2B_9H_{12}]^-$ chemical shifts with experimental values, and consideration of two principal isomers of the 7,8-$[C_2B_9H_{12}]^-$ ion by comparison of geometry optimization data, NMR chemical shift data, and NMR coupling data to respective experimental data, **289**, 11
synthesis and characterization of the anionic fluorocarboranes 6,7,8,9,10-$CB_9H_5F_5^-$, 6,7,8,9-$CB_9H_4F_4$-10-OH^-, and 6,7,8,9-$CB_9H_4F_4$-10-$NHCOCH_3^-$, **289**, 76
Carbosilane dendrimers
carbosilane dendrimers functionalized with $AuFe_3$ clusters, **291**, 247
Carboxylate cluster complexes
variations in capping the $Zr_6O_4(OH)_4$ cluster core: X-ray structure analyses of $[Zr_6(OH)_4O_4(OOC\text{–}CH\text{=}CH_2)_{10}]_2(\mu\text{-}OOC\text{–}CH\text{=}CH_2)_4$ and $Zr_6(OH)_4O_4(OOCR)_{12}(PrOH)$ (R = Ph, CMe = CH_2), **284**, 1
Carboxylate complexes
a dinuclear copper(II) complex involving monoatomic *O*-carboxylate bridging and Cu–S(thioether) bonds: $[Cu(tda)(phen)]_2 \cdot H_2tda$ (tda = thiodiacetate, phen = phenanthroline), **286**, 74
magneto–structural correlation of dimeric copper(II) carboxylates with pyridyl-substituted nitronyl nitroxides, **293**, 30
mixed-valence states of iron long-chain carboxylate complexes, **291**, 127
Carboxylato complexes
facile synthesis and structural properties of (diamine)-tetracarboxylatoplatinum(IV) complexes, **292**, 52
Carboxylic acid complexes
spectroscopic study of lanthanide(III) complexes with carboxylic acids, **293**, 167
Catalysis
kinetics of the molybdate and tungstate catalyzed oxidation of iodide by hydrogen peroxide, **288**, 229
synthesis of chiral diphosphite ligands and comparison of their rhodium and platinum complexes in the asymmetric hydroformylation of styrene, **296**, 222
Catalysts
hydrothermal synthesis and crystal structure of a new molybdenum(VI) arsenate(III), $Co^{III}(en)_3H_3O[(Co^{II}O_6)Mo^{VI}{}_6O_{18}(As^{III}{}_3O_3)_2]\cdot 2H_2O$, **295**, 244
optically active β-ketoiminato cobalt(II) complexes as highly efficient catalysts in enantioselective borohydride reduction of ketones, imines, and α,β-unsaturated carboxylates, **296**, 86
Catalytic cyclization
triphenylsilanolato complexes of Group 4 metals: syntheses, X-ray crystal structures and catalytic cyclization reaction of 1,5-hexadiene, **286**, 14
Catalytic hydrogenation
catalytic hydrogenation and deuteration of phospholipid model membranes with a water-soluble chlorotris(1,3,5-triaza-7-phosphaadamantane)rhodium(I) complex catalyst, **293**, 218

Catalytic oxidation
catalytic oxidation of 1,2-diaminobenzene by the dinuclear cobalt(II)–OBISDIEN dioxygen complex, **287**, 193
Catechol dioxygenase model complexes
novel iron(III) complexes with imidazole containing tripodal ligands as model systems for catechol dioxygenases, **291**, 289
Cationic complexes
P ^ N versus N ^ N ligands for the palladium-catalyzed alternating copolymerization of styrene and carbon monoxide, **296**, 45
C–C Bond formation
photochemical reactions of transition metal organyl complexes with olefins: XXII. Photoreactions of pentacarbonyl(μ-$\eta^{5:3}$-4*H*-indenyl)diiron(Fe–Fe) with alkynes—substituent controlled C–H transfer or C–C bond formation, **296**, 127
Cesium complexes
structural aspects of cesium ion selectivity by tetrabenzo-24-crown-8, **290**, 86
Cesium compounds
syntheses and structures of Group 1 mellitate compounds, $K_6[C_6(COO)_6]\cdot 8H_2O$ and $Cs_5[C_6(COO)_6H]\cdot 7H_2O$, **286**, 207
Chain catalysis
organometallic electron reservoir sandwich iron complexes as potential agents for redox and electron transfer chain catalysis, **291**, 1
Chain complexes
structure and magnetic properties of *catena*-(μ-CrO_4-*O*,*O*′)[M^{II}(L)]·$5H_2O$ (M = Ni^{II}, Cu^{II}; L = 3,14-dimethyl-2,6,13,17-tetraazatricyclo[14,4,0^{1,18},0^{7,12}]docosane), **293**, 100
Chalcogenide complexes
synthesis and characterisation of mono- and binuclear palladium and platinum complexes with organochalcogenides, **285**, 190
synthesis and structural characterization of two new Zintl $[M_7As_3Q_{13}]^{4-}$ anions (M = Cu, Ag; Q = Se, Te) containing distorted cubane $[M_7Q]$ clusters, **288**, 1
Chalcogenoimido ligands
new versatile phosphorus-containing ligands—asymmetric $(XPR_2)(YPR'_2)NH$ (X, Y = O, S; R, R′ = Ph, OEt) and their potassium salts: the crystal and molecular structures of $(SPPh_2)[OP(OEt)_2]NH$, $K[(SPPh_2)\{OP(OEt)_2\}N]\cdot H_2O$ and $(OPPh_2)[OP(OEt)_2]NH\cdot 1/2HCl\cdot 1/4H_2O$, **287**, 61
Charge density wave
synthetic control of intrinsic ground-state defects in a mixed valence quasi-one-dimensional Pt halide chain, **294**, 275
Charge transfer
electronic spectra of 1,2-diiminetricarbonylrhenium(I)chloride complexes with imidazole derivatives as ligands, **288**, 150
preparation spectroscopy and conductivity of [Pt(SS)(NN)]-type complexes and their iodine oxidized complexes, **284**, 112
Charge transfer sensitizer
synthesis and photophysical properties of *trans*-dithiocyanato bis(4,4′-dicarboxylic acid-2,2′-bipyridine) ruthenium(II) charge transfer sensitizer, **296**, 250
Chelate complexes
chemoselective synthesis of (allyloxy)carbene chelate complexes of iron, **291**, 449
Chelating agents
new chelating agents suitable for the treatment of iron overload, **291**, 238
Chelating ligands
X-ray crystal structures of copper(II) nitrate complexes of two isomeric N,O-chelating ligands, **284**, 273
Chemical vapour deposition
preliminary studies of $Ni(dmen)_2(CF_3CO_2)_2$, $Ni(deen)_2(CF_3CO_2)_2$ and $Ni(en)(CF_3C(O)CHC(O)CF_3)_2$ as precursors for metalorganic chemical vapour deposition: structures, thermal behaviours and vapour pressures, **288**, 233
Chiral complexes
hydrogen bonding in *cis*-bis(L-alaninato)copper(II): a single crystal neutron diffraction study, **288**, 29
the synthesis and structure of some, *rac*–*trans*-, *meso*–*trans*- and (*S*,*S*)-*trans*- Co(III) complexes with Me_6-[14]-diene-(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene), **292**, 73
Chirality
the age of chiral deltahedral borane derivatives, **289**, 45
Chiral ligands
aminophosphanes derived from chiral β-aminoalcohols and carbonyliron complexes thereof, **291**, 300
Chiral phosphine complexes
formation of donor–acceptor Fe(0)–Hg(II) bond in separation and stabilization of optically active iron(0) phosphine complexes. Absolute configuration of (+)-(*R*)-$(CO)_4$Fe(μ-EtPhPpy)$HgCl_2$, **293**, 106
Chlorinated complexes
synthesis and characterization of a series of hexadecachlorinated phthalocyanines, **293**, 80
Chlorinated hydrocarbons
chemistry of polyfunctional molecules : CXXXI. Crystal structure and preparation of [bis(diphenyl–methyl–phosphino)iminium] tetraphenyloborate, $[(C_6H_5)_2(CH_3)P{=}N{=}P(CH_3)(C_6H_5)_2]^+[B(C_6H_5)_4]^-$, **284**, 288
Chlorination
different chlorination modes of oximes: chlorination of salicylaldoxime coordinated to platinum, **285**, 116
Chloro complexes
crystal structure and magnetic properties of the single-μ-chloro copper(II) chain $[Cu(bipy)Cl_2]$ (bipy = 2,2′-bipyridine), **284**, 258
Chromium complexes
diastereoselective *ortho*-lithiation of tricarbonylchromium complexes of chiral benzaldehyde acetals, **296**, 80
hexadentate chromium(III) complexes containing unsymmetrical edta-type ligands: Part IV. Crystal structures of $(-)_{589}$-*trans*(O_5)-Na[Cr(ed3ap)]·$3H_2O$ and Na[Cr(*u*-eddadp)]·$3H_2O$ and CD spectra correlation: structural parameters of [Cr(edta-type)]$^-$ complexes and their octahedral distortion in relation to the structure of the ligand, **292**, 16
infinite two-dimensional sheet-like array of a 2,2′-bipyrimidine-bis(oxalato)chromate(III) complex: synthesis, crystal structure and magnetic properties of $[K(H_2O)Cr(C_2O_4)_2(bpym)]$, **284**, 8
intramolecular coupling of conjugated enynes with cyclopropylcarbene–chromium complexes; a complex reaction pathway, **296**, 273
nitrite ion-promoted hydration of acetonitrile by the bis(μ-hydroxo)bis{tris(2-pyridylmethyl)amine}chromium(III)} ion, **288**, 211
solvatochromism in ligand-field absorption bands and solvent-dependent deuteron nuclear magnetic resonance spectra of $[CrF_3$(tri(2-pyridyl)amine)]: implication of trigonal splitting for the first spin-allowed d–d band and evaluation of π bonding peculiarities, **295**, 84
synthesis of *N*-allyl-2,3-dihydrobenzoxazol-2-ylidene complexes of chromium(0) and tungsten(0), **288**, 47
the crystal structure and doublet emission properties of *trans*-$[Cr(NH_3)_2$(cyclam)$](ClO_4)Cl_2$, **285**, 313
C–H Transfer
photochemical reactions of transition metal organyl complexes with olefins: XXII. Photoreactions of pentacarbonyl(μ-$\eta^{5:3}$-4*H*-indenyl)diiron(Fe–Fe) with alkynes—substituent controlled C–H transfer or C–C bond formation, **296**, 127
Citrate complexes
the first binuclear Mo–S cluster compound containing citrate ligands, $K_5(NH_4)[Mo_2O_2(\mu\text{-}S)_2(C_6H_4O_7)_2]\cdot CH_3OH\cdot 5H_2O$,

characterized by X-ray single crystal structure determination, **285**, 152
Clathrochelate complexes
new capping agents for oximehydrazonate clathrochelates: sterically controlled synthesis, structural characterization and intramolecular reactions, **284**, 180
Cluster complexes
about the synthesis of the prismane $[NEt_4]_2[Fe_6S_6I_6]$, **284**, 296
a novel copper(I) cluster stabilized by a dithiolate ligand having a conjugated TTF moiety, **285**, 332
a novel oligomerization versus metal chelation: synthesis and crystal structure of pentakis[di-μ-(diisopropylaminoethanethiolato)nickel], **294**, 83
carbosilane dendrimers functionalized with $AuFe_3$ clusters, **291**, 247
cluster synthesis by the reactions of $[Cp'_2Fe_2S_4]$ with transition-metal complexes ($Cp' = C_5Me_5$, 1,3-$C_5H_3(SiMe_3)_2$), **291**, 395
dipyridylamine-ligated octamolybdate clusters aggregated into extended ribbons and layers via hydrogen bonding, **292**, 131
lattice effects in the Mössbauer spectra of salts of $[Fe_4S_4\{S(CH_2)_nOH\}_4]^{2-}$. Crystal structures of $[PPh_4]_2[Fe_4S_4\{S(CH_2)_nOH\}_4]$ ($n = 2$, 3 and 4), **291**, 101
preparation of cyclopentadienyl molybdenum clusters linked by a cyclotetradeca-1,8-diyne ligand, **294**, 232
protonation and substitution reactions of Fe–S 'basket' clusters including extracted FeMo-cofactor of nitrogenase, **291**, 212
'ship-in-bottle' synthesis of platinum carbonyl clusters in titania- or zirconia-modified mesoporous channels of FSM-16. Structural characterization and catalysis in CO hydrogenation, **294**, 281
spontaneous formation of trinickel clusters containing μ^3-imido bridges from 2-nitrophenols and $Ni(PMe_3)_4$, **287**, 199
syntheses and X-ray structures of mixed-metal gold phosphine clusters including an example having a highly asymmetric Re_2Au_2 skeleton, **294**, 255
syntheses, crystal structures and magnetic properties of heterometallic copper–lanthanide clusters $[Cu_{12}Ln_6(\mu_3\text{-}OH)_{24}(\mu\text{-}O_2CR)_{12}(H_2O)_{18}(\mu_{12}\text{-}ClO_4)]^{5+}$ (Ln = La, Nd; R = CH_2Cl, CCl_3), **293**, 129
syntheses, crystal structures and nonlinear optical properties of heteronuclear clusters $[MS_4Cu_4(SCN)_2(NC_5H_5)_6]$ (M = W, Mo), **294**, 73
synthesis and structural characterization of $[NEt_4][Fe_3(\mu_3\text{-}O)(\mu_3\text{-}AuPPh_3)(\mu\text{-}CO)_3(CO)_6]$, the new $[Au_6(\mu_3\text{-}S)_2(PPh_3)_6][Fe_3(\mu_3\text{-}S)(\mu\text{-}AuPPh_3)(CO)_9]_2$ and $[Au_6(\mu_3\text{-}S)_2(PPh_3)_6][Fe_5(\mu_3\text{-}S)_2(CO)_{14}]$ ionic solids containing assemblages of cluster-cations and cluster-anions, **291**, 372
synthesis and structural characterization of two new Zintl $[M_7As_3Q_{13}]^{4-}$ anions (M = Cu, Ag; Q = Se, Te) containing distorted cubane $[M_7Q]$ clusters, **288**, 1
synthesis, structure, and redox properties of the sulfido bridged tetranuclear cluster complex $[(Ph_3P)_2N][Fe_3(CO)_9\{\mu_3\text{-}SFeCp(CO)_2\}]$, **286**, 142
the relationship between surface derivatization and the physical properties of bisarylphosphinidenetriironnonacarbonyl clusters, **294**, 140
Cluster growth reactions
mechanistic aspects of boron hydride reactions, **289**, 1
Cluster structures
cluster synthesis by the reactions of $[Cp'_2Fe_2S_4]$ with transition-metal complexes ($Cp' = C_5Me_5$, 1,3-$C_5H_3(SiMe_3)_2$), **291**, 395
^{13}C NMR
^{13}C NMR confirmation that $[Ru^{II}(hedta)(6\text{-azauridine})]^-$ has $\eta^2(5,6)$ and not $\sigma(C\text{-}5)$ coordination, **293**, 123
Cobalamins
vitamin B_{12}–transport protein interaction: electrochemistry of aquo- and glutathionyl-cobalamins adsorbed on carbon electrodes; role of the nucleotide chain, **292**, 204
Cobaloxime complexes
reactions of cobaloximes with vinyl allenes as a route to 1-cobaloxime substituted 1,3-dienyl complexes, **296**, 13
Cobaloximes
pyridines from allyl cobaloximes: a new decomposition pathway, **288**, 112
Cobalt complexes
anionopentaaminecobalt(III) complexes with polyamine ligands: Part 30. The base hydrolysis kinetics and structure determination of some $[CoCl(N)_5]^{2+}$ complexes containing pyridine ligands, **288**, 181
base hydrolysis of $[Co(triamine)(diamine)Cl]^{2+}$ and epimerization of $[Co(triamine)(diamine)OH]^{2+}$: studies on resolved cobalt(III) complexes of 3-aza-hexane-1,6-diamine (2,3-tri), **284**, 37
carbon–oxygen bond cleavage: a new reaction mode for amides bonded through oxygen to cobalt(III), **287**, 1
catalytic oxidation of 1,2-diaminobenzene by the dinuclear cobalt(II)–OBISDIEN dioxygen complex, **287**, 193
coordination chemistry of 2,4′-bipyridine. Synthesis and structures of Co(2,4′-bipyridine)$_2$$(NO_3)_2(H_2O)$ and Cd(2,4′-bipyridine)$_2$$(NO_3)_2(H_2O)_2$, **288**, 215
coordination of *ortho* and *para* quinones and oxidation reduction processes in trimethylphosphine complexes of cobalt and nickel, **287**, 167
electrochemical synthesis of cobalt, nickel, copper, zinc and cadmium complexes with *N*[(2-hydroxyphenyl)methylidine]-*N*′-tosylbenzene-1,2-diamine. The crystal structures of {(1, 10-phenanthroline)[*N*-(2-oxophenyl)methylidine]-*N*-tosylbenzene-1,2-diaminato}nickel(II) and {(1,10-phenanthroline)[*N* - (2 - oxophenyl)methylidine] - *N*′ - tosylbenzene-1,2-diaminato}copper(II), **295**, 39
equilibrium studies of reactions of 4-chloro-1,2-diaminobenzene, 4-methyl-1,2-diaminobenzene, 4-methoxy-1,2-diaminobenzene and 4,5-dimethyl-1,2-diaminobenzene with dinuclear cobalt(II) 1,4,7,13,16,19-hexaaza-10,22-dioxacyclotetracosane, **284**, 30
facile and stereoselective condensation of acetone with ammonia ligands on cobalt(III): structure of a N-bonded cyanate complex containing the 2-methyl-2-amino-4-imino-pentane ligand, **290**, 133
hydrothermal synthesis and crystal structure of a new molybdenum(VI) arsenate(III), $Co^{III}(en)_3H_3O[(Co^{II}O_6)Mo^{VI}{}_6O_{18}(As^{III}{}_3O_3)_2]\cdot 2H_2O$, **295**, 244
isocyanide derivatives of manganese cobalt nonacarbonyl: synthesis and characterization, **288**, 159
metal(II) complexes with monodentate S and N ligands as structural models for zinc–sulfur DNA-binding proteins, **285**, 262
molecular structure of $[PPh_4]_2[Co(C_3S_5)_2]$ [$C_3S_5^{2-}$ = 4,5-disulfanyl-1,3-dithiole-2-thionate(2-)] and spectroscopic and electrical properties of its oxidized species, **295**, 239
$[(NH_3)_5CoOP(H)(O)OCo(NH_3)_5]^{4+}$ and its conjugate acid; evidence for the P^+–O^- description of a terminal P–O bond, **294**, 14
optically active β-ketoiminato cobalt(II) complexes as highly efficient catalysts in enantioselective borohydride reduction of ketones, imines, and α,β-unsaturated carboxylates, **296**, 86
oxidation state dependent *cis*/*trans* isomerization in ML_2 complexes of the heteroscorpionate ligand, (2-hydroxyphenyl)bis(pyrazolyl)methane and its derivatives with iron and cobalt, **286**, 80
reaction of formaldehyde with cyano(ethylenediamine)-(glycylglycinato)cobalt(III): crystal structures of cyano-(ethylenediamine) [*N*-(4-carbonyl-4-hydroxymethyl-oxazo-

lidine)glycinato]cobalt(III)·3.5H_2O and dicyanobis(ethylenediamine) {μ - *N* - [*N* - (((methylimino)methyl)carbonyl) - α - methylserinato)glycylglycinato}dicobalt(III) tetrahydrate, **292**, 96
receptors based on 2,2′:6′,2″-terpyridine fragments containing peripheral amino groups, **292**, 28
structural characterization of a new ligand mode of 2,6-diacetylpyridine bis(semicarbazone), H_2daps, **292**, 121
syntheses and anti-*Candida* activity of cobalt(II) complexes of octanedioic acid (odaH_2) and nonanedioic acid (ndaH_2); X-ray crystal structures of [Co(phen)$_3$]oda·14H_2O and [Co(phen)$_3$]nda·11.5H_2O (phen = 1,10-phenanthroline), **293**, 160
synthesis, crystal structure and aquation kinetics of cobalt(III) complex of N-substituted tetra-aza macrocycle: C-*rac*-1,5,8,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane, **285**, 107
synthesis of 6-amino-6-(4-aminobenzyl)-1,4,8,11-tetra-azacyclotetradecane, a new pentadentate polyamine ligand, and crystal structure of its Co(III) complex, **288**, 40
synthesis of the {Co[(6-amino-6-(4-aminobenzyl)-1,4,8,11-tetra-azacyclotetradecane)][PO_4]}, a potential phosphate-binding site for molecular recognition, **293**, 245
the first X-ray crystal structures of cobalt complexes containing monodentate and bridging ethylenediamine ligands, **288**, 53
the kinetics of the sequential hydrolysis reactions of the *xxm*-[$CoCl_2$(dien)(py)]$^+$ cation and the crystal structure of the $ZnCl_4^{2-}$·2H_2O salt, **292**, 198
the methylcobalt(III) complex of a tetrapodal pentadentate amine ligand, 2,6-bis(1′,3′-diamino-2′-methyl-prop-2′-yl)pyridine, **286**, 98
the syntheses, spectra and structures of five-coordinate cobalt(II) complexes of pyrazolyl-containing ligands, **284**, 266
the synthesis and structure of some, *rac*–*trans*-, *meso*–*trans*- and (*S*,*S*)-*trans*- Co(III) complexes with Me_6-[14]-diene-(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene), **292**, 73
tridentate N_3 capping coordination behaviour of potentially multi-site coordinating cyclotriphosphazenes: synthesis and spectroscopic studies of 2,2-spiro(1,3-propanediamino)-4,4,6,6-tetrakis(3,5-dimethylpyrazolyl)cyclotriphosphazene, 2,2-diamino-4,4,6,6-tetrakis(3,5-dimethylpyrazolyl)-cyclotriphosphazene and their mononuclear complexes; X-ray structure of ATPCTP·$CoCl_2$, **286**, 127

Condensation
preparation and characterization of various *N*-substituted-2-aza-[3]-ferrocenophanes and their chemical and electrochemical properties, **296**, 176

Conductivity
polyiodides of transition metal trischelate cations: syntheses, structures, and spectral and electrical conductivity studies of [Mn(phen)$_3$](I_3)$_2$ and [Mn(bpy)$_3$](I_3)$_{1.5}$(I_8)$_{0.25}$, **284**, 158

Conjugated enynes
intramolecular coupling of conjugated enynes with cyclopropylcarbene–chromium complexes; a complex reaction pathway, **296**, 273

π-Conjugated polymers
preparation of poly(aryleneethynylene) type π-conjugated polymers constituted of 2-alkylbenzimidazole-4,7-diyl units by using palladium catalyzed cross coupling reaction, **296**, 254

Cooley's anemia
new chelating agents suitable for the treatment of iron overload, **291**, 238

Coordination chemistry
the preparation and coordination chemistry of $^tPr_2P(E)NHP(E')^tPr_2$ (E, E′ = Se; E = Se, E′ = S; E = S, E′ = O; E,E′ = O, **290**, 1

Coordination number
high coordination numbers of metal ions in chelate complexes: molecular adducts of scandium tris-α-diketonates in solution, **286**, 175
structural diversity in thallium chemistry: III. The first structurally characterized examples of the pentabromothallate(III) anion in the 1,1,4,4-tetramethylpiperazinium and *N*,*N*′-diethyltriethylenediammonium salts contrasted with the mixed anions in the 1,1,3,3-tetramethylimidazolidinium salt, **285**, 122

Coordination polymer complexes
[Cd_2(4,4′-bipy)$_2$(H_2O)$_3$(SO_4)$_2$]·3H_2O: a novel three-dimensional coordination polymer with nonlinear optical properties, **294**, 95
studies of the construction of coordination polymers using linear pyridyl-donor ligands, **292**, 231

Copolymerization
palladium(II) complexes with hemilabile etherdiphos ligands in the alternating copolymerization of carbon monoxide with olefins, **296**, 103
P ^ N versus N ^ N ligands for the palladium-catalyzed alternating copolymerization of styrene and carbon monoxide, **296**, 45

Copper and silver complexes
electrochemical synthesis of copper and silver pyridine-2-selenolato complexes: crystal structure of (6-trimethylsilylpyridine-2-selenolato)silver(I), [Ag_6(6-Me_3SipySe)$_6$], **284**, 133

Copper complexes
ab initio calculations and X-ray structure determinations of dichloro(*O*-*O*′-*O*″-1,3,5-cyclohexanetriol)copper(II) and bis(*O*-*O*′-*O*″-1,3,5-hexanetriol)nickel(II), **284**, 284
a dinuclear copper(II) complex involving monoatomic *O*-carboxylate bridging and Cu–S(thioether) bonds: [Cu(tda)(phen)]$_2$ · H_2tda (tda = thiodiacetate, phen = phenanthroline), **286**, 74
an alternating copper(II) chain with bridging oxamidato and nitrito ligands: crystal structure and magnetic properties of [$Cu(NO_2)_2CuL]_n$ (L = *N*,*N*′-bis(2-methyl-2-aminopropyl) oxamide), **294**, 8
a new bidimensional compound containing (μ-thiocyanate)(bpy) copper(II) molecules: synthesis, crystal structure and magnetic properties of [Cu(bpy)($NCS)_2]_n$ (bpy = 2,2′-bipyridyl), **286**, 108
a new dioxotetraamine ligand appended with fluorenyl and its copper(II) complex. Synthesis, crystal structure and solution behavior, **295**, 48
a new self-assembled copper(I) complex of 2-thioxo-5H-1,3-dithiolo[4,5-*b*][1,4]dithiepin-6(7*H*)-one showing three-dimensional supramolecular networks, **292**, 117
a novel copper(I) cluster stabilized by a dithiolate ligand having a conjugated TTF moiety, **285**, 332
aromatic carboxylate-controlled self-organization of copper(II)–L-arginine complexes, **286**, 37
binding ability of aldaric acid toward metal(II). X-ray study and solution state investigation on Cu(II)-galactaric acid system and its 2,2′-bypiridine adduct, **292**, 189
bridging properties of disulfoxide ligands: crystal structure of copper(1,3-bis(n-propylsulfinyl)propane) perchlorate, **292**, 144
complexes of copper(II) dihalide with 2,2′-dipyridylamine. X-ray diffraction structures of the [dibromo-bis(dipyam)-copper-(II)]–water (1:2) and di[chloro-bis(dipyam)-copper(II)]diiodide–acetonitrile (1:2) complexes, **290**, 180
coordination complexes of bifunctional compounds: I. Synthesis and properties of bis[5-(2-oxyphenyl)-3-phenyl-1,2,4-oxadiazolyl]copper(II). A fluorescent coordination compound of Cu(II), **292**, 1

coordination compounds with 4-aminopyridine. An unusual in situ formation of a bridging formato compound. The structure of *catena*-poly[bis(4-amino-pyridine)(formato-*O*)-copper(II)-μ-formato(*O*,*O'*)], **287**, 226

copper(I) and copper(II) complexes of the bidentate imidazole/thioether ligand 1-methyl-2-(methylthiomethyl)-1*H*-benzimidazole, **287**, 204

copper(II) and zinc(II) complexes of a macrocyclic bis-(amine–amide–hydroxamate) siderophore analogue. Formation constants and coordination chemistry, **284**, 20

copper(II) complexes with (2-hydroxybenzyl-2-pyridylmethyl)-amine–Hbpa: syntheses, characterization and crystal structures of the ligand and $[Cu(II)(Hbpa)_2](ClO_4)_2{\cdot}2H_2O$, **290**, 207

crystallographic and ESR studies of π–π stacked binuclear copper(II) complexes of 1,3-bis(quinoline-2-carboxamido) and 1,3-bis(isoquinoline-3-carboxamido)propane, **288**, 90

crystal structure and magnetic properties of the single-μ-chloro copper(II) chain $[Cu(bipy)Cl_2]$ (bipy = 2,2'-bipyridine), **284**, 258

crystal structures and magnetic properties of the squarate-O^1,O''-bridged dinuclear copper(II) complexes $[Cu_2(phen)_4(C_4O_4)](CF_3SO_3)_2{\cdot}3H_2O$ ($n=2$) and $[Cu_2(bipy)_4(C_4O_4)](CF_3SO_3)_2{\cdot}6H_2O$ ($n=3$), **287**, 173

Cu–O stretching frequency correlation with phenanthroline pK_a values in mixed copper complexes, **288**, 106

cyanamidonitrate–copper(II) complexes of imidazole ligands: X-ray crystallography and physical investigation, **287**, 186

displacement of triphenylphosphine from $Cu(PPh_3)_2NO_3$ and $Co(PPh_3)_2Cl_2$ by a diselenoimidodiphosphinato ligand. X-ray crystal structure of $(PPh_3)Cu[Ph_2P(Se)NP(Se)Ph_2]$ and $Co[Ph_2P(Se)NP(Se)Ph_2]_2$ containing the novel $CuSe_2P_2N$ and $CoSe_2P_2N$ inorganic metallocycles, **290**, 256

electrochemical synthesis of cobalt, nickel, copper, zinc and cadmium complexes with *N*[(2-hydroxyphenyl)methylidine]-*N'*-tosylbenzene-1,2-diamine. The crystal structures of {(1,10 - phenanthroline)[*N* - (2 - oxophenyl)methylidine]-*N*-tosylbenzene-1,2-diaminato}nickel(II) and {(1,10-phenanthroline)[*N* - (2 - oxophenyl)methylidine] - *N'* - tosylbenzene - 1,2-diaminato}copper(II), **295**, 39

electrospray mass spectrometric investigation of the relative ligand properties of EPh_3 ligands (E = P, As, Sb or Bi) towards Ag^+ and Cu^+, **290**, 213

fluctuations between square-planar and tetrahedral coordination geometry with bis(2-benzimidazolyl)alkane ligands. Synthesis, spectroscopic properties and X-ray crystal structure of four representative examples, **290**, 105

hydrogen bonding in *cis*-bis(L-alaninato)copper(II): a single crystal neutron diffraction study, **288**, 29

interaction between heterobinuclear molecules and nature of the ground spin state in oximato-bridged $[Cu^{II}M^{II}]_2$ bis-binuclear complexes (M = Cu, Ni, Mn): crystal structure of $[Cu(pdmg)Ni(Me_3[12]N_3)(EtOH)](ClO_4)_2$, **288**, 57

magneto–structural correlation of dimeric copper(II) carboxylates with pyridyl-substituted nitronyl nitroxides, **293**, 30

potentiometric and spectroscopic studies of copper(II) complexes of bis(1,4,7-triazacyclononane) ligands containing polymethylene and xylyl linker groups, **292**, 172

solid and solution studies of ternary copper(II) complexes formed by heteroaromatic N bases and the anion of *N*,*N*-bis(2-hydroxyethyl)glycine (bicine), **285**, 326

studies of the construction of coordination polymers using linear pyridyl-donor ligands, **292**, 231

syntheses and structures of nickel(II) and copper(II) complexes with a 14-membered tetraaza macrocyclic ligand, **286**, 67

syntheses, crystal structures and autoreduction behavior of antiferromagnetically coupled dicopper(II) oximato complexes, **293**, 20

syntheses, crystal structures and magnetic properties of heterometallic copper–lanthanide clusters $[Cu_{12}Ln_6(\mu_3\text{-}OH)_{24}(\mu\text{-}O_2CR)_{12}(H_2O)_{18}(\mu_{12}\text{-}ClO_4)]^{5+}$ (Ln = La, Nd; R = CH_2Cl, CCl_3), **293**, 129

syntheses, crystal structures and nonlinear optical properties of heteronuclear clusters $[MS_4Cu_4(SCN)_2(NC_5H_5)_6]$ (M = W, Mo), **294**, 73

synthesis and coordination chemistry of 1-phenyl-1-phospha-4,7-dithiacyclononane, **294**, 170

synthesis and crystal structure of tetranuclear copper(I) and silver(I) complexes bridged by 2-amino-1,3,4-thiadiazole (atdz): $[Cu_4(atdz)_6](ClO_4)_4{\cdot}2CH_3OH$ and $[Ag_4(atdz)_6](ClO_4)_4$, **290**, 153

synthesis, characterization, spectroscopy, magnetism and X-ray structures of red and green Cu(II) chloride adducts with bis(2-benzimidazolyl)propane, **288**, 220

synthesis, crystal structure and magnetic behaviour of the new linear copper(II) tetranuclear compound $[Cu_4(\mu_2\text{-}AcO)_6(AcO)_2(DPQ)_2]$ (DPQ = 2,3 - bis(2 - pyridyl)quinoxaline), **286**, 62

synthesis, crystal structure and magnetic properties of a copper(II)–nitroxide linear chain complex, **287**, 181

synthesis, crystal structure and magnetic properties of *catena*-poly[(2,2'-bipyridylcopper)-di-μ-thiocyanato] and bis(2,2'-bipyridyl)thiocyanatocopper(II) perchlorate, **286**, 169

synthesis of 6-amino-6-(4-aminobenzyl)-1,4,8,11-tetra-azacyclo-tetradecane, a new pentadentate polyamine ligand, and crystal structure of its Co(III) complex, **288**, 40

synthesis, spectroscopic, magnetic properties and X-ray crystal structure of di-μ-hydroxo-bis(μ-perchlorato-*O*,*O'*)-bis-[bis(2-amino-4-methylpyrimidine)copper(II)]: bridging perchlorate results in significant deviation from the predicted magnetic exchange, **286**, 24

synthesis, structural characterization and biological activity of helicin thiosemicarbazone monohydrate and a copper(II) complex of salicylaldehyde thiosemicarbazone, **286**, 134

synthesis, structure and magnetic behavior of a di-μ-chloro-di-μ-imidazolato-tetracopper(II) complex with two dinucleating macrocycles, **287**, 52

synthetic, spectroscopic and X-ray crystallographic structural studies on some copper(II) complexes of the 6-methylpyridine-2-carboxaldehyde Schiff base of *S*-methyldithiocarbazate, **287**, 8

the crystal structures and electronic properties of bis(di-2-pyridylamine)copper(II), bis(tetrafluoroborate) and bis[bis-aquabis(di-2-pyridylamine)copper(II)] sulfate heptahydrate, **292**, 57

the interaction of BMXD and its copper(II) complexes with glycine, aspartic acid, and histidine, **286**, 55

three mononuclear copper complexes of 9-hydroxy-9*H*-fluorene-9-carboxylic acid, **294**, 163

tridentate N_3 capping coordination behaviour of potentially multi-site coordinating cyclotriphosphazenes: synthesis and spectroscopic studies of 2,2-spiro(1,3-propanediamino)-4,4,6,6 - tetrakis(3,5 - dimethylpyrazolyl)cyclotriphosphazene, 2,2-diamino-4,4,6,6-tetrakis(3,5-dimethylpyrazolyl)-cyclotriphosphazene and their mononuclear complexes; X-ray structure of ATPCTP·$CoCl_2$, **286**, 127

triethanolamine complexes of copper, **294**, 153

versatility of the cyanato ligand: structure and magnetic properties of four new copper(II)–cyanato compounds derived from tridentate amines, **286**, 189

X-ray crystal structures of copper(II) nitrate complexes of two isomeric N,O-chelating ligands, **284**, 273

Copper(I) complexes
stability constants of copper(I) and silver(I) complexes with open-chain, macrocyclic and -bicyclic aza-ligands in acetonitrile and comparison with results in dimethylsulfoxide, **286**, 160
Copper(II) complexes
a two-dimensional network constructed via 72-membered heart-shaped macrocycles, a copper(II) complex with 1,3,5-tris-(imidazol-1-ylmethyl)-2,4,6-trimethylbenzene and diethylenetriamine ligands, **295**, 129
characterization and crystal structure of copper(II) complex with betaine having a water-wheel-like structure, **295**, 121
Copper sulfite
spectroscopic and magnetic properties of Chevreul's salt, a mixed valence copper sulfite $Cu_3(SO_3)_2 \cdot 2H_2O$, **295**, 125
Cross-coupling
Kumada couplings catalyzed by nickel on charcoal (Ni/C), **296**, 164
cross-coupling reactions proceeding through η^1- and η^3-propargyl/allenyl–palladium(II) intermediates, **296**, 37
Crown ether complexes
a novel polymeric disilver(I) complex containing both macrocyclic- and linear-coordinate silver(I) centres: $[Ag_2(mn\text{-}15S_2O_3)]_x(ClO_4)_{2x}$ (mn-15S_2O_3 = maleonitriledithia[15]crown-5), **285**, 305
solution state chemistry of scandium(III): a study of scandium-crown ether complexes by ^{45}Sc-NMR spectroscopy, **284**, 71
structural aspects of cesium ion selectivity by tetrabenzo-24-crown-8, **290**, 86
vibrational spectra of the series CdX_2(18-crown-6) and CdXY(18-crown-6) (X or Y = Cl, Br or I), **292**, 108
Crown ethers
$NH_4TlI_4 \cdot 2$(18c6)$\cdot NH_3$, a new type of a potential double-shell lattice clathrate, **293**, 234
Cryptands
stability constants of copper(I) and silver(I) complexes with open-chain, macrocyclic and -bicyclic aza-ligands in acetonitrile and comparison with results in dimethylsulfoxide, **286**, 160
Crystal engineering
hydrothermal assembly and structural characterization of two low-dimensional molybdenum oxide/2,2′-dipyridylamine hybrid materials, **295**, 222
Crystal structure complexes
reactions of Ti(O-*i*-Pr)Cl_3 with bidentate ligands containing hydroxy and methoxy groups. : The crystal structure of Ti(O-*i*-Pr)Cl_3($HOCH_2CH_2OCH_3$) and the ^{1}H variable-temperature NMR studies of solution structures of Ti(O-*i*-Pr)Cl_3($HOCH_2CH_2OCH_3$) and Ti(O-*i*-Pr)Cl_3(CH_3OCH_2-CH_2OCH_3), **285**, 254
Crystal structures
ab initio calculations and X-ray structure determinations of dichloro(*O*-*O*′-*O*″-1,3,5-cyclohexanetriol)copper(II) and bis-(*O*-*O*′-*O*″-1,3,5-hexanetriol)nickel(II), **284**, 284
a *cyclo*-tetraphosphoxane cage complex of molybdenum(0) and palladium(II), **287**, 223
adduct formation between zinc and cadmium oxydiacetates and imidazol. Crystal structures of $[Zn(Him)_2(oda)]_n$ and $[Cd_2(Him)_4(oda)_2(H_2O)]_2 \cdot 6H_2O$ (Him = imidazol, oda = oxydiacetate), **284**, 49
1:1 adducts of triphenyltin chloride with oxovanadium(IV) tetradentate Schiff-base complexes, **293**, 147
a dinuclear copper(II) complex involving monoatomic *O*-carboxylate bridging and Cu–S(thioether) bonds: $[Cu(tda)(phen)]_2 \cdot H_2tda$ (tda = thiodiacetate, phen = phenanthroline), **286**, 74
alkyl derivatives of $[B_6H_6]^{2-}$: NMR and vibrational spectra and crystal structure of $(Ph_4P)[B_6H_6CH_2Ph]$, **289**, 70
aminophosphanes derived from chiral β-aminoalcohols and carbonyliron complexes thereof, **291**, 300
a molybdenum(0) complex of bis(hydroxymethyl)phenylphosphine, **287**, 209
an alternating copper(II) chain with bridging oxamidato and nitrito ligands: crystal structure and magnetic properties of $[Cu(NO_2)_2CuL]_n$ (L = *N*,*N*′-bis(2-methyl-2-aminopropyl) oxamide), **294**, 8
a neutral molecule of I_2 as a bridge for dimerization of iodine iron(III) phthalocyanine, **287**, 101
a new bidimensional compound containing (μ-thiocyanate)(bpy) copper(II) molecules: synthesis, crystal structure and magnetic properties of $[Cu(bpy)(NCS)_2]_n$ (bpy = 2,2′-bipyridyl), **286**, 108
a new carbonate bridged dinuclear zinc complex with tripodal amine ligands, **288**, 206
a new dioxotetraamine ligand appended with fluorenyl and its copper(II) complex. Synthesis, crystal structure and solution behavior, **295**, 48
a new linear polymer chain iron complex; synthesis, crystal and molecular structural characterization of catena-μ-acetato-[*NN*′-ethylenebis(salicylaldiminato)]iron(III), **287**, 109
a new self-assembled copper(I) complex of 2-thioxo-5H-1,3-dithiolo[4,5-*b*][1,4]dithiepin-6(7*H*)-one showing three-dimensional supramolecular networks, **292**, 117
a new structurally constrained dioxocyclam ligand: synthesis, characterization and molecular structure comparison of its nickel(II) complex with the nickel(II) complex of dioxocyclam, **292**, 254
an optically active nickel(II) Schiff base coordination compound *N*,*N*′-(1*R*,2*R*)-(−)-1,2-cyclohexylenebis(2-hydroxyacetophenonylideneiminato)nickel(II), **293**, 239
a novel copper(I) cluster stabilized by a dithiolate ligand having a conjugated TTF moiety, **285**, 332
a novel oligomerization versus metal chelation: synthesis and crystal structure of pentakis[di-μ-(diisopropylaminoethanethiolato)nickel], **294**, 83
a novel polymeric disilver(I) complex containing both macrocyclic- and linear-coordinate silver(I) centres: $[Ag_2(mn\text{-}15S_2O_3)]_x(ClO_4)_{2x}$ (mn-15S_2O_3 = maleonitrile-dithia-[15]crown-5), **285**, 305
π-arene complexes: 11. Site control affected by steric and electronic factors of ring substituents in σ,π-coordinated bimetallic complexes, **284**, 127
aromatic carboxylate-controlled self-organization of copper(II)–L-arginine complexes, **286**, 37
a series of heteroleptic complexes of the type *fac*-$[Mn^{III}L_2]^-$ [H_2L = derivatives of *N*-(2-hydroxybenzyl)glycine or *N*-(5-nitro-2-hydroxybenzyl)sarcosine] possessing unusual Mn(III) co-ordination spheres, **290**, 228
a seven-coordinate manganese(II) complex formed with the single heptadentate ligand *N*,*N*,*N*′-tris(2-pyridylmethyl)-*N*′-(2-salicylideneethyl)ethane-1,2-diamine, **290**, 139
asymmetric compartmental macrocyclic ligands and related mononuclear and hetero-dinuclear complexes with d- and/or f-metal ions, **293**, 178
B and B–C as interstitials in reduced rare earth halides, **289**, 174
bdmpp complexes of Ru(II) as structural analogs for Ru(II)-2,2′:6′,2″-terpyridine species (where bdmpp is 2,6-bis(3,5-dimethyl-*N*-pyrazoyl)pyridine), **286**, 181

B-frame-supported bimetallics. Isoelectronic *arachno*-structured $[(PMe_2Ph)_4Pd_2B_8H_{10}]$ and *closo*-structured $[(PMe_3)_4(CO)_2Ir_2B_8H_8]$, **285**, 290

binding ability of aldaric acid toward metal(II). X-ray study and solution state investigation on Cu(II)-galactaric acid system and its 2,2′-bypiridine adduct, **292**, 189

bisphenolate iron(II) complexes with intramolecularly coordinating nitrogen Lewis bases, **291**, 326

bridging properties of disulfoxide ligands: crystal structure of copper(1,3-bis(n-propylsulfinyl)propane) perchlorate, **292**, 144

cadmium(II) complexes with [(4-methylphenyl)sulfonyl]-2-pyridyl-amine: crystal structures of bis{[(4-methylphenyl)-sulfonyl]-2-pyridyl-amide}cadmium(II); aquo-bis(pyridine) bis{[(4-methylphenyl)sulfonyl]-2-pyridyl-amide}cadmium(II) and 1,10-phenanthroline bis{[(4-methylphenyl)sulfonyl]-2-pyridyl-amide}cadmium(II), **294**, 87

cationic 16-electron half-sandwich ruthenium complexes containing asymmetric diamines: understanding the stability and reactivity of coordinatively unsaturated two-legged piano stool complexes, **286**, 114

$[Cd_2(4,4'\text{-bipy})_2(H_2O)_3(SO_4)_2]\cdot 3H_2O$: a novel three-dimensional coordination polymer with nonlinear optical properties, **294**, 95

chelating and ligand-transfer studies of the complexes *cis*-$[Mn(CO)_4(SeR)_2]^-$: crystal structures of heterotrinuclear $[(CO)_4Mn(\mu\text{-SeMe})_2Ni(\mu\text{-SeMe})_2Mn(CO)_4]$ with a distorted square planar Ni^{II}–selenolate core and $[PPN]_2[Cl_2Pd(\mu\text{-SePh})_2PdCl_2]$, **286**, 155

chemistry of functionalized thiolates: syntheses and structures of neutral, monomeric $[Zn(SC_6H_4X\text{-}4)_2(N\text{–}N)]$, where X = H, NH_2 and OH, and (N–N) = nitrogen bases 2,2′-bpy, phen and tmen, **295**, 9

chiral phosphine ligands derived from sugars. 15. Synthesis, structure and spectroscopic studies of platinum(II) complexes with chiral phosphine substituted carbohydrate ligands, **290**, 121

comparing structures and reactivity in analogous Fe and Ru complexes. (^{i}Pr-DAB)Fe(CO)$_2$I$_2$ and (^{i}Pr-DAB)FeI$_2$: a perfectly reversible CO-carrier system. (R-DAB = *N*,*N*′-R_2-1,4-diaza-1,3-butadiene), **291**, 438

complexes of copper(II) dihalide with 2,2′-dipyridylamine. X-ray diffraction structures of the [dibromo-bis(dipyam) copper(II)]–water (1:2) and di[chloro-bis(dipyam) copper(II)]diiodide–acetonitrile (1:2) complexes, **290**, 180

complexes of lanthanide dithionates 2. Electrospray mass spectrometry of $Ln_2(S_2O_6)_3$ and their Ph_3PO complexes; the X-ray crystal structure of $Pr_2(S_2O_6)_3(Ph_3PO)_6(H_2O)_6$, **295**, 146

conformational diversity in (octaethylporphinato)(trichloroacetato)iron(III) derivatives, **291**, 49

control of the coordination mode of NO_2^- by the nature of the amine ligand: the importance of H-bond. Crystal structures of the [*mer*-Ni(dpt)$(NO_2)_2$] (**I**) and [*trans*-Ni(333-tet)$(NO_2)_2$] (**II**), **292**, 149

coordination chemistry and X-ray crystal structure of (ferrocenylmethyl)diphenylphosphine, $FcCH_2PPh_2$ [Fc = (η^5-C_5H_5)Fe(η^5-C_5H_4)], **295**, 18

coordination compounds with 4-aminopyridine. An unusual in situ formation of a bridging formato compound. The structure of *catena*-poly[bis(4-amino-pyridine)(formato-*O*)-copper(II)-μ-formato(*O*,*O*′)], **287**, 226

coordination geometries for palladium and platinum: theoretical studies and the synthesis and structure of tris(indazol-1-yl)borate complexes $MMe_3\{(ind)_3BH\}$, **287**, 27

coordination mode of dihydroxybenzoic acids in oxorhenium(V) complexes, **294**, 114

copper(I) and copper(II) complexes of the bidentate imidazole/thioether ligand 1-methyl-2-(methylthiomethyl)-1*H*-benzimidazole, **287**, 204

copper(II) complexes with (2-hydroxybenzyl-2-pyridylmethyl)amine–Hbpa: syntheses, characterization and crystal structures of the ligand and $[Cu(II)(Hbpa)_2](ClO_4)_2\cdot 2H_2O$, **290**, 207

crystallographic and ESR studies of π–π stacked binuclear copper(II) complexes of 1,3-bis(quinoline-2-carboxamido) and 1,3-bis(isoquinoline-3-carboxamido)propane, **288**, 90

crystallographic evidence for the coordination performance of $WO_2S_2^{2-}$. Synthesis, characterization and crystal structure of palladium complexes with mixed-ligand thiotungstates and 1,2-dithiolates: $[NBu_4]_2[PdWO_2S_2(C_3S_5)]$ and $[NBu_4]_2[PdWS_4(C_4N_2S_2)]$, **285**, 241

crystal structure and characterization of an unexpected dinuclear Mn(II)Mn(III) azacryptate, **287**, 14

crystal structure and magnetic properties of the single-μ-chloro copper(II) chain $[Cu(bipy)Cl_2]$ (bipy = 2,2′-bipyridine), **284**, 258

crystal structure of (μ-oxo)bis{aquatris(2-pyridylmethyl)amine}-iron(III) perchlorate, **284**, 124

crystal structures and magnetic properties of the squarate-O^1,O^n-bridged dinuclear copper(II) complexes $[Cu_2(phen)_4(C_4O_4)](CF_3SO_3)_2\cdot 3H_2O$ ($n = 2$) and $[Cu_2(bipy)_4(C_4O_4)](CF_3SO_3)_2\cdot 6H_2O$ ($n = 3$), **287**, 173

crystal structures and stereochemical properties of lanthanide(III) complexes with ethylenediamine-*N*,*N*,*N*′,*N*′-tetraacetate, **288**, 7

C–S, C–H, and N–H bond cleavage of heterocycles by a zero-valent iron complex, $Fe(N_2)(depe)_2$ [depe = 1,2-bis(diethylphosphino)ethane], **291**, 341

cyanamidonitrate–copper(II) complexes of imidazole ligands: X-ray crystallography and physical investigation, **287**, 186

deboronation of $HB(3,5\text{-}Me_2C_3HN_2)_3^-$ in a reaction with $[Re(O)Cl_4]^-$, the X-ray crystal structures of $[\{Re(O)Cl(3,5\text{-}Me_2C_3HN_2H)_3\}(\mu\text{-}O)\{Re(O)Cl_3(3,5\text{-}Me_2C_3HN_2H)\}]$ and $[\{Re(O)Cl_2(3,5\text{-}Me_2C_3HN_2H)_2\}(\mu\text{-}O)\{Re(O)Cl_2(3,5\text{-}Me_2C_3HN_2H)_2\}]$, **285**, 55

di- and trinuclear Pd(II) and Pt(II) complexes containing bridging 2,7-disubstituted naphthyridines, **287**, 72

different chlorination modes of oximes: chlorination of salicylaldoxime coordinated to platinum, **285**, 116

dimolybdenum complexes of bis(2-pyridyl)amine (Hdpa) ligand and its anion (dpa^-); transformation of chelating η^2-Hdpa to bridging η^2-dpa^-, **288**, 82

diphenoxo-bridged dinuclear nickel(II) complex of a dinucleating macrocycle with $N(amine)_6O_2$ metal-binding sites: synthesis, X-ray crystal structure and electrochemistry, **287**, 212

diphosphinito-bridged ruthenium(II) and rhodium(III) complexes with stereogenic metal centers, **292**, 84

dipyridylamine-ligated octamolybdate clusters aggregated into extended ribbons and layers via hydrogen bonding, **292**, 131

displacement of triphenylphosphine from $Cu(PPh_3)_2NO_3$ and $Co(PPh_3)_2Cl_2$ by a diselenoimidodiphosphinato ligand. X-ray crystal structure of $(PPh_3)Cu[Ph_2P(Se)NP(Se)Ph_2]$ and $Co[Ph_2P(Se)NP(Se)Ph_2]_2$ containing the novel $CuSe_2P_2N$ and $CoSe_2P_2N$ inorganic metallocycles, **290**, 256

effect of N–H···S amide–thiolate hydrogen bonding on bond lengths in rubredoxin models $[(CH_3)_3NCH_2CONH_2]_2[M(S_2\text{-}o\text{-}xyl)_2]$, **291**, 388

effects of β-cyclodextrin inclusion on the kinetics of the oxidation of bisferrocenyl cations by bis(pyridine-2,6-dicarboxylato)cobaltate(III) in aqueous solution: crystal structure of dimethyldi-(1-methylferrocene)ammonium bromide, **291**, 365

effects of electronic structure on chiral induction in outer-sphere electron transfer reactions of complexes with the C_3 symmetric ligand, 1,4,7-triazacyclononane-1,4,7-tris[2′(*R*)-2′-propionate](-3), **290**, 197

electrochemical and ^{95}Mo NMR studies of triply-bridged dinuclear oxomolybdenum(V) complexes with various dithiocarbamates, **292**, 7

electrochemical synthesis of cobalt, nickel, copper, zinc and cadmium complexes with *N*[(2-hydroxyphenyl)methylidine]-*N*′-tosylbenzene-1,2-diamine. The crystal structures of {(1,10 - phenanthroline)[*N* - (2 - oxophenyl)methylidine]-*N*-tosylbenzene-1,2-diaminato}nickel(II) and {(1,10-phenanthroline)[*N*-(2-oxophenyl)methylidine]-*N*′-tosylbenzene-1,2-diaminato}copper(II), **295**, 39

electrochemical synthesis of copper and silver pyridine-2-selenolato complexes: crystal structure of (6-trimethylsilylpyridine-2-selenolato)silver(I), $[Ag_6(6\text{-}Me_3SipySe)_6]$, **284**, 133

evidence for a new nitrosyl-to-nitro reaction and an acetamide formation in the $\{RuNO\}^6$-type complex, *cis*-$[Ru(NO)(CH_3CN)(bpy)_2]^{3+}$, **290**, 113

facile and stereoselective condensation of acetone with ammonia ligands on cobalt(III): structure of a N-bonded cyanate complex containing the 2-methyl-2-amino-4-imino-pentane ligand, **290**, 133

facile synthesis and structural properties of (diamine)-tetracarboxylatoplatinum(IV) complexes, **292**, 52

five-coordinate rhenium(III)-thiolato complexes: the structure of $[Re(SCH_2PhOCH_3)_3(PPh_3)_2]$, **290**, 247

fluctuations between square-planar and tetrahedral coordination geometry with bis(2-benzimidazolyl)alkane ligands. Synthesis, spectroscopic properties and X-ray crystal structure of four representative examples, **290**, 105

formation of donor–acceptor Fe(0)–Hg(II) bond in separation and stabilization of optically active iron(0) phosphine complexes. Absolute configuration of (+)-(*R*)-$(CO)_4Fe(\mu\text{-}EtPhPpy)HgCl_2$, **293**, 106

formation of two stable seven-membered chelate rings bound to one metal center due to addition of bifunctional amine to coordinated nitriles, **285**, 203

germanium-containing ferrocenes and ferrocenophanes. Potential precursors for ring-opening polymerizations and 'germaferrocenes', **291**, 355

halfsandwich complexes of ruthenium(II), rhodium(III) and iridium(III) with *N*-substituted 3-hydroxy-2-methyl-4-pyridone ligands, **294**, 62

heterobimetallic nitrido-bridged Ru(II)NOs(VIII) and Ru(II)-NOs(VI) complexes containing ruthenium porphyrins, **290**, 28

heteroleptic poly(pyrazol-1-yl)borate derivatives of the lanthanides. The syntheses of picolinate-*N*-oxide complexes and the X-ray crystal structure of $[Tb\{HB(C_3H_3N_2)_3\}_2(ONC_5H_4CO_2\text{-}2)]$, **285**, 283

hexadentate chromium(III) complexes containing unsymmetrical edta-type ligands: Part IV. Crystal structures of $(-)_{589}$-*trans*(O_5)-Na[Cr(ed3ap)]·$3H_2O$ and Na[Cr(*u*-eddadp)]·$3H_2O$ and CD spectra correlation: structural parameters of $[Cr(edta\text{-}type)]^-$ complexes and their octahedral distortion in relation to the structure of the ligand, **292**, 16

hydrothermal synthesis and characterization of a new vanadyl sulfate compound: crystal structure of $[V(IV)OSO_4(H_2O)_4]\cdot SO_4\cdot[H_2N(C_2H_4)_2NH_2]$, **292**, 112

hydrothermal synthesis and crystal structure of a new molybdenum(VI) arsenate(III), $Co^{III}(en)_3H_3O[(Co^{II}O_6)Mo^{VI}{}_6O_{18}(As^{III}{}_3O_3)_2]\cdot 2H_2O$, **295**, 244

iminoacylation: 2. Addition of alkylated hydroxylamines via oxygen to platinum(IV)-bound nitriles, **292**, 272

indate(III) phthalocyanines: synthesis, spectroscopy and crystal structure of iodophthalocyaninato(2-) indate(III), **288**, 174

infinite two-dimensional sheet-like array of a 2,2′-bipyrimidine-bis(oxalato)chromate(III) complex: synthesis, crystal structure and magnetic properties of $[K(H_2O)Cr(C_2O_4)_2(bpym)]$, **284**, 8

interaction between heterobinuclear molecules and nature of the ground spin state in oximato-bridged $[Cu^{II}M^{II}]_2$ bis-binuclear complexes (M = Cu, Ni, Mn): crystal structure of $[Cu(pdmg)Ni(Me_3[12]N_3)(EtOH)](ClO_4)_2$, **288**, 57

interactions between thiamine and large anions. Crystal structures of (H-thiamine)$[Pt^{II}(SCN)_4]\cdot 3H_2O$ and (H-thiamine)$[Pt^{IV}(SCN)_6]\cdot H_2O$, **290**, 145

iron carbonyls with bulky thiolate ligands: crystal structures of $[Fe_2(CO)_6(\mu\text{-}SC_6H_2{}^tPr_3\text{-}2,4,6)_2]$ and $(C_6H_2{}^tPr_3\text{-}2,4,6)_2S_2$, **284**, 14

iron(II/III) complexes of non-porphyrin macrocyclic $[N_4]^{2-}$ ligands: structure and axial reactivity, **291**, 190

isocyanide derivatives of manganese cobalt nonacarbonyl: synthesis and characterization, **288**, 159

lattice effects in the Mössbauer spectra of salts of $[Fe_4S_4\{S(CH_2)_nOH\}_4]^{2-}$. Crystal structures of $[PPh_4]_2[Fe_4S_4\{S(CH_2)_nOH\}_4]$ (n = 2, 3 and 4), **291**, 101

low-spin iron(III) complexes with N,S coordination: syntheses, structures, and properties of bis(*N*-2-mercaptophenyl-2′-pyridylmethyleniminato)iron(III) tetraphenylborate and bis(*N* - 2 - mercapto - 2 - methylpropyl - 2′ - pyridylmethyleni - minato)iron(III) tetraphenylborate, **285**, 269

macrocyclic complexes derived from the cyclocondensation of sodium 2,6-diformyl-4-substituted phenolates with *N*,*N*-bis(2-aminoethyl)-2-hydroxybenzylamine and in situ transmetallation. An anticipated dinuclear Cd(II) pendant-arm macrocyclic complex and an unexpected mononuclear Fe(III) macrocyclic complex, **294**, 1

magneto–structural correlation of dimeric copper(II) carboxylates with pyridyl-substituted nitronyl nitroxides, **293**, 30

main group metal halide complexes with sterically hindered thioureas XVI: The synthesis, characterization, and crystal structures of two new complexes of 1,3-dimethyl-2(3*H*)-imidazolethione with indium trihalides, **285**, 217

18-membered tetraimino macrocyclic complexes containing furan groups: stacking of two macrocycles and characterization of organic derivatives, **284**, 149

mercury(II) compounds with 1,3-imidazole-2-thione and its 1-methyl analogue. Preparative and NMR spectroscopic studies. The crystal structures of di-μ-iodo-bis[iodo(1,3-imidazolium-2-thiolato-*S*)mercury(II)], bis[bromo(1,3-imidazolium-2-thiolato-*S*)]mercury(II) and bis[μ-(1-*N*-methyl-1,3-imidazole-2-thiolato-*S*)]mercury(II), **294**, 35

metal(II) complexes of the carbonylnitronate ligands $O_2NCHC(O)R^-$ (R = Ph, OMe or OEt) and X-ray structure of (DMANH)$[Ni(O_2NCHC(O)Ph)_3]$ (DMAN = 1,8-bis(dimethylamino)naphthalene), **285**, 18

metal(II) complexes with monodentate S and N ligands as structural models for zinc–sulfur DNA-binding proteins, **285**, 262

metal–metal multiply-bonded complexes of technetium. Part 8. Synthesis and characterization of the α- and β-isomers of $Tc_2Cl_4(dppe)_2$, **288**, 69

molecular structure of $[PPh_4]_2[Co(C_3S_5)_2]$ [$C_3S_5^{2-}$ = 4,5-disulfanyl-1,3-dithiole-2-thionate(2-)] and spectroscopic and electrical properties of its oxidized species, **295**, 239

monoindenyl titanium alkyl halides. The synthesis and molecular structures of $(\eta^5\text{-}C_9H_7)TiBr_3$, $(\eta^5\text{-}C_9H_7)Ti(CH_3)Br_2$, and $(\eta^5\text{-}C_9H_7)Ti(CH_3)Cl_2$, **292**, 220

multinuclear NMR studies of diphosphine, diphosphine–dioxide and diarsine complexes of tin(IV) halides. Structures of $[SnI_4\{o\text{-}C_6H_4(AsMe_2)_2\}]$ and $[SnI_4\{o\text{-}C_6H_4(P(O)Ph_2)_2\}]$, **288**, 142

multinuclear NMR study and crystal structures of complexes of the types *cis*- and *trans*-$Pt(Ypy)_2X_2$, where Ypy = pyridine derivative and X = Cl and I, **295**, 25

neodymium and europium picrate complexes with 3-picoline-*N*-oxide (3-picNO), **292**, 249

new capping agents for oximehydrazonate clathrochelates: sterically controlled synthesis, structural characterization and intramolecular reactions, **284**, 180

new ferracyclic π-allyl and η^3-pentadienyl complexes by carbonylative lactonization of unsaturated diols, **291**, 91

new palladium(II) complexes with bis(pyrazol-1-yl)alkanes, **295**, 136

new phosphino silver(I) derivatives of hydrotris(3-methyl-2-thioxo-1-imidazolyl)borate. X-ray crystal structure of tricyclohexylphosphinesilver(I)-hydrotris(3-methyl-2-thioxo-1-imidazolyl)borate, **285**, 81

new synthesis of $[RuH(CO)(PPh_3)_2L_2][BF_4]$ [L_2 = 2py, 2(4-Mepy) or bpy] and X-ray crystal structure of $[RuH(CO)(PPh_3)_2(4\text{-}Mepy)_2][BF_4]$, **285**, 318

new versatile phosphorus-containing ligands—asymmetric $(XPR_2)(YPR'_2)NH$ (X, Y = O, S; R, R′ = Ph, OEt) and their potassium salts: the crystal and molecular structures of $(SPPh_2)[OP(OEt)_2]NH$, $K[(SPPh_2)\{OP(OEt)_2\}N]\cdot H_2O$ and $(OPPh_2)[OP(OEt)_2]NH\cdot 1/2HCl\cdot 1/4H_2O$, **287**, 61

$[(NH_3)_5CoOP(H)(O)OCo(NH_3)_5]^{4+}$ and its conjugate acid; evidence for the $P^+\text{–}O^-$ description of a terminal P–O bond, **294**, 14

$NH_4TlI_4\cdot 2(18c6)\cdot NH_3$, a new type of a potential double-shell lattice clathrate, **293**, 234

nickel complexes of tetradentate asymmetric ligands, **290**, 14

non-linear optical absorption of $[\{Ni(mnt)_2\}_2]^{2-}$ and its unique counter ion, a disodium triple-decker dication $[Na_2(benzo\text{-}15\text{-}crown\text{-}5)_3(H_2O)_2]^{2+}$, **293**, 89

novel iron(III) complexes with imidazole containing tripodal ligands as model systems for catechol dioxygenases, **291**, 289

novel structural aspects of oxythiamine, an antagonist of thiamine. Crystal structures of three salts, (oxythiamineH)-(picrolonate)$_2\cdot 2H_2O$, (oxythiamineH)$(PtCl_6)$, (oxythiamineH)$_2(V_{10}O_{28}H_2)\cdot 8H_2O$, and a metal complex Pt(oxythiamine)$Cl_3\cdot H_2O$, **295**, 71

on the contrasting reactivity of the phosphaalkene $[(\eta^5\text{-}C_5Me_5)(CO)_2FeP{=}C(NMe_2)_2]$ and the arsaalkene $[(\eta^5\text{-}C_5Me_5)(CO)_2FeAs{=}C(NMe_2)_2]$ towards $[(\eta^5\text{-}C_5H_5)Rh(CO)_2]$, **291**, 311

optically active β-ketoiminato cobalt(II) complexes as highly efficient catalysts in enantioselective borohydride reduction of ketones, imines, and α,β-unsaturated carboxylates, **296**, 86

oxidation state dependent *cis*/*trans* isomerization in ML_2 complexes of the heteroscorpionate ligand, (2-hydroxyphenyl)-bis(pyrazolyl)methane and its derivatives with iron and cobalt, **286**, 80

platina- and palladalactam complexes derived from 2-benzoylacetanilide; syntheses and X-ray structure of $[Pd\{NPhC(O)CHC(O)Ph\}(bipy)]\cdot CH_2Cl_2$, **292**, 260

platinum(II) and palladium(II) complexes with 3,3,8,8-tetramethyl-1,6-dioxa-4,9-diaza-5λ^5-phosphaspiro[4,4]nonane, **295**, 164

platinum(II) and palladium(II) saccharinate complexes, **285**, 145

platinum(II) complexes with the tridentate phosphino–thioether ligand bis(2-diphenylphosphinoethyl)sulfide, $(PPh_2CH_2CH_2)_2S$. X-ray crystal structures of $[Pt\{(PPh_2CH_2CH_2)_2S\}Cl]ClO_4$ and $[Pt\{(PPh_2CH_2CH_2)_2S\}I]I$, **295**, 153

polyiodides of transition metal trischelate cations: syntheses, structures, and spectral and electrical conductivity studies of $[Mn(phen)_3](I_3)_2$ and $[Mn(bpy)_3](I_3)_{1.5}(I_8)_{0.25}$, **284**, 158

potentiometric and spectroscopic studies of copper(II) complexes of bis(1,4,7-triazacyclononane) ligands containing polymethylene and xylyl linker groups, **292**, 172

preliminary studies of $Ni(dmen)_2(CF_3CO_2)_2$, $Ni(deen)_2(CF_3CO_2)_2$ and $Ni(en)(CF_3C(O)CHC(O)CF_3)_2$ as precursors for metalorganic chemical vapour deposition: structures, thermal behaviours and vapour pressures, **288**, 233

preparation and characterisation of new oxovanadium(IV) Schiff base complexes derived from amino acids and aromatic *o*-hydroxyaldehydes, **293**, 1

preparation and spectroscopic properties of the complexes of mercuric thiocyanate with pyridine-2-thione and pyridine-2-carboxylic acid. Crystal and molecular structure of two polymorphs of $Hg(SCN)_2(C_5H_5NS)_2$, **285**, 208

preparation of cyclopentadienyl molybdenum clusters linked by a cyclotetradeca-1,8-diyne ligand, **294**, 232

preparation, structure and electrochemical behavior of dinuclear cyclooctadiene-chelated Ir(I) complexes with 2-aminopyridinato bridges, **292**, 244

reaction of formaldehyde with cyano(ethylenediamine)-(glycylglycinato)cobalt(III): crystal structures of cyano-(ethylenediamine) [*N*-(4-carbonyl-4-hydroxymethyl-oxazolidine)glycinato]cobalt(III)·3.5H_2O and dicyanobis(ethylenediamine) {μ - *N* - [*N* - (((methylimino)methyl)carbonyl)-α-methylserinato)glycylglycinato}dicobalt(III) tetrahydrate, **292**, 96

reaction of hydrous nitrosylruthenium trichloride with 2-pyridinemethanol, **290**, 36

reaction patterns of 'tetraferrio-azaallenium', $[(OC)_3Cp_2Fe_2(\mu_4\text{-}C{=}N{=}C)Fe_2Cp_2(CO)_3]^+$, and its 'activated cyanide'-precursor (μ-phthalimidocarbyne)tricarbonyldicyclopentadienyl-diiron(1 +), **291**, 266

reactions of σ-π alkenyl complexes $[Fe_2(CO)_6(\mu\text{-alkenyl})(\mu\text{-}PPh_2)]$ with bis(diphenylphosphino)methane (dppm); carbonyl substitution, migratory carbonyl insertion, phosphorus–carbon bond formation and cleavage, proton shift and alkenyl isomerisation reactions, **291**, 178

reactions of bis(silyl)palladium(II) complexes with allyl halides. Synthesis of mono(silyl)palladium(II) halides and X-ray structure of *trans*-$PdCl(SiF_2Ph)(PMe_2Ph)_2$, **296**, 19

reactions of H_2, silanes, and olefins with superelectrophilic cationic rhenium complexes: heterolytic cleavage of H_2 and relation to the structure and function of hydrogenases, **294**, 240

reactivity towards some weak protic electrophiles of the di-μ-hydroxo complex $[\{Pd(\mu\text{-}OH)(tmeda)\}_2][BPh_4]_2$: crystal structure of $[\{Pd(tmeda)\}_2(\mu\text{-}OH)(\mu\text{-}HNC_6H_4Cl\text{-}p)][BPh_4]_2$, **287**, 37

receptors based on 2,2′:6′,2″-terpyridine fragments containing peripheral amino groups, **292**, 28

silver(I) tetraiodoethylene complexes with twisted olefin moiety, **290**, 251

solid and solution studies of ternary copper(II) complexes formed by heteroaromatic N bases and the anion of *N*,*N*-bis(2-hydroxyethyl)glycine (bicine), **285**, 326

some complexes derived from zinc salicylate or 3,5-di-tert-butylsalicylate. The crystal structure of (2,2′-bipyridyl)-(methanol)(*O*-salicylato)(*O*,*O*′-salicylato)zinc, **287**, 89

spin-crossover in the $[Fe(abpt)_2(NCX)_2]$ (X = S, Se) system: structural, magnetic, calorimetric and photomagnetic studies, **291**, 279

spontaneous formation of trinickel clusters containing μ^3-imido bridges from 2-nitrophenols and $Ni(PMe_3)_4$, **287**, 199

structural and spectroscopic characterisation of the mixed valence platinum complex $PtdabI_3$ (dab = *o*-diaminobenzene), **287**, 218
structural aspects of cesium ion selectivity by tetrabenzo-24-crown-8, **290**, 86
structural characterization of a new ligand mode of 2,6-diacetylpyridine bis(semicarbazone), H_2daps, **292**, 121
structural diversity in thallium chemistry: III. The first structurally characterized examples of the pentabromothallate(III) anion in the 1,1,4,4-tetramethylpiperazinium and *N*,*N*′-diethyltriethylenediammonium salts contrasted with the mixed anions in the 1,1,3,3-tetramethylimidazolidinium salt, **285**, 122
structural diversity in thallium chemistry: II. Potential pentabromothallate(III) derivatives: examples of incipient bromide ion coordination?, **284**, 215
structural, spectroscopic and catalytic properties of water-soluble hydride rhodium complexes $[RhH(Rtpa^+I^-)_4]H_2O$ (R = Me, Et), **293**, 110
structural studies of ReO(V) mixed ligand [SNS][Cl] and [SNS][S] complexes, **287**, 142
structural study of the electrochemically synthesized binuclear complex bis{1-phenylglyoxal bis(3-piperidylthiosemicarbazone)zinc(II)}, **294**, 79
structural trends in a series of divalent transition metal triazole complexes, **293**, 61
structure and characterization of a mononuclear Fe^{II} dialkyldithiocarbamate complex $Fe(OC_4H_8dtc)_2(DMF)$, **293**, 95
structure and DNA binding properties of the gold(III) complex $[AuCl_2(esal)]$, **285**, 309
structure and magnetic properties of *catena*-(μ-CrO_4-*O*,*O*′)$[M^{II}(L)]\cdot 5H_2O$ ($M = Ni^{II}$, Cu^{II}; L = 3,14-dimethyl-2,6,13,17-tetraazatricyclo[14,4,$0^{1,18}$,$0^{7,12}$]docosane), **293**, 100
structure and magnetic properties of dinuclear [Mn(III)(salen)$(H_2O)]_2(ClO_4)_2$ and polynuclear [Mn(III)(salen)$(NO_3)]_n$, **290**, 8
structure and magnetic properties of (μ-oxo)bis[*meso*-tetrakis(*p*-tolyl-porphyrinato)iron(III)] and (μ-oxo)bis[*N*,*N*′-ethylenebis(2-acetylphenoliminato)iron(III)], **290**, 51
structure and physical properties of hexacyanomanganate(III), $[Mn^{III}(CN)_6]^{3-}$, **284**, 175
structure and properties of a novel OH^- bridged dimetal helical complex with 6,6‴-dimethyl-2,2′:6′,2″:6″,2‴:6‴,2⁗-quinquepyridine ligand, **290**, 127
structure and stability of aminoiminomethanesulfonic acid, **286**, 149
structures of 1,3-bis(diphenylphosphino)propane platinum(II) alditolate complexes, **285**, 10
studies in aryltin chemistry: Part XIV. Crystal and molecular structures of tris(*o*-methoxyphenyl)tin(IV) chloride, fluoride and oxide, **294**, 224
studies of the binding modes of carboxylate donors with manganese(III) complexes containing tetradentate Schiff base ligands. Crystal structures of the complexes $[Mn(3\text{-}CH_3Osalpn)(HO_2CC_6H_4CO_2)]_n$ and $[\{Mn(5\text{-}NO_2salpn)(OH)(H_2O)\}\{Mn(5\text{-}NO_2salpn)(HO_2CC_6H_4CO_2)(H_2O)\}]\cdot C_2H_5OH$, **293**, 210
studies of the construction of coordination polymers using linear pyridyl-donor ligands, **292**, 231
study on the formation of mixed ligand oxorhenium and oxotechnetium complexes (SNS/S combination), **295**, 1
syntheses and anti-*Candida* activity of cobalt(II) complexes of octanedioic acid ($odaH_2$) and nonanedioic acid ($ndaH_2$); X-ray crystal structures of $[Co(phen)_3]oda\cdot 14H_2O$ and $[Co(phen)_3]nda\cdot 11.5H_2O$ (phen = 1,10-phenanthroline), **293**, 160
syntheses and structures of Group 1 mellitate compounds, $K_6[C_6(COO)_6]\cdot 8H_2O$ and $Cs_5[C_6(COO)_6H]\cdot 7H_2O$, **286**, 207
syntheses and structures of nickel(II) and copper(II) complexes with a 14-membered tetraaza macrocyclic ligand, **286**, 67
syntheses and X-ray structures of mixed-metal gold phosphine clusters including an example having a highly asymmetric Re_2Au_2 skeleton, **294**, 255
syntheses, crystal structures and autoreduction behavior of antiferromagnetically coupled dicopper(II) oximato complexes, **293**, 20
syntheses, crystal structures and coordination compounds of some 2-hetarylcyanoximes, **284**, 85
syntheses, crystal structures and magnetic properties of heterometallic copper–lanthanide clusters $[Cu_{12}Ln_6(\mu_3\text{-}OH)_{24}(\mu\text{-}O_2CR)_{12}(H_2O)_{18}(\mu_{12}\text{-}ClO_4)]^{5+}$ (Ln = La, Nd; R = CH_2Cl, CCl_3), **293**, 129
syntheses, crystal structures and nonlinear optical properties of heteronuclear clusters $[MS_4Cu_4(SCN)_2(NC_5H_5)_6]$ (M = W, Mo), **294**, 73
syntheses, properties and Mössbauer studies of cyanamide and cyanoguanidine complexes of iron(II). Crystal structures of *trans*-$[FeH(NCNH_2)(Ph_2PCH_2CH_2PPh_2)_2][BF_4]$ and *trans*-$[Fe(NCNEt_2)_2(Et_2PCH_2CH_2PEt_2)_2][BF_4]_2$, **291**, 39
synthesis and characterisation of mono- and binuclear palladium and platinum complexes with organochalcogenides, **285**, 190
synthesis and characterization of a series of new thiocarboxylate complexes of molybdenum(V), **284**, 223
synthesis and characterization of five-coordinate rhenium(V) and technetium(V) mixed ligand bifunctional complexes carrying the SNS/S or the SNN/S donor atom set. Crystal structure of $ReO\{[(C_2H_5)_2NCH_2CH_2N(CH_2CH_2S)_2](p\text{-}H_2N\text{–}PhS)\}$ and $ReO\{[(CH_2)_4NCH_2CH_2NCH_2CH_2S](p\text{-}H_2N\text{–}PhS)\}$, **285**, 97
synthesis and characterization of oxorhenium-'3 + 1' mixed-thiolate complexes. Crystal and molecular structures of $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}(C_6H_4X\text{-}4\text{-}CH_2S)]$ (X = F, Cl, Br, OMe) and of the pendant thiolate compounds $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}(\eta^1\text{-}SCH_2CH_2SCH_2CH_2SH)]$ and $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}\{\eta^1\text{-}SCH_2CH(OH)CH(OH)CH_2SH\}]$, **284**, 252
synthesis and characterization of ruthenium 'pincer' complexes containing sulfur atom donor ligands. The X-ray crystal structure of $[RuCl_2(\eta^3\text{-}NN'N)(SMe_2)]$ (NN′N = {2,6-$(Me_2NCH_2)_2C_5H_3N$}), **287**, 113
synthesis and characterization of siloxy-substituted diene and dienyl complexes of the $Ru(C_5Me_5)$ fragment, **296**, 170
synthesis and characterization of the triethylphosphine adduct of bis(2,4-dimethylpentadienyl)tungsten, $[W(2,4\text{-}C_7H_{11})_2(PEt_3)]$, a complex with one η^5-**U** and one η^5-**S** pentadienyl ligand, **288**, 154
synthesis and crystal structure of $Na[Pd(Hpba)]\cdot 2H_2O$ [H_4pba = 1,3-propylenebis(oxamic acid)]. The first bis(oxamato)-palladium(II) complex and an unexpected trideprotonated form of the ligand, **284**, 292
synthesis and crystal structure of $[Re(CO)_3(\mu\text{-}CH_3CO_2)(THF)]_4$, **284**, 304
synthesis and crystal structure of tetranuclear copper(I) and silver(I) complexes bridged by 2-amino-1,3,4-thiadiazole (atdz): $[Cu_4(atdz)_6](ClO_4)_4\cdot 2CH_3OH$ and $[Ag_4(atdz)_6](ClO_4)_4$, **290**, 153
synthesis and crystal structure of two new rare earth trimesate complexes: $ErTMA(H_2O)_5\cdot 3.5H_2O$ and $YTMA(H_2O)_6$, **284**, 139
synthesis and formation of metal complexes of 4-alkynyl and 4-cyano-2,6-diisopropylphenylisocyanides, **284**, 205
synthesis and properties of (diamine)platinum(II) complexes of pyridine carboxylate isomers and their antitumor activity, **292**, 238

synthesis and structural characterisation of rhenium(V) oxo-thioether thiolate complexes with pendant ester or carboxylate groups, **285**, 76

synthesis and structural characterization of binuclear silver(I) complexes bridged by three polydentate phosphine ligands, **284**, 278

synthesis and structural characterization of face-sharing bioctahedral complexes containing poly(pyrazolyl)borate ligands: $[HB(Me_2Pz)_3BH][X_3Mo(\mu\text{-}X)_2(\mu\text{-}H)MoTp^*]$ (X = Cl or Br; $Tp^* = HB(Me_2Pz)_3$; Pz = pyrazolyl), **292**, 182

synthesis and structural characterization of $[NEt_4][Fe_3(\mu_3\text{-}O)(\mu_3\text{-}AuPPh_3)(\mu\text{-}CO)_3(CO)_6]$, the new $[Au_6(\mu_3\text{-}S)_2(PPh_3)_6][Fe_3(\mu_3\text{-}S)(\mu\text{-}AuPPh_3)(CO)_9]_2$ and $[Au_6(\mu_3\text{-}S)_2(PPh_3)_6][Fe_5(\mu_3\text{-}S)_2(CO)_{14}]$ ionic solids containing assemblages of cluster-cations and cluster-anions, **291**, 372

synthesis and structural characterization of two new Zintl $[M_7As_3Q_{13}]^{4-}$ anions (M = Cu, Ag; Q = Se, Te) containing distorted cubane $[M_7Q]$ clusters, **288**, 1

synthesis and structure of a heptacoordinated cadmium(II) complex, **288**, 118

synthesis and structure of a novel terminal imido complex of titanium, **292**, 105

synthesis and structure of niobium and tantalum derivatives of bis(dicyclohexylphosphino)methane (dcpm), **288**, 35

synthesis and structures of 2-$[Me_2NCH_2CH_2N(Me)CH_2]C_6H_4$ substituted tin compounds, **290**, 100

synthesis and structures of (*SP*-4-2)-diiodobis(2-hydroxyethylamine)platinum(II), (*SP*-4-2)-dichlorobis(2-hydroxyethylamine)platinum(II) and (*OC* - 6 - 22) - bis(2 - hydroxyethylamine)tetrachloroplatinum(IV) in the crystal, **292**, 127

synthesis and structures of $TiCl_2(2\text{-}OC_6H_4OCH_3)_2$ and $[TiCl_2(2\text{-}OC_6H_4OCH_3)(\mu\text{-}Cl)]_2$ and stepwise reactions of $[TiCl_2(2\text{-}OC_6H_4OCH_3)(\mu\text{-}Cl)]_2$ with 2-propanol, **293**, 12

synthesis and structures of tungsten ferrocenyldiazoalkane complexes derived from a dinitrogen complex. Part 58. Preparation and properties of molybdenum and tungsten dinitrogen complexes, **285**, 336

synthesis and X-ray characterization of phthalocyaninato(2)-antimony(III) triiodide, **293**, 155

synthesis and X-ray crystallographic structures of dichloronickel(II) and -palladium(II) complexes containing the biphenyl diamine ligand 6-(2-dimethylaminoethyl)-6,7-dihydro-5-H-dibenz[c,e]azepin, **294**, 28

synthesis, characterization and crystal structures of rhenium(V) complexes with diphosphines, **294**, 47

synthesis, characterization, and structural analyses of three high-coordination tetra-n-butylammonium lanthanide(III) complexes, **292**, 137

synthesis, characterization, spectroscopy, magnetism and X-ray structures of red and green Cu(II) chloride adducts with bis(2-benzimidazolyl)propane, **288**, 220

synthesis, crystal structure and aquation kinetics of cobalt(III) complex of N-substituted tetra-aza macrocycle: C-*rac*-1,5,8,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane, **285**, 107

synthesis, crystal structure and magnetic behaviour of the new linear copper(II) tetranuclear compound $[Cu_4(\mu_2\text{-}AcO)_6(AcO)_2(DPQ)_2]$ (DPQ = 2,3-bis(2-pyridyl)quinoxaline), **286**, 62

synthesis, crystal structure and magnetic properties of a copper(II)–nitroxide linear chain complex, **287**, 181

synthesis, crystal structure and magnetic properties of *catena*-poly[(2,2′-bipyridylcopper)-di-μ-thiocyanato] and bis(2,2′-bipyridyl)thiocyanatocopper(II) perchlorate, **286**, 169

synthesis, crystal structures and magnetic properties of trinuclear oxo-centered manganese complexes of the general formula $[Mn_3O(O_2CR)_6L_3]^{z+}$ (R = Me, Et, Ph, CH_2Cl; L = 3-methylpyridine or water $z = 1, 0$), **287**, 82

synthesis, molecular and crystal structure of the zwitterionic Re(V)–oxo complex $ReOCl_3(L)(PPh_3)$ with L = 2-(diethylaminomethyl)-4-methylphenol, **295**, 209

synthesis of a dinuclear tungsten complex, $W_2(\mu\text{-}H)(\mu\text{-}OC_2H_5)(\mu\text{-}O_2CC_6H_5)(\eta^1\text{-}OC_2H_5)_2Cl_2(\eta^2\text{-}P,P'\text{-dppp})$ where dppp is 1,3 bis(diphenylphosphino) propane, **284**, 303

synthesis of 6-amino-6-(4-aminobenzyl)-1,4,8,11-tetra-azacyclotetradecane, a new pentadentate polyamine ligand, and crystal structure of its Co(III) complex, **288**, 40

synthesis of bis-(acyl, carbamoyl or alkoxycarbonyl) iron monomer or dimer complexes by reaction of oxalyl chloride with various tetracarbonyl ferrates: $\{Fe[C(O)R](CO)_4\}^-$, **291**, 166

synthesis of (ferrocenyl-indenyl)cyclopentadienyliron compounds with and without a bridging group via a CpFe transfer reaction, **286**, 215

synthesis of five-coordinate Pt(II) complexes containing organogermanium fragments in axial position. Structure of $[PtCl(GePh_2Cl)(2,9\text{-}Me_2\text{-}1,10\text{-phen})(CH_2{=}CH_2)]$, **285**, 70

synthesis of hexametallic $CH_2\{(\eta^5\text{-}C_5H_4)Fe(CO)_2SnPh_2(\eta^5\text{-}C_5H_5)Fe(CO)_2\}_2$, **291**, 252

synthesis of *N*-allyl-2,3-dihydrobenzoxazol-2-ylidene complexes of chromium(0) and tungsten(0), **288**, 47

synthesis of organo(siloxo)platinum and -palladium complexes and preparation of supported nanoclusters by facile ligand reduction, **294**, 266

synthesis of the ligand *N*,*N*′-bis(2-tert-butylthiobenzenylidene)-diethylenetriamine: its reactivity with nickel(II) salts, **287**, 105

synthesis, platinum(II) complexes and structural aspects of the new tetradentate phosphine *cis*,*trans*,*cis*-1,2,3,4-tetrakis-(diphenylphosphino)cyclobutane, **290**, 167

synthesis, reactivity, and structures of (μ-acetamidato)hexakis-(acetonitrilo)dimolybdenum(II) tris(tetrafluoroborate) and derivatives, **287**, 159

synthesis, spectroscopic, magnetic properties and X-ray crystal structure of di-μ-hydroxo-bis(μ-perchlorato-*O*,*O*′)-bis-[bis(2-amino-4-methylpyrimidine)copper(II)]: bridging perchlorate results in significant deviation from the predicted magnetic exchange, **286**, 24

synthesis, structural characterization and biological activity of helicin thiosemicarbazone monohydrate and a copper(II) complex of salicylaldehyde thiosemicarbazone, **286**, 134

synthesis, structure and EPR spectroscopy of a vanadium(IV)–amide metallacyclic complex, **285**, 134

synthesis, structure and magnetic behavior of a di-μ-chloro-di-μ-imidazolato-tetracopper(II) complex with two dinucleating macrocycles, **287**, 52

synthesis, structure and magnetic properties of a linear trimanganese(II,III,II) complex with hydroxo and 1,2,4-triazole as bridging ligands, **295**, 229

synthesis, structure and magnetic properties of oxo-centered trinuclear manganese complex $[Mn_3O(O_2CC_3H_7)_6(C_5H_5N)_3]\cdot ClO_4$, **294**, 109

synthesis, structure, and redox properties of the sulfido bridged tetranuclear cluster complex $[(Ph_3P)_2N][Fe_3(CO)_9\{\mu_3\text{-}SFeCp(CO)_2\}]$, **286**, 142

synthesis, studies and molecular structure of trinuclear cyanide-bridged copper–iron complexes, **292**, 157

synthesis, vibrational study, crystal structure and density functional calculations of $[Ni(dien)_2]^{2+}$ complexes. Configurational and conformational study of $[M(dien)_2]^{n+}$ complexes, **290**, 57

synthesis, X-ray crystal structure, and reactivity of the monomeric dithio-*o*-carboranyl iridium complex [Ir(η^5-C_5Me_5)(η^2-$S_2C_2B_{10}H_{10}$)], **289**, 141
synthesis, X-ray structural determination and Mössbauer characterization of Schiff bases bearing ferrocene groups, their reduced analogues and related complexes, **287**, 117
synthetic approaches to Fe(II) complexes of the bulky phosphine P^{t}Bu$_2$Me, **291**, 226
synthetic routes for hydrotris(pyrazolyl)borate complexes of rhodium(III), **284**, 191
synthetic, spectroscopic and X-ray crystallographic structural studies on some copper(II) complexes of the 6-methylpyridine-2-carboxaldehyde Schiff base of *S*-methyldithiocarbazate, **287**, 8
template synthesis, crystal structure, and solution behavior of a hexaaza macrocyclic nickel(II) complex containing two *N*-aminoethyl pendant arms, **293**, 140
terminal S-coordinated thiocyanate in a nickel(II) complex. X-ray structures of Ni(poph)(H_2O)(SCN)(NCS) and Ni(poqh)(H_2O)$(NCS)_2$ (poph = 2-pyridinecarboxaldehyde 1-oxide 2′-pyridinylhydrazone, poqh = 2-pyridinecarboxaldehyde 1-oxide 2′-quinolynylhydrazone), **285**, 297
the crystal structure and doublet emission properties of *trans*-[Cr$(NH_3)_2$(cyclam)]$(ClO_4)Cl_2$, **285**, 313
the crystal structures and electronic properties of bis(di-2-pyridylamine)copper(II), bis(tetrafluoroborate) and bis[bisaquabis(di-2-pyridylamine)copper(II)] sulfate heptahydrate, **292**, 57
the effect of replacing carbonyl with acetonitrile on the reactivity of the bimetallic system $[Mo_2Cp_2(\mu\text{-}SMe)_3(L)_2]^+$ (L = CO, CH_3CN). Crystal structure of $[Mo_2Cp_2(\mu\text{-}SMe)_2(CH_3CN)_4](BF_4)_2$, **284**, 107
the first binuclear Mo–S cluster compound containing citrate ligands, $K_5(NH_4)[Mo_2O_2(\mu\text{-}S)_2(C_6H_4O_7)_2]\cdot CH_3OH\cdot 5H_2O$, characterized by X-ray single crystal structure determination, **285**, 152
the first crystal structure of a complex from a lower rim calix(4)arene alicyclic amino derivative: {tetrakis[1-(ethyl)piperidinium]oxyl}*p*-tert-butylcalix(4)arene-bis-[tetrachloroaurate(III)]dichloro dihydrate, **285**, 142
the first X-ray crystal structures of cobalt complexes containing monodentate and bridging ethylenediamine ligands, **288**, 53
the hydrothermal synthesis and structural characterization of a new class of compounds: nickel organoamine-halocadmates, **292**, 41
the kinetics of the sequential hydrolysis reactions of the *xxm*-$[CoCl_2(dien)(py)]^+$ cation and the crystal structure of the $ZnCl_4^{2-}\cdot 2H_2O$ salt, **292**, 198
the methylcobalt(III) complex of a tetrapodal pentadentate amine ligand, 2,6-bis(1′,3′-diamino-2′-methyl-prop-2′-yl)-pyridine, **286**, 98
the molecular and electronic structure of the asymmetric, trinuclear complex $[LFe^{III}(\mu\text{-}O)(\mu\text{-}piv)_2Fe^{III}(\mu\text{-}OH)(\mu\text{-}piv)_2Fe^{III}(piv)_2]\cdot C_6H_5CH_3$ (L = 1,4,7-trimethyl-1,4,7-triazacyclononane, piv = pivalate(1-)), **291**, 258
the nickel(II) complex of a new multidentate ligand containing both imidazole and pyrazole moieties, **288**, 239
the nitrido complex $(NEt_4)[MoCl_3(N)(OH)(MeCN)]$, **295**, 249
the preparation and coordination chemistry of ^{i}Pr$_2$P(E)NHP(E′)^{i}Pr$_2$ (E, E′ = Se; E = Se, E′ = S; E = S, E′ = O; E,E′ = O, **290**, 1
the reaction of triorganophosphorus diiodides, R_3PI_2, with zinc metal powder; dependency of product on R; the X-ray crystal structures of dimeric $\{ZnI_2[P(NMe_2)_3]\}_2$ and monomeric $ZnI_2(PPh_2Me)_2$, **292**, 213
the relationship between surface derivatization and the physical properties of bisarylphosphinidenetriironnonacarbonyl clusters, **294**, 140
the syntheses, spectra and structures of five-coordinate cobalt(II) complexes of pyrazolyl-containing ligands, **284**, 266
the synthesis and structure of some, *rac–trans*-, *meso–trans*- and (*S,S*)-*trans*- Co(III) complexes with Me_6-[14]-diene(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene), **292**, 73
three mononuclear copper complexes of 9-hydroxy-9*H*-fluorene-9-carboxylic acid, **294**, 163
transition metal complexes of 1,3-bis(thiomorpholino)propane: crystal structure and dynamic ^{1}H NMR study, **285**, 31
tricarbonylchromium complexes with methoxy substituted benzocyclobutenone and benzocyclobutenedione ligands, **296**, 150
tridentate N_3 capping coordination behaviour of potentially multi-site coordinating cyclotriphosphazenes: synthesis and spectroscopic studies of 2,2-spiro(1,3-propanediamino)-4,4,6,6-tetrakis(3,5-dimethylpyrazolyl)cyclotriphosphazene, 2,2-diamino-4,4,6,6-tetrakis(3,5-dimethylpyrazolyl)-cyclotriphosphazene and their mononuclear complexes; X-ray structure of ATPCTP·$CoCl_2$, **286**, 127
triethanolamine complexes of copper, **294**, 153
trinuclear vanadium dialkyldithiocarbamate complexes. Synthesis, structure and spectroscopic characterization of $[Et_4N][V_3(\mu_3\text{-}S)(\eta^2\text{-}\mu\text{-}S_2)_3(Et_2dtc)_3]\cdot 2CH_3CN$, **286**, 7
triphenylsilanolato complexes of Group 4 metals: syntheses, X-ray crystal structures and catalytic cyclization reaction of 1,5-hexadiene, **286**, 14
two diphenylformamidinate compounds of the quadruply-bonded Re_2^{6+} core, **285**, 149
unusual coordination mode of a 2-pyridyl ketone oxime ligand in bis(4-butylphenyl 2-pyridyl ketone oximate)palladium(II), **285**, 138
vanadium phenoxide complexes with oxide, nitride or hydrazide co-ligands: preparation and crystal structures of $[V(OC_6H_3Pr^i_2\text{-}2,6)_3NLi(C_4H_8O)_3]$, $[\{VO_2(OC_6H_3Pr^i_2\text{-}2,6)_2\}_2\text{-}\{\mu\text{-}Li(C_4H_8O)_2\}_2]$ and $[V(NNMe_2)(OC_6H_3Pr^i_2\text{-}2,6)_3]$, **285**, 178
variations in capping the $Zr_6O_4(OH)_4$ cluster core: X-ray structure analyses of $[Zr_6(OH)_4O_4(OOC\text{–}CH\text{=}CH_2)_{10}]_2(\mu\text{-}OOC\text{–}CH\text{=}CH_2)_4$ and $Zr_6(OH)_4O_4(OOCR)_{12}(PrOH)$ (R = Ph, CMe = CH_2), **284**, 1
versatility of the cyanato ligand: structure and magnetic properties of four new copper(II)–cyanato compounds derived from tridentate amines, **286**, 189

C_2-Symmetric ligands
stereoselective reduction of ketones using extracoordinate silicon: C_2-symmetric ligands, **296**, 208

CT complexes
Rh-complex promoted synthesis of poly(arylacetylene) with pendant TTF units and formation of CT complexes of the polymer with π-acceptors, **296**, 278

Cyanamide complexes
syntheses, properties and Mössbauer studies of cyanamide and cyanoguanidine complexes of iron(II). Crystal structures of *trans*-$[FeH(NCNH_2)(Ph_2PCH_2CH_2PPh_2)_2][BF_4]$ and *trans*-$[Fe(NCNEt_2)_2(Et_2PCH_2CH_2PEt_2)_2][BF_4]_2$, **291**, 39

Cyanamidonitrate complexes
cyanamidonitrate–copper(II) complexes of imidazole ligands: X-ray crystallography and physical investigation, **287**, 186

Cyanato bridge complexes
versatility of the cyanato ligand: structure and magnetic properties of four new copper(II)–cyanato compounds derived from tridentate amines, **286**, 189

Cyanide bridged complexes
synthesis, studies and molecular structure of trinuclear cyanide-bridged copper–iron complexes, **292**, 157
Cyanide complexes
reduction of the NO^+ ligand in the pentacyanonitrosylosmate(II) ion, **285**, 129
Cyano-bridged complexes
cyano-bridged dinuclear iron complexes in different oxidation states, **287**, 95
Cyano complexes
widely differing stabilities of molecule-bridged cyanodiiron(III, II) species in non-aqueous solvents, **291**, 66
Cyanoguanidine complexes
syntheses, properties and Mössbauer studies of cyanamide and cyanoguanidine complexes of iron(II). Crystal structures of *trans*-$[FeH(NCNH_2)(Ph_2PCH_2CH_2PPh_2)_2][BF_4]$ and *trans*-$[Fe(NCNEt_2)_2(Et_2PCH_2CH_2PEt_2)_2][BF_4]_2$, **291**, 39
Cyanometallate
structure and physical properties of hexacyanomanganate(III), $[Mn^{III}(CN)_6]^{3-}$, **284**, 175
Cyanoxime complexes
syntheses, crystal structures and coordination compounds of some 2-hetarylcyanoximes, **284**, 85
Cyclam complexes
synthesis of 6-amino-6-(4-aminobenzyl)-1,4,8,11-tetra-azacyclotetradecane, a new pentadentate polyamine ligand, and crystal structure of its Co(III) complex, **288**, 40
the crystal structure and doublet emission properties of *trans*-$[Cr(NH_3)_2(cyclam)](ClO_4)Cl_2$, **285**, 313
Cyclic carbonylation
rhodium-catalyzed cyclic carbonylation of 2-phenylethynylbenzamide under water–gas shift reaction conditions, **296**, 195
Cycloaddition
$Ni(COD)_2$: an efficient promoter in the construction of 5-cyclopent-2-enone derivatives from allyl halides, alkynes and carbon monoxide, **296**, 94
Cycloalkadiyne complexes
preparation of cyclopentadienyl molybdenum clusters linked by a cyclotetradeca-1,8-diyne ligand, **294**, 232
Cyclobutadienyl complexes
preparation of cyclopentadienyl molybdenum clusters linked by a cyclotetradeca-1,8-diyne ligand, **294**, 232
Cyclodextrin
effects of β-cyclodextrin inclusion on the kinetics of the oxidation of bisferrocenyl cations by bis(pyridine-2,6-dicarboxylato)cobaltate(III) in aqueous solution: crystal structure of dimethyldi-(1-methylferrocene)ammonium bromide, **291**, 365
Cyclodextrin complexes
thermodynamic and structural study of inclusion complexes between trivalent lanthanide ions and native cyclodextrins, **293**, 53
Cycloheptatrienyl complexes
ring-opening polymerization of norbornene by η-cycloheptatrienyl molybdenum and tungsten complexes, **288**, 226
Cyclohexadienyliron electrophiles
addition of aryl substituents to cyclohexadienyliron electrophiles in the development of routes to C_{12} central building blocks for alkaloid synthesis, **296**, 139
Cyclopalladated complexes
synthesis and characterisation of main-chain oligomeric cyclopalladated azobenzene complexes, **292**, 163
Cyclopentadienyl complexes
novel features associated with the electrochemically driven bis(η^5-pentaphenylcyclopentadienyl)iron(II)–iron(III) redox transformation at an electrode–microcrystal–solvent (electrolyte) interface, **291**, 21
photo-assisted preparation of double isocyanide insertion complexes of η^5-cyclopentadienyliron(II) bearing only isocyanide: the molecular structure of $(\eta^5\text{-}C_5H_5)Fe(XylNC)_2[C(=NXyl)C(=NXyl)CH_3]$ (XylNC = 2,6-$Me_2C_6H_3$-NC), **286**, 233
reactions of $Cp'(CO)_2MnPPh_2H$ with CH_3COCl and $CH_3S(O)_2Cl$ in THF/triethylamine: evidence of the first complex stabilization of a phosphorus homolog of the sulfonamides, **288**, 101
reduction of cationic P-coordinated (diphenylphosphino)alkyne iron complexes with sodium borohydride, **291**, 207
synthesis and characterization of long chain acyl and alkyl complexes of the types $[(\eta^5\text{-}C_5R'_5)Fe(CO)_2\{C(O)R\}]$ (R' = H, CH_3), $[CpW(CO)_3\{C(O)R\}]$ and $[CpW(CO)_3R]$ (R = *n*-alkyl), **286**, 221
synthesis of hexametallic $CH_2\{(\eta^5\text{-}C_5H_4)Fe(CO)_2SnPh_2(\eta^5\text{-}C_5H_5)Fe(CO)_2\}_2$, **291**, 252
Cyclopentenone
$Ni(COD)_2$: an efficient promoter in the construction of 5-cyclopent-2-enone derivatives from allyl halides, alkynes and carbon monoxide, **296**, 94
Cyclophosphazene complexes
iron and ruthenium organometallic derivatives of spiro-substituted cyclotriphosphazenes bearing nitrile spacer: synthesis and electronic structure, **294**, 20
tridentate N_3 capping coordination behaviour of potentially multi-site coordinating cyclotriphosphazenes: synthesis and spectroscopic studies of 2,2-spiro(1,3-propanediamino)-4,4,6,6-tetrakis(3,5-dimethylpyrazolyl)cyclotriphosphazene, 2,2-diamino-4,4,6,6-tetrakis(3,5-dimethylpyrazolyl)-cyclotriphosphazene and their mononuclear complexes: X-ray structure of ATPCTP·$CoCl_2$, **286**, 127
Cyclopropylcarbene complexes
intramolecular coupling of conjugated enynes with cyclopropylcarbene–chromium complexes; a complex reaction pathway, **296**, 273

1D and 2D-NMR
mononuclear lanthanide complexes of tripodal ligands: synthesis and spectroscopic studies, **285**, 49
Deboronation reaction
deboronation of $HB(3,5\text{-}Me_2C_3HN_2)_3^-$ in a reaction with $[Re(O)Cl_4]^-$, the X-ray crystal structures of $[\{Re(O)Cl(3,5\text{-}Me_2C_3HN_2H)_3\}(\mu\text{-}O)\{Re(O)Cl_3(3,5\text{-}Me_2C_3HN_2H)\}]$ and $[\{Re(O)Cl_2(3,5\text{-}Me_2C_3HN_2H)_2\}(\mu\text{-}O)\{Re(O)Cl_2(3,5\text{-}Me_2C_3HN_2H)_2\}]$, **285**, 55
Decavanadate
novel structural aspects of oxythiamine, an antagonist of thiamine. Crystal structures of three salts, (oxythiamineH)(picrolonate)$_2$·$2H_2O$, (oxythiamineH)$(PtCl_6)$, (oxythiamineH)$_2(V_{10}O_{28}H_2)$·$8H_2O$, and a metal complex Pt(oxythiamine)Cl_3·H_2O, **295**, 71
Deltahedral boranes
alkyl derivatives of $[B_6H_6]^{2-}$: NMR and vibrational spectra and crystal structure of $(Ph_4P)[B_6H_6CH_2Ph]$, **289**, 70
the age of chiral deltahedral borane derivatives, **289**, 45
Density functional calculations
relativistic density functional study on the bimetallic cluster $[Pt_3Fe_3(CO)_{15}]^{n-}$ (n = 0, 1, 2), **286**, 30
Density functional theory
a dinuclear platinum(II) complex with hydrazine and chloro bridges: $(NPr^n_4)[PtCl_2(\mu\text{-}Cl)(\mu\text{-}N_2H_4)PtCl_2]$, **284**, 103
density functional theory and fluorocarboranes: I. Trends in B–H and B–F distances and dissociation energies for $CB_{11}H_{12-n}F_n^-$ anions (n = 0, 1, 6, 11), **294**, 193
formation of FeSi and $FeSi_2$ films from *cis*-$Fe(SiCl_3)_2(CO)_4$ by MOCVD–precursor versus substrate control, **291**, 380

theoretical study of photochromic dinuclear molybdenum complexes, **295**, 180
Deposition mechanism
formation of FeSi and $FeSi_2$ films from *cis*-$Fe(SiCl_3)_2(CO)_4$ by MOCVD –precursor versus substrate control, **291**, 380
Desilylation
cleavage of substituents upon oxidative polymerization of substituted thiophenes, **287**, 232
Diamide complexes
crystallographic and ESR studies of π–π stacked binuclear copper(II) complexes of 1,3-bis(quinoline-2-carboxamido) and 1,3-bis(isoquinoline-3-carboxamido)propane, **288**, 90
Diamine complexes
base hydrolysis of $[Co(triamine)(diamine)Cl]^{2+}$ and epimerization of $[Co(triamine)(diamine)OH]^{2+}$: studies on resolved cobalt(III) complexes of 3-aza-hexane-1,6-diamine (2,3-tri), **284**, 37
cationic 16-electron half-sandwich ruthenium complexes containing asymmetric diamines: understanding the stability and reactivity of coordinatively unsaturated two-legged piano stool complexes, **286**, 114
facile synthesis and structural properties of (diamine)-tetracarboxylatoplatinum(IV) complexes, **292**, 52
stereochemical investigation of a kinetically labile chiral organopalladium complex in solution, **284**, 99
structural and spectroscopic characterisation of the mixed valence platinum complex $PtdabI_3$ (dab = *o*-diaminobenzene), **287**, 218
synthesis and properties of (diamine)platinum(II) complexes of pyridine carboxylate isomers and their antitumor activity, **292**, 238
Diammine complexes
[*meso*- and *rac*-1,2-bis(4-fluorophenyl)ethylenediamine]chloro-[sulfinylbis(methane)-S]platinum(II) chloride new water soluble platinum complexes with high anti-breast cancer activities, **285**, 184
Diarsine complexes
multinuclear NMR studies of diphosphine, diphosphine–dioxide and diarsine complexes of tin(IV) halides. Structures of $[SnI_4\{o\text{-}C_6H_4(AsMe_2)_2\}]$ and $[SnI_4\{o\text{-}C_6H_4(P(O)Ph_2)_2\}]$, **288**, 142
Diastereoselective aldol additions
synthesis of the first chiral phosphorous chelate aminophosphino–carbene complex and its applications in stereoselective addition reactions, **296**, 236
Diastereoselectivity
diastereoselective *ortho*-lithiation of tricarbonylchromium complexes of chiral benzaldehyde acetals, **296**, 80
Diaza-crown complexes
stability constants of copper(I) and silver(I) complexes with open-chain, macrocyclic and -bicyclic aza-ligands in acetonitrile and comparison with results in dimethylsulfoxide, **286**, 160
Diazadiene complexes
comparing structures and reactivity in analogous Fe and Ru complexes. (^{i}Pr-DAB)$Fe(CO)_2I_2$ and (^{i}Pr-DAB)FeI_2: a perfectly reversible CO-carrier system. (R-DAB = *N*,*N*′-R_2-1,4-diaza-1,3-butadiene), **291**, 438
Diazoalkane complexes
synthesis and structures of tungsten ferrocenyldiazoalkane complexes derived from a dinitrogen complex. Part 58. Preparation and properties of molybdenum and tungsten dinitrogen complexes, **285**, 336
Dibenzofulvalene
novel organoiron compounds resulting from the attempted syntheses of dibenzofulvalene complexes, **291**, 109
Dibenzothiophene
oxidative addition reactions of benzo[*b*]thiophene and dibenzothiophene with $[Os_3(CO)_{10}(MeCN)_2]$: room-temperature ring opening of benzo[*b*]thiophene at triosmium clusters, **285**, 277
Diborane(6)
mechanochemical reactions in the chemistry of boranes, **289**, 51
Diborane compounds
matrix isolation investigation of the reaction of B_2H_6 with carbon–carbon multiple bonds, **286**, 1
1,3-Dicarbonyl group complexes
transformation of the carboxyl group adjacent to the (η^4-diene)$Fe(CO)_3$ moiety: a convenient route to (η^4-diene)$Fe(CO)_3$ complexes containing α,β- and β,γ-unsaturated carbonyl groups, and 1,3-dicarbonyl groups, **291**, 231
Dicarboxylic acid complexes
syntheses and anti-*Candida* activity of cobalt(II) complexes of octanedioic acid ($odaH_2$) and nonanedioic acid ($ndaH_2$); X-ray crystal structures of $[Co(phen)_3]oda{\cdot}14H_2O$ and $[Co(phen)_3]nda{\cdot}11.5H_2O$ (phen = 1,10-phenanthroline), **293**, 160
Diels–Alder reactions
reactions of cobaloximes with vinyl allenes as a route to 1-cobaloxime substituted 1,3-dienyl complexes, **296**, 13
Diene complexes
synthesis and characterization of siloxy-substituted diene and dienyl complexes of the $Ru(C_5Me_5)$ fragment, **296**, 170
transformation of the carboxyl group adjacent to the (η^4-diene)-$Fe(CO)_3$ moiety: a convenient route to (η^4-diene)$Fe(CO)_3$ complexes containing α,β- and β,γ-unsaturated carbonyl groups, and 1,3-dicarbonyl groups, **291**, 231
(1,3-Diene)iron complexes
synthesis and reactivity of acyclic (pentadienyl)iron(1 +) cations: model studies for the preparation of the 8*E*,10*Z*,16*E*,18*E*-tetraene segment of macrolactin A, **296**, 261
Diethylenetriamine complexes
studies on Pt–S bonds. Complexes of methylsulfanyl- and methylsulfynil-acetate, benzoate and phenolate with the diethylenetriamineplatinum(II) moiety, **285**, 249
Dihydrogen complexes
reactions of H_2, silanes, and olefins with superelectrophilic cationic rhenium complexes: heterolytic cleavage of H_2 and relation to the structure and function of hydrogenases, **294**, 240
Dihydroxybenzoic acid complexes
coordination mode of dihydroxybenzoic acids in oxorhenium(V) complexes, **294**, 114
Diimine complexes
electronic spectra of 1,2-diiminetricarbonylrhenium(I)chloride complexes with imidazole derivatives as ligands, **288**, 150
syntheses, spectroscopic properties and photochemistry of [Re(R)$(CO)_3$(4,4′-Me_2-bpy)] (R = alkyl) complexes, **284**, 61
Diimine ligand complexes
synthesis, crystal structure and magnetic behaviour of the new linear copper(II) tetranuclear compound $[Cu_4(\mu_2\text{-}AcO)_6(AcO)_2(DPQ)_2]$ (DPQ = 2,3-bis(2-pyridyl)quinoxaline), **286**, 62
Diiron complexes
photochemical reactions of transition metal organyl complexes with olefins: XXII. Photoreactions of pentacarbonyl(μ-$\eta^{5:3}$-4*H*-indenyl)diiron(Fe–Fe) with alkynes—substituent controlled C–H transfer or C–C bond formation, **296**, 127
Diketonate complexes
high coordination numbers of metal ions in chelate complexes: molecular adducts of scandium tris-α-diketonates in solution, **286**, 175

β-Diketone complexes
synthesis and characterization of a metal coordination polymer consisting of ruthenium(III) β-diketone units linked by butadiyne bridges, **292**, 64
Dimer complexes
nickel(II) and palladium(II) complexes with acylhydrazone ligands of α-diketones: the electronic and steric factors affecting the formation of the dimeric palladium(II) complexes, **295**, 171
Dimers
crystal structures and magnetic studies of six new nickel(II) amine μ-oxalato compounds, **295**, 91
Dimetallic complexes
novel organoiron compounds resulting from the attempted syntheses of dibenzofulvalene complexes, **291**, 109
Dimethylpyrazole complexes
deboronation of $HB(3,5\text{-}Me_2C_3HN_2)_3^-$ in a reaction with $[Re(O)Cl_4]^-$, the X-ray crystal structures of $[\{Re(O)Cl(3,5\text{-}Me_2C_3HN_2H)_3\}(\mu\text{-}O)\{Re(O)Cl_3(3,5\text{-}Me_2C_3HN_2H)\}]$ and $[\{Re(O)Cl_2(3,5\text{-}Me_2C_3HN_2H)_2\}(\mu\text{-}O)\{Re(O)Cl_2(3,5\text{-}Me_2\text{-}C_3HN_2H)_2\}]$, **285**, 55
Dinitrogen complexes
synthesis and structures of tungsten ferrocenyldiazoalkane complexes derived from a dinitrogen complex. Part 58. Preparation and properties of molybdenum and tungsten dinitrogen complexes, **285**, 336
Dinuclear complexes
a dinuclear copper(II) complex involving monoatomic *O*-carboxylate bridging and Cu–S(thioether) bonds: $[Cu(tda)(phen)]_2 \cdot H_2tda$ (tda = thiodiacetate, phen = phenanthroline), **286**, 74
crystallographic and ESR studies of π–π stacked binuclear copper(II) complexes of 1,3-bis(quinoline-2-carboxamido) and 1,3-bis(isoquinoline-3-carboxamido)propane, **288**, 90
crystal structure of (μ-oxo)bis{aquatris(2-pyridylmethyl)amine}-iron(III) perchlorate, **284**, 124
cyano-bridged dinuclear iron complexes in different oxidation states, **287**, 95
dimolybdenum complexes of bis(2-pyridyl)amine (Hdpa) ligand and its anion (dpa^-); transformation of chelating η^2-Hdpa to bridging η^2-dpa^-, **288**, 82
diphenoxo-bridged dinuclear nickel(II) complex of a dinucleating macrocycle with $N(amine)_6O_2$ metal-binding sites: synthesis, X-ray crystal structure and electrochemistry, **287**, 212
electrochemical and ^{95}Mo NMR studies of triply-bridged dinuclear oxomolybdenum(V) complexes with various dithiocarbamates, **292**, 7
formation pathway of a Mn(IV),Mn(IV) bis(μ-oxo) dimer that incorporates alkali and alkaline earth cations and electron transfer properties of the dimer, **294**, 133
magneto–structural correlation of dimeric copper(II) carboxylates with pyridyl-substituted nitronyl nitroxides, **293**, 30
molecular recognition of dipeptides. Catalysis of deuteration and hydrolysis of glycylglycine by dinuclear OBISDIEN Zn(II) complexes, **287**, 134
nitrite ion-promoted hydration of acetonitrile by the bis(μ-hydroxo)bis{tris(2-pyridylmethyl)amine)chromium(III)} ion, **288**, 211
palladium(I) phosphido-bridged dinuclear derivatives with both phosphine and amine ligands, **284**, 246
preparation, structure and electrochemical behavior of dinuclear cyclooctadiene-chelated Ir(I) complexes with 2-aminopyridinato bridges, **292**, 244
reactions of σ-π alkenyl complexes $[Fe_2(CO)_6(\mu\text{-alkenyl})(\mu\text{-}PPh_2)]$ with bis(diphenylphosphino)methane (dppm); carbonyl substitution, migratory carbonyl insertion, phosphorus–carbon bond formation and cleavage, proton shift and alkenyl isomerisation reactions, **291**, 178
reactivity towards some weak protic electrophiles of the di-μ-hydroxo complex $[\{Pd(\mu\text{-}OH)(tmeda)\}_2][BPh_4]_2$: crystal structure of $[\{Pd(tmeda)\}_2(\mu\text{-}OH)(\mu\text{-}HNC_6H_4Cl\text{-}p)][BPh_4]_2$, **287**, 37
syntheses, crystal structures and autoreduction behavior of antiferromagnetically coupled dicopper(II) oximato complexes, **293**, 20
synthesis of a dinuclear tungsten complex, $W_2(\mu\text{-}H)(\mu\text{-}OC_2H_5)(\mu\text{-}O_2CC_6H_5)(\eta^1\text{-}OC_2H_5)_2Cl_2(\eta^2\text{-}P,P'\text{-dppp})$ where dppp is 1,3 bis(diphenylphosphino) propane, **284**, 303
synthesis of diphenylphosphine palladium complexes, **284**, 116
synthesis, reactivity, and structures of (μ-acetamidato)-hexakis(acetonitrilo)dimolybdenum(II) tris(tetrafluoroborate) and derivatives, **287**, 159
the effect of replacing carbonyl with acetonitrile on the reactivity of the bimetallic system $[Mo_2Cp_2(\mu\text{-}SMe)_3(L)_2]^+$ (L = CO, CH_3CN). Crystal structure of $[Mo_2Cp_2(\mu\text{-}SMe)_2(CH_3CN)_4](BF_4)_2$, **284**, 107
the μ-sulfonium–methylidene diiron complexes $[Fe_2\{\mu\text{-}C(X)SMe_2\}(\mu\text{-}CO)(CO)_2(Cp)_2]SO_3CF_3$ (X = CN, H) as precursors of μ-alkylidene complexes, **291**, 333
two diphenylformamidinate compounds of the quadruply-bonded Re_2^{6+} core, **285**, 149
versatility of the cyanato ligand: structure and magnetic properties of four new copper(II)–cyanato compounds derived from tridentate amines, **286**, 189
Dinuclear molybdenum complexes
theoretical study of photochromic dinuclear molybdenum complexes, **295**, 180
Dioxolene complexes
dinuclear complexes of a new bridging ligand containing 2,2′-bipyridyl and dioxolene binding sites: syntheses, electrochemical and electronic spectroscopic properties, **285**, 89
Dioxygen complexes
catalytic oxidation of 1,2-diaminobenzene by the dinuclear cobalt(II)–OBISDIEN dioxygen complex, **287**, 193
equilibrium studies of reactions of 4-chloro-1,2-diaminobenzene, 4-methyl-1,2-diaminobenzene, 4-methoxy-1,2-diaminobenzene and 4,5-dimethyl-1,2-diaminobenzene with dinuclear cobalt(II) 1,4,7,13,16,19-hexaaza-10,22-dioxacyclotetracosane, **284**, 30
Dipeptide complexes
molecular recognition of dipeptides. Catalysis of deuteration and hydrolysis of glycylglycine by dinuclear OBISDIEN Zn(II) complexes, **287**, 134
reaction of formaldehyde with cyano(ethylenediamine)-(glycylglycinato)cobalt(III): crystal structures of cyano-(ethylenediamine) [*N*-(4-carbonyl-4-hydroxymethyl-oxazolidine)glycinato]cobalt(III)·$3.5H_2O$ and dicyanobis(ethyl-enediamine) {μ-*N*-[*N*-(((methylimino)methyl)carbonyl)-α-methylserinato)glycylglycinato}dicobalt(III) tetrahydrate, **292**, 96
Diphenylformadinate complexes
two diphenylformamidinate compounds of the quadruply-bonded Re_2^{6+} core, **285**, 149
Diphenylphosphine complexes
synthesis of diphenylphosphine palladium complexes, **284**, 116
Diphosphine complexes
C–S, C–H, and N–H bond cleavage of heterocycles by a zero-valent iron complex, $Fe(N_2)(depe)_2$ [depe = 1,2-bis-(diethylphosphino)ethane], **291**, 341
ligand substitution kinetics in $M(CO)_4$ ($\eta^{2,2}$-1,5-cyclooctadiene) complexes (M = Cr, Mo, W) — substitution of 1,5-cyclooctadiene by bis(diphenylphosphino)alkanes, **284**, 229

multinuclear NMR studies of diphosphine, diphosphine–dioxide and diarsine complexes of tin(IV) halides. Structures of $[SnI_4\{o\text{-}C_6H_4(AsMe_2)_2\}]$ and $[SnI_4\{o\text{-}C_6H_4(P(O)Ph_2)_2\}]$, **288**, 142
synthesis, characterization and crystal structures of rhenium(V) complexes with diphosphines, **294**, 47
synthesis, platinum(II) complexes and structural aspects of the new tetradentate phosphine *cis,trans,cis*-1,2,3,4-tetrakis-(diphenylphosphino)cyclobutane, **290**, 167

Diphosphinito-bridged complexes
diphosphinito-bridged ruthenium(II) and rhodium(III) complexes with stereogenic metal centers, **292**, 84

Diphosphinomethane complexes
synthesis and structure of niobium and tantalum derivatives of bis(dicyclohexylphosphino)methane (dcpm), **288**, 35

Diphosphite
synthesis of chiral diphosphite ligands and comparison of their rhodium and platinum complexes in the asymmetric hydroformylation of styrene, **296**, 222

Dipyridylamine complexes
dipyridylamine-ligated octamolybdate clusters aggregated into extended ribbons and layers via hydrogen bonding, **292**, 131

Diselenoimidodiphosphinato complexes
displacement of triphenylphosphine from $Cu(PPh_3)_2NO_3$ and $Co(PPh_3)_2Cl_2$ by a diselenoimidodiphosphinato ligand. X-ray crystal structure of $(PPh_3)Cu[Ph_2P(Se)NP(Se)Ph_2]$ and $Co[Ph_2P(Se)NP(Se)Ph_2]_2$ containing the novel $CuSe_2P_2N$ and $CoSe_2P_2N$ inorganic metallocycles, **290**, 256

1,3-Disilacyclohexanes
a facile synthesis of 1,3-disilacyclohexanes, **296**, 231

Dispersion curves
lattice vibrations of the icosahedral solid boron arsenide, $B_{12}As_2$, **289**, 198

Dissociation energies
density functional theory and fluorocarboranes: I. Trends in B–H and B–F distances and dissociation energies for $CB_{11}H_{12-n}F_n^-$ anions ($n = 0, 1, 6, 11$), **294**, 193

Disulfoxide complexes
bridging properties of disulfoxide ligands: crystal structure of copper(1,3-bis(n-propylsulfinyl)propane) perchlorate, **292**, 144

Dithiocarbamate complexes
an EXAFS study of Mo(VI) and Mo(V) complexes with proline dithiocarbamate, **288**, 127
electrochemical and ^{95}Mo NMR studies of triply-bridged dinuclear oxomolybdenum(V) complexes with various dithiocarbamates, **292**, 7
structure and characterization of a mononuclear Fe^{II} dialkyldithiocarbamate complex $Fe(OC_4H_8dtc)_2(DMF)$, **293**, 95
trinuclear vanadium dialkyldithiocarbamate complexes. Synthesis, structure and spectroscopic characterization of $[Et_4N][V_3(\mu_3\text{-}S)(\eta^2\text{-}\mu\text{-}S_2)_3(Et_2dtc)_3] \cdot 2CH_3CN$, **286**, 7

Dithiocarbazate complexes
synthetic, spectroscopic and X-ray crystallographic structural studies on some copper(II) complexes of the 6-methylpyridine-2-carboxaldehyde Schiff base of *S*-methyldithiocarbazate, **287**, 8

Dithiocyanato complex
synthesis and photophysical properties of *trans*-dithiocyanato bis(4,4′-dicarboxylic acid-2,2′-bipyridine) ruthenium(II) charge transfer sensitizer, **296**, 250

Dithioether ligands
synthesis, reactivity and catalytic properties of rhodium complexes of (*R,R*)-1-benzyl-3,4-dithioetherpyrrolidines, **295**, 64

Dithiolate complexes
a novel copper(I) cluster stabilized by a dithiolate ligand having a conjugated TTF moiety, **285**, 332
crystallographic evidence for the coordination performance of $WO_2S_2^{2-}$. Synthesis, characterization and crystal structure of palladium complexes with mixed-ligand thiotungstates and 1,2-dithiolates: $[NBu_4]_2[PdWO_2S_2(C_3S_5)]$ and $[NBu_4]_2[PdWS_4(C_4N_2S_2)]$, **285**, 241
molecular structure of $[PPh_4]_2[Co(C_3S_5)_2]$ [$C_3S_5^{2-}$ = 4,5-disulfanyl-1,3-dithiole-2-thionate(2-)] and spectroscopic and electrical properties of its oxidized species, **295**, 239
properties of Pt(II) complexes containing both a pyridyl N–N chelate ligand having a long alkyl chain and a sulfur-rich dithiolate ligand and their molecular interactions in the solid state, **284**, 55

Dithiolato complexes
non-linear optical absorption of $[\{Ni(mnt)_2\}_2]^{2-}$ and its unique counter ion, a disodium triple-decker dication $[Na_2(benzo\text{-}15\text{-}crown\text{-}5)_3(H_2O)_2]^{2+}$, **293**, 89

Dithio-*o*-carboranyl complexes
synthesis, X-ray crystal structure, and reactivity of the monomeric dithio-*o*-carboranyl iridium complex $[Ir(\eta^5\text{-}C_5Me_5)(\eta^2\text{-}S_2C_2B_{10}H_{10})]$, **289**, 141

Divalent metal complexes
3-hydroxypropylation of tetraazamacrocycles and its effect on the stability of divalent metal complexes, **294**, 103

DNA cleavage
synthesis, spectroscopic, redox properties, and DNA cleavage activity of low-spin iron(III) complexes of bleomycin analogues, **294**, 56

DNA determination
spectroscopic characterization of porphyrin supramolecular aggregates on poly-lysine and their application to quantitative DNA determination, **286**, 121

Double allylation of carboxylic ester
Barbier-type allylation of carbonyl derivatives by use of aluminum as an electron pool. Double allylation of carboxylic esters, **296**, 204

Double bond reduction
intramolecular coupling of conjugated enynes with cyclopropylcarbene–chromium complexes; a complex reaction pathway, **296**, 273

Dynamic NMR
synthesis and dynamic NMR studies of stereochemical nonrigidity in rhenium(I) complexes of 2,6-bis[1-(phenylimino)-ethyl]pyridine derivatives, **295**, 56

Edta complexes
crystal structures and stereochemical properties of lanthanide(III) complexes with ethylenediamine-*N,N,N′,N′*-tetraacetate, **288**, 7
detection of N-3 and N-7-coordinated $[Ru^{II}(edta)(5'\text{-}GMP)]^{4-}$ complexes and the N-1 protonation equilibrium of the Ru^{III} derivative, **285**, 170
intervalence transfer properties of the binuclear μ-benzotriazolate- and μ-benzimidazolate-bis{ruthenium(II)/(III)-edta} complexes, **285**, 197

Edta-type complexes
hexadentate chromium(III) complexes containing unsymmetrical edta-type ligands: Part IV. Crystal structures of $(-)_{589}$-*trans*(O_5)-Na[Cr(ed3ap)]·$3H_2O$ and Na[Cr(*u*-eddadp)]·$3H_2O$ and CD spectra correlation: structural parameters of [Cr(edta-type)]$^-$ complexes and their octahedral distortion in relation to the structure of the ligand, **292**, 16

Electrical conductivity
molecular structure of $[PPh_4]_2[Co(C_3S_5)_2]$ [$C_3S_5^{2-}$ = 4,5-disulfanyl-1,3-dithiole-2-thionate(2-)] and spectroscopic and electrical properties of its oxidized species, **295**, 239
properties of Pt(II) complexes containing both a pyridyl N–N chelate ligand having a long alkyl chain and a sulfur-rich dithiolate ligand and their molecular interactions in the solid state, **284**, 55
Electrical resistivity
B and B–C as interstitials in reduced rare earth halides, **289**, 174
Electrochemical synthesis
electrochemical synthesis of copper and silver pyridine-2-selenolato complexes: crystal structure of (6-trimethylsilylpyridine-2-selenolato)silver(I), $[Ag_6(6\text{-}Me_3SipySe)_6]$, **284**, 133
Electrochemistry
cadmium(II) complexes with [(4-methylphenyl)sulfonyl]-2-pyridylamine: crystal structures of bis{[(4-methylphenyl)-sulfonyl]-2-pyridyl-amide}cadmium(II); aquo-bis(pyridine) bis{[(4-methylphenyl)sulfonyl]-2-pyridyl-amide}cadmium(II) and 1,10-phenanthroline bis{[(4-methylphenyl)sulfonyl]-2-pyridyl-amide}cadmium(II), **294**, 87
complexation of macrocyclic ligands with relatively non-solvated metal ions generated in dichloromethane by electrochemical oxidation of amalgam electrodes, **293**, 223
dinuclear complexes of a new bridging ligand containing 2,2′-bipyridyl and dioxolene binding sites: syntheses, electrochemical and electronic spectroscopic properties, **285**, 89
diphenoxo-bridged dinuclear nickel(II) complex of a dinucleating macrocycle with $N(amine)_6O_2$ metal-binding sites: synthesis, X-ray crystal structure and electrochemistry, **287**, 212
electrochemical and ^{95}Mo NMR studies of triply-bridged dinuclear oxomolybdenum(V) complexes with various dithiocarbamates, **292**, 7
electrochemical synthesis of cobalt, nickel, copper, zinc and cadmium complexes with *N*[(2-hydroxyphenyl)methylidine]-*N*′-tosylbenzene-1,2-diamine. The crystal structures of {(1,10-phenanthroline)[*N* - (2 - oxophenyl)methylidine] - *N*-tosylbenzene-1,2-diaminato}nickel(II) and {(1,10-phenanthroline)[*N*-(2-oxophenyl)methylidine]-*N*′-tosylbenzene-1,2-diaminato}copper(II), **295**, 39
electrochemical tuning of indium phosphide|poly(acetylene) interfaces, **294**, 207
electrochemistry of Au-complexes, **290**, 44
evidence for a charge-transfer intermediate in the catalytic epoxidation of alkenes by ruthenium complexes: a comparison between electrochemical and kinetic data, **288**, 122
novel features associated with the electrochemically driven bis(η^5-pentaphenylcyclopentadienyl)iron(II)–iron(III) redox transformation at an electrode–microcrystal–solvent (electrolyte) interface, **291**, 21
preparation and characterization of various *N*-substituted-2-aza-[3]-ferrocenophanes and their chemical and electrochemical properties, **296**, 176
preparation, structure and electrochemical behavior of dinuclear cyclooctadiene-chelated Ir(I) complexes with 2-aminopyridinato bridges, **292**, 244
solution studies of complexes of iron(III) with iminodiacetic, alkyl-substituted iminodiacetic and nitrilotriacetic acids by potentiometry and cyclic voltammetry, **291**, 158
structural study of the electrochemically synthesized binuclear complex bis{1-phenylglyoxal bis(3-piperidylthiosemicarbazone)zinc(II)}, **294**, 79
synthesis and electrochemical properties of a new C_{60}-phenanthroline derivative and its dimeric complex with silver, **292**, 34
synthesis, structure, and redox properties of the sulfido bridged tetranuclear cluster complex $[(Ph_3P)_2N][Fe_3(CO)_9\{\mu_3\text{-}SFeCp(CO)_2\}]$, **286**, 142
the relationship between surface derivatization and the physical properties of bisarylphosphinidenetriironnonacarbonyl clusters, **294**, 140
vitamin B_{12}–transport protein interaction: electrochemistry of aquo- and glutathionyl-cobalamins adsorbed on carbon electrodes; role of the nucleotide chain, **292**, 204
Electronic spectra
potentiometric and spectroscopic studies of reactions of histidine with 13-membered amide-based macrocyclic Cu^{2+} chelates, **295**, 115
Electronic structure
early versus late transition metals. Electronic structure of *nido*-2-$CpML_nB_4H_8$, $CpML_n$ = $CpTaCl_2$, $CpWH_3$ and CpCo, **289**, 85
iron and ruthenium organometallic derivatives of spiro-substituted cyclotriphosphazenes bearing nitrile spacer: synthesis and electronic structure, **294**, 20
relativistic density functional study on the bimetallic cluster $[Pt_3Fe_3(CO)_{15}]^{n-}$ (n = 0, 1, 2), **286**, 30
Electron transfer
effects of β-cyclodextrin inclusion on the kinetics of the oxidation of bisferrocenyl cations by bis(pyridine-2,6-dicarboxylato)cobaltate(III) in aqueous solution: crystal structure of dimethyldi-(1-methylferrocene)ammonium bromide, **291**, 365
effects of electronic structure on chiral induction in outer-sphere electron transfer reactions of complexes with the C_3 symmetric ligand, 1,4,7-triazacyclononane-1,4,7-tris[2′(*R*)-2′-propionate](-3), **290**, 197
organometallic electron reservoir sandwich iron complexes as potential agents for redox and electron transfer chain catalysis, **291**, 1
Electrospray mass spectrometry
complexes of lanthanide dithionates 2. Electrospray mass spectrometry of $Ln_2(S_2O_6)_3$ and their Ph_3PO complexes; the X-ray crystal structure of $Pr_2(S_2O_6)_3(Ph_3PO)_6(H_2O)_6$, **295**, 146
electrospray mass spectrometric analysis of transition-metal halide complexes, **294**, 183
electrospray mass spectrometric investigation of the relative ligand properties of EPh_3 ligands (E = P, As, Sb or Bi) towards Ag^+ and Cu^+, **290**, 213
on tuning the copper(I) coordination number in halocuprate(I) anions: new insights into cation control, **292**, 266
Enantioselective catalysis
highly enantioselective catalytic isopropenylation of aldehydes using diisopropenylzinc, **296**, 33
Enantioselective reduction
optically active β-ketoiminato cobalt(II) complexes as highly efficient catalysts in enantioselective borohydride reduction of ketones, imines, and α,β-unsaturated carboxylates, **296**, 86
Ene reactions
molecular assembly of BINOL–Ti complexes into an active μ_3-oxo titanium catalyst, **296**, 267
Enterobactin analog complexes
thermodynamic and kinetic studies of the sulfonated derivative of the iron chelator TRENCAM, an analog of enterobactin, **291**, 148
Epimerization
base hydrolysis of $[Co(triamine)(diamine)Cl]^{2+}$ and epimerization of $[Co(triamine)(diamine)OH]^{2+}$: studies on resolved cobalt(III) complexes of 3-aza-hexane-1,6-diamine (2,3-tri), **284**, 37
Epoxidation
evidence for a charge-transfer intermediate in the catalytic epoxidation of alkenes by ruthenium complexes: a comparison between electrochemical and kinetic data, **288**, 122

synthesis and properties of hydrophobic $[Mn^{IV}_2(\mu\text{-}O)_3(L)_2]^{2+}$ complexes, derived from alkyl substituted 1,4,7-triazacyclononane ligands, **295**, 189

EPR
copper(I) and copper(II) complexes of the bidentate imidazole/thioether ligand 1-methyl-2-(methylthiomethyl)-1*H*-benzimidazole, **287**, 204
reduction of the NO^+ ligand in the pentacyanonitrosylosmate(II) ion, **285**, 129

Equilibrium constants
potentiometric determination of the equilibrium constants for the formation of Fe(III)–sorbitol complexes, **293**, 206

Equilibrium study
potentiometric and spectroscopic studies of copper(II) complexes of bis(1,4,7-triazacyclononane) ligands containing polymethylene and xylyl linker groups, **292**, 172

ESMS
interaction of beryllium(II) in aqueous solution with bidentate ligands containing phosphonate groups, **285**, 39

Ethene
palladium(II) complexes with hemilabile etherdiphos ligands in the alternating copolymerization of carbon monoxide with olefins, **296**, 103

Ethylenediamine complexes
synthesis and biological activity of *cis*-dichloro mono- and bis(platinum) complexes with *N*-alkyl-ethylenediamine ligands, **290**, 237
the first X-ray crystal structures of cobalt complexes containing monodentate and bridging ethylenediamine ligands, **288**, 53

Europium complexes
neodymium and europium picrate complexes with 3-picoline-*N*-oxide (3-picNO), **292**, 249

EXAFS
an EXAFS study of Mo(VI) and Mo(V) complexes with proline dithiocarbamate, **288**, 127

Extracoordinate silanes
stereoselective reduction of ketones using extracoordinate silicon: C_2-symmetric ligands, **296**, 208

Extraction equilibria
complexation of uranyl ion by tetrahexylmalonamides: an equilibrium modeling and infrared spectroscopic study, **293**, 195

FAB mass spectra
triethanolamine complexes of copper, **294**, 153

4f complexes
mononuclear lanthanide complexes of tripodal ligands: synthesis and spectroscopic studies, **285**, 49

FeMo–cofactor
the first binuclear Mo–S cluster compound containing citrate ligands, $K_5(NH_4)[Mo_2O_2(\mu\text{-}S)_2(C_6H_4O_7)_2]\cdot CH_3OH\cdot 5H_2O$, characterized by X-ray single crystal structure determination, **285**, 152

Ferrate ion
kinetics and mechanism of the reduction of ferrate by one-electron reductants, **293**, 229

Ferrocene
synthesis, structure, and reactivity of new chiral ferrocenyl pyrazole ligands, **296**, 183
the quadrupole hyperfine interaction of iron organometallics: an interpretation of its temperature dependence, **291**, 74

Ferrocene complexes
coordination chemistry and X-ray crystal structure of (ferrocenylmethyl)diphenylphosphine, $FcCH_2PPh_2$ [$Fc = (\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4)$], **295**, 18
germanium-containing ferrocenes and ferrocenophanes. Potential precursors for ring-opening polymerizations and 'germaferrocenes', **291**, 355
synthesis, X-ray structural determination and Mössbauer characterization of Schiff bases bearing ferrocene groups, their reduced analogues and related complexes, **287**, 117

Ferrocenophane
preparation and characterization of various *N*-substituted-2-aza-[3]-ferrocenophanes and their chemical and electrochemical properties, **296**, 176

Ferrocenophane complexes
germanium-containing ferrocenes and ferrocenophanes. Potential precursors for ring-opening polymerizations and 'germaferrocenes', **291**, 355

[2]Ferrocenophanes
strained, ring tilted dicarbon-bridged [2]ferrocenophanes and ferrocene revisited: ^{57}Fe Mössbauer spectroscopic study of bonding, hyperfine interactions, and lattice dynamics, **287**, 152

Ferrocenyl complexes
synthesis and structures of tungsten ferrocenyldiazoalkane complexes derived from a dinitrogen complex. Part 58. Preparation and properties of molybdenum and tungsten dinitrogen complexes, **285**, 336
synthesis of (ferrocenyl-indenyl)cyclopentadienyliron compounds with and without a bridging group via a CpFe transfer reaction, **286**, 215
synthesis of gold ferrocenyl substituted ylides and methanides: crystal structure of $[FcCH(AuPPh_3)PPh_3]ClO_4$ [$Fc = (\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4)$], **291**, 60

Five-coordinate complexes
five-coordinate rhenium(III)-thiolato complexes: the structure of $[Re(SCH_2PhOCH_3)_3(PPh_3)_2]$, **290**, 247
synthesis of five-coordinate Pt(II) complexes containing organogermanium fragments in axial position. Structure of $[PtCl(GePh_2Cl)(2.9\text{-}Me_2\text{-}1,10\text{-}phen)(CH_2{=}CH_2)]$, **285**, 70
synthesis of the ligand *N*,*N*′-bis(2-tert-butylthiobenzenylidene)-diethylenetriamine; its reactivity with nickel(II) salts, **287**, 105

Five coordination
the syntheses, spectra and structures of five-coordinate cobalt(II) complexes of pyrazolyl-containing ligands, **284**, 266

Flash photolysis
volume and thermodynamic profiles of CO-binding to Fe(II) protoporphyrin IX in detergent micelles, **288**, 74

Fluorenyl complexes
a new dioxotetraamine ligand appended with fluorenyl and its copper(II) complex. Synthesis, crystal structure and solution behavior, **295**, 48

Fluorescence
a new dioxotetraamine ligand appended with fluorenyl and its copper(II) complex. Synthesis, crystal structure and solution behavior, **295**, 48
coordination complexes of bifunctional compounds: I. Synthesis and properties of bis[5-(2-oxyphenyl)-3-phenyl-1,2,4-oxadiazolyl]copper(II). A fluorescent coordination compound of Cu(II), **292**, 1

Fluoride complexes
solvatochromism in ligand-field absorption bands and solvent-dependent deuteron nuclear magnetic resonance spectra of $[CrF_3(tri(2\text{-}pyridyl)amine)]$: implication of trigonal splitting for the first spin-allowed d–d band and evaluation of π bonding peculiarities, **295**, 84

Fluorination
synthesis and characterization of the anionic fluorocarboranes 6,7,8,9,10-$CB_9H_5F_5^-$, 6,7,8,9-$CB_9H_5F_4$-10-OH^-, and 6,7,8,9-$CB_9H_5F_4$-10-$NHCOCH_3^-$, **289**, 76

Fluoroalkoxy P-ligand complexes
solution thermochemical study of ligand substitution reactions of hybrid alkyl/fluoroalkoxy phosphorus ligands in the $L_2Fe(CO)_3$ system, **291**, 32

Fluorocarborane
density functional theory and fluorocarboranes: I. Trends in B–H and B–F distances and dissociation energies for $CB_{11}H_{12-n}F_n^-$ anions ($n = 0, 1, 6, 11$), **294**, 193
Fluorocarboxylate complexes
preliminary studies of $Ni(dmen)_2(CF_3CO_2)_2$, $Ni(deen)_2(CF_3CO_2)_2$ and $Ni(en)(CF_3C(O)CHC(O)CF_3)_2$ as precursors for metalorganic chemical vapour deposition: structures, thermal behaviours and vapour pressures, **288**, 233
Formation constants
3-hydroxypropylation of tetraazamacrocycles and its effect on the stability of divalent metal complexes, **294**, 103
potentiometric and spectroscopic studies of reactions of histidine with 13-membered amide-based macrocyclic Cu^{2+} chelates, **295**, 115
thermodynamic and kinetic studies of the sulfonated derivative of the iron chelator TRENCAM, an analog of enterobactin, **291**, 148
Formato complexes
coordination compounds with 4-aminopyridine. An unusual in situ formation of a bridging formato compound. The structure of *catena*-poly[bis(4-amino-pyridine)(formato-*O*)-copper(II)-μ-formato(*O*,*O'*)], **287**, 226
Fullerenes complexes
synthesis and electrochemical properties of a new C_{60}-phenanthroline derivative and its dimeric complex with silver, **292**, 34
Furanone
rhodium-catalyzed cyclic carbonylation of 2-phenylethynylbenzamide under water–gas shift reaction conditions, **296**, 195

Gadolinium complexes
interaction of ATP with a Gd^{3+} complex employed as paramagnetic contrast agent in NMR imaging, **288**, 244
Gallium trichloride
β-silylethenylation of phenol with silylethyne using gallium trichloride, **296**, 67
Geometry optimization
theoretical study of photochromic dinuclear molybdenum complexes, **295**, 180
Germanium complexes
germanium-containing ferrocenes and ferrocenophanes. Potential precursors for ring-opening polymerizations and 'germaferrocenes', **291**, 355
Glasses
NMR and NQR studies of boron in vitreous and crystalline borates, **289**, 158
Glycine derivative complexes
a series of heteroleptic complexes of the type *fac*-$[Mn^{III}L_2]^-$ [H_2L = derivatives of *N*-(2-hydroxybenzyl)glycine or *N*-(5-nitro-2-hydroxybenzyl)sarcosine] possessing unusual Mn(III) co-ordination spheres, **290**, 228
GMP complexes
detection of N-3 and N-7-coordinated $[Ru^{II}(edta)(5'\text{-GMP})]^{4-}$ complexes and the N-1 protonation equilibrium of the Ru^{III} derivative, **285**, 170
Gold complexes
carbosilane dendrimers functionalized with $AuFe_3$ clusters, **291**, 247
electrochemistry of Au-complexes, **290**, 44
structure and DNA binding properties of the gold(III) complex $[AuCl_2(esal)]$, **285**, 309
syntheses and X-ray structures of mixed-metal gold phosphine clusters including an example having a highly asymmetric Re_2Au_2 skeleton, **294**, 255
synthesis and structural characterization of $[NEt_4][Fe_3(\mu_3\text{-}O)(\mu_3\text{-}AuPPh_3)(\mu\text{-}CO)_3(CO)_6]$, the new $[Au_6(\mu_3\text{-}S)_2(PPh_3)_6][Fe_3(\mu_3\text{-}S)(\mu\text{-}AuPPh_3)(CO)_9]_2$ and $[Au_6(\mu_3\text{-}S)_2(PPh_3)_6][Fe_6(\mu_3\text{-}S)_2(CO)_{14}]$ ionic solids containing assemblages of cluster-cations and cluster-anions, **291**, 372
synthesis of gold ferrocenyl substituted ylides and methanides: crystal structure of $[FcCH(AuPPh_3)PPh_3]ClO_4$ [$Fc = (\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4)$], **291**, 60
the first crystal structure of a complex from a lower rim calix(4)arene alicyclic amino derivative: {tetrakis[1-(ethyl)-piperidinium]oxyl}*p*-tert-butylcalix(4)arene-bis-[tetrachloro-aurate(III)]dichloro dihydrate, **285**, 142
Gold(I) phosphine
the versatile nature of [ClAu(L–L)AuCl] (L–L = bis-(diphenylphosphino)methane (dppm) or -butane (dppb)) in reactions with $[Rh_2Ru_4(CO)_{16}B]^-$: crystal structures of $[\{Rh_2Ru_4(CO)_{16}B\}_2\{\mu\text{-}Au(dppb)Au\}]$ and $[RhRu_4(CO)_{14}B\{Au(dppm)Au\}]$, **289**, 149
Grignard reagent
Grignard reagent formation from aryl halides. There is no aryl radical intermediate along the dominant reaction channel, **296**, 52
Group 4 metal complexes
triphenylsilanolato complexes of Group 4 metals: syntheses, X-ray crystal structures and catalytic cyclization reaction of 1,5-hexadiene, **286**, 14
Guanine mononucleotides
synthesis and characterization of compounds of di- and tributyltin chloride with adenine and guanine mononucleotides, **285**, 60

Hafnium
comparison of the surface chemical reactivity of hafnium diboride and hafnium, **289**, 191
Half-sandwich complexes
cationic 16-electron half-sandwich ruthenium complexes containing asymmetric diamines: understanding the stability and reactivity of coordinatively unsaturated two-legged piano stool complexes, **286**, 114
Halide complexes
electrospray mass spectrometric analysis of transition-metal halide complexes, **294**, 183
monoindenyl titanium alkyl halides. The synthesis and molecular structures of $(\eta^5\text{-}C_9H_7)TiBr_3$, $(\eta^5\text{-}C_9H_7)Ti(CH_3)Br_2$, and $(\eta^5\text{-}C_9H_7)Ti(CH_3)Cl_2$, **292**, 220
nucleophilic substitutions of iodide with bromides in the $Fe(CO)_2(PMe_3)_2MeI$ complex, **288**, 200
structural diversity in thallium chemistry: III. The first structurally characterized examples of the pentabromothallate(III) anion in the 1,1,4,4-tetramethylpiperazinium and *N*,*N'*-diethyltriethylenediammonium salts contrasted with the mixed anions in the 1,1,3,3-tetramethylimidazolidinium salt, **285**, 122
Halocadmate complexes
the hydrothermal synthesis and structural characterization of a new class of compounds: nickel organoamine-halocadmates, **292**, 41
Halocuprate complexes
on tuning the copper(I) coordination number in halocuprate(I) anions: new insights into cation control, **292**, 266
Hapticity change
photochemical reactions of transition metal organyl complexes with olefins: XXII. Photoreactions of pentacarbonyl(μ-$\eta^{5,3}$-4*H*-indenyl)diiron(Fe–Fe) with alkynes—substituent controlled C–H transfer or C–C bond formation, **296**, 127
Helical complexes
multimetallic, double-stranded helical complexes derived from hexa(*n*-propylthio)novipyridine: synthesis, structure and redox properties, **288**, 189

structure and properties of a novel OH^- bridged dimetal helical complex with 6,6''''-dimethyl-2,2':6',2'':6'',2''':6''',2''''-quinquepyridine ligand, **290**, 127

β-Hematin
Mössbauer studies of different types of hematin, **286**, 229

Heme
volume and thermodynamic profiles of CO-binding to Fe(II) protoporphyrin IX in detergent micelles, **288**, 74

Hemilabile ligands
palladium(II) complexes with hemilabile etherdiphos ligands in the alternating copolymerization of carbon monoxide with olefins, **296**, 103

Heptacoordinate complexes
synthesis and structure of a heptacoordinated cadmium(II) complex, **288**, 118

Heptadentate ligands
a seven-coordinate manganese(II) complex formed with the single heptadentate ligand *N,N,N'*-tris(2-pyridylmethyl)-*N'*-(2-salicylideneethyl)ethane-1,2-diamine, **290**, 139

Heptamolybdate
crystal structures of two new heptamolybdates and of a pyrazole incorporating a γ-octamolybdate anion, **295**, 106

Heterocycles
spirocyclic stannole derivatives by 1,1-organoboration of 2,2-bis(1-alkynyl)-1,3-di-tert-butyl-1,3,2-diazastannacyclohexanes, **296**, 26

Heterodinuclear complexes
^1H NMR study of new heterodinuclear macrocyclic complexes, **285**, 233

Heterogeneous catalysis
Kumada couplings catalyzed by nickel on charcoal (Ni/C), **296**, 164

Heterometallic clusters
the versatile nature of [ClAu(L–L)AuCl] (L–L = bis(diphenylphosphino)methane (dppm) or -butane (dppb)) in reactions with $[Rh_5Ru_4(CO)_{16}B]^-$: crystal structures of $[\{Rh_5Ru_4(CO)_{16}B\}_2\{\mu\text{-Au(dppb)Au}\}]$ and $[RhRu_4(CO)_{14}B\{Au(dppm)Au\}]$, **289**, 149

Heterometallic complexes
chelating and ligand-transfer studies of the complexes *cis*-$[Mn(CO)_4(SeR)_2]^-$: crystal structures of heterotrinuclear $[(CO)_4Mn(\mu\text{-SeMe})_2Ni(\mu\text{-SeMe})_2Mn(CO)_4]$ with a distorted square planar Ni^{II}–selenolate core and $[PPN]_2[Cl_2Pd(\mu\text{-SePh})_2PdCl_2]$, **286**, 155

Heteroscorpionate complexes
oxidation state dependent *cis*/*trans* isomerization in ML_2 complexes of the heteroscorpionate ligand, (2-hydroxyphenyl)-bis(pyrazolyl)methane and its derivatives with iron and cobalt, **286**, 80

Hexaborane
$[4\text{-}\{(p\text{-cym})Ru\}\text{-}5\text{-}\{(Ph_3P)_2(CO)Os\}B_4H_8]$: the first *nido*-heterobimetallahexaborane cluster, **289**, 134

Hexamine complexes
molecular recognition of dipeptides. Catalysis of deuteration and hydrolysis of glycylglycine by dinuclear OBISDIEN Zn(II) complexes, **287**, 134

Hexanuclear complexes
electrochemical synthesis of copper and silver pyridine-2-selenolato complexes: crystal structure of (6-trimethylsilylpyridine-2-selenolato)silver(I), $[Ag_6(6\text{-}Me_3SipySe)_6]$, **284**, 133

High coordination complexes
synthesis, characterization, and structural analyses of three high-coordination tetra-n-butylammonium lanthanide(III) complexes, **292**, 137

High pressure NMR
high pressure NMR studies on ligand exchange reactions in the equatorial sites of $[VO(L)_5]^{2+}$ (L = acetonitrile, *N,N*-dimethylformamide, and *N,N*-dimethylacetamide), **294**, 119

4*H*-Indenyl complexes
photochemical reactions of transition metal organyl complexes with olefins: XXII. Photoreactions of pentacarbonyl(μ-$\eta^{5:3}$-4*H*-indenyl)diiron(Fe–Fe) with alkynes—substituent controlled C–H transfer or C–C bond formation, **296**, 127

Histidine complexes
potentiometric and spectroscopic studies of reactions of histidine with 13-membered amide-based macrocyclic Cu^{2+} chelates, **295**, 115

H-resorcinol
mechanistic study on the reaction of boronic acids with H-resorcinol, **290**, 222

Hydration
nitrite ion-promoted hydration of acetonitrile by the bis(μ-hydroxo)bis{tris(2-pyridylmethyl)amine)chromium(III)} ion, **288**, 211

Hydrazide complexes
vanadium phenoxide complexes with oxide, nitride or hydrazide co-ligands: preparation and crystal structures of $[V(OC_6H_3Pr^i_2\text{-}2,6)_3NLi(C_4H_8O)_3]$, $[\{VO_2(OC_6H_3Pr^i_2\text{-}2,6)_2\}_2\{\mu\text{-}Li(C_4H_8O)_2\}_2]$ and $[V(NNMe_2)(OC_6H_3Pr^i_2\text{-}2,6)_3]$, **285**, 178

Hydrazine complexes
a dinuclear platinum(II) complex with hydrazine and chloro bridges: $(NPr^n_4)[PtCl_2(\mu\text{-}Cl)(\mu\text{-}N_2H_4)PtCl_2]$, **284**, 103

Hydrazones
nickel(II) and palladium(II) complexes with acylhydrazone ligands of α-diketones: the electronic and steric factors affecting the formation of the dimeric palladium(II) complexes, **295**, 171
synthesis of transition metal isocyanide complexes containing hydrogen bonding sites in peripheral locations, **285**, 223

Hydride complexes
new synthesis of $[RuH(CO)(PPh_3)_2L_2][BF_4]$ [L_2 = 2py, 2(4-Mepy) or bpy] and X-ray crystal structure of $[RuH(CO)(PPh_3)_2(4\text{-Mepy})_2][BF_4]$, **285**, 318
soluble complexes of $[FeH_6]^{4-}$, **291**, 142
structural, spectroscopic and catalytic properties of water-soluble hydride rhodium complexes $[RhH(Rtpa^+I^-)_4]H_2O$ (R = Me, Et), **293**, 110

Hydroboration
synthesis, reactivity and catalytic properties of rhodium complexes of (*R,R*)-1-benzyl-3,4-dithioetherpyrrolidines, **295**, 64

Hydroformylation
synthesis of chiral diphosphite ligands and comparison of their rhodium and platinum complexes in the asymmetric hydroformylation of styrene, **296**, 222
synthesis, reactivity and catalytic properties of rhodium complexes of (*R,R*)-1-benzyl-3,4-dithioetherpyrrolidines, **295**, 64

Hydrogenation
synthesis, reactivity and catalytic properties of rhodium complexes of (*R,R*)-1-benzyl-3,4-dithioetherpyrrolidines, **295**, 64

Hydrogen bond
synthesis, molecular and crystal structure of the zwitterionic Re(V)–oxo complex $ReOCl_3(L)(PPh_3)$ with L = 2-(diethylaminomethyl)-4-methylphenol, **295**, 209

Hydrogen bonding
effect of N–H···S amide–thiolate hydrogen bonding on bond lengths in rubredoxin models $[(CH_3)_3NCH_2CONH_2]_2[M(S_2\text{-}o\text{-xyl})_2]$, **291**, 388
hydrogen bonding in *cis*-bis(L-alaninato)copper(II): a single crystal neutron diffraction study, **288**, 29

Hydrogen bonding interactions
synthetic control of intrinsic ground-state defects in a mixed valence quasi-one-dimensional Pt halide chain, **294**, 275

Hydrogen peroxide
kinetics of the molybdate and tungstate catalyzed oxidation of iodide by hydrogen peroxide, **288**, 229
Hydrogen transfer
soluble complexes of $[FeH_6]^{4-}$, **291**, 142
Hydrolysis
anionopentaaminecobalt(III) complexes with polyamine ligands: Part 30. The base hydrolysis kinetics and structure determination of some $[CoCl(N)_5]^{2+}$ complexes containing pyridine ligands, **288**, 181
catalysis of hydrolysis of a phosphate diester by mono- and dinuclear macrocyclic zinc(II) complexes, **287**, 47
molecular recognition of dipeptides. Catalysis of deuteration and hydrolysis of glycylglycine by dinuclear OBISDIEN Zn(II) complexes, **287**, 134
synthesis and comparative reactivity of thiomorpholine–borane: aqueous hydrolysis and oxidation by hypochlorite, **290**, 80
the kinetics of the sequential hydrolysis reactions of the *xxm*-$[CoCl_2(dien)(py)]^+$ cation and the crystal structure of the $ZnCl_4^{2-}\cdot 2H_2O$ salt, **292**, 198
the nitrido complex $(NEt_4)[MoCl_3(N)(OH)(MeCN)]$, **295**, 249
Hydrometallation
reactions of cobaloximes with vinyl allenes as a route to 1-cobaloxime substituted 1,3-dienyl complexes, **296**, 13
Hydrophosphoranes
platinum(II) and palladium(II) complexes with 3,3,8,8-tetramethyl-1,6-dioxa-4,9-diaza-5λ^5-phosphaspiro[4,4]nonane, **295**, 164
Hydrosilane
synthesis of 2-silylmethyl-2-alkenals by rhodium-catalyzed interaction among 2-propynylamine, hydrosilane, and CO, **296**, 72
Hydrothermal assembly
hydrothermal assembly and structural characterization of two low-dimensional molybdenum oxide/2,2'-dipyridylamine hybrid materials, **295**, 222
Hydrothermal synthesis
hydrothermal synthesis and characterization of a new vanadyl sulfate compound: crystal structure of $[V(IV)OSO_4(H_2O)_4]\cdot SO_4\cdot[H_2N(C_2H_4)_2NH_2]$, **292**, 112
hydrothermal synthesis and crystal structure of a new molybdenum(VI) arsenate(III), $Co^{III}(en)_3H_3O[(Co^{II}O_6)Mo^{VI}{}_6O_{18}(As^{III}{}_3O_3)_2]\cdot 2H_2O$, **295**, 244
the hydrothermal synthesis and structural characterization of a new class of compounds: nickel organoamine-halocadmates, **292**, 41
Hydrotris(pyrazolyl)borate complexes
synthetic routes for hydrotris(pyrazolyl)borate complexes of rhodium(III), **284**, 191
Hydroxo-bridged complexes
reactivity towards some weak protic electrophiles of the di-μ-hydroxo complex $[\{Pd(\mu\text{-}OH)(tmeda)\}_2][BPh_4]_2$: crystal structure of $[\{Pd(tmeda)\}_2(\mu\text{-}OH)(\mu\text{-}HNC_6H_4Cl\text{-}p)][BPh_4]_2$, **287**, 37
structure and properties of a novel OH^- bridged dimetal helical complex with 6,6''''-dimethyl-2,2':6',2'':6'',2''':6''',2'''':6'''',2''''-quinquepyridine ligand, **290**, 127
synthesis, spectroscopic, magnetic properties and X-ray crystal structure of di-μ-hydroxo-bis(μ-perchlorato-*O,O'*)-bis-[bis(2-amino-4-methylpyrimidine)copper(II)]: bridging perchlorate results in significant deviation from the predicted magnetic exchange, **286**, 24
the molecular and electronic structure of the asymmetric, trinuclear complex $[LFe^{III}(\mu\text{-}O)(\mu\text{-}piv)_2Fe^{III}(\mu\text{-}OH)(\mu\text{-}piv)_2Fe^{III}(piv)_2]\cdot C_6H_5CH_3$ (L = 1,4,7-trimethyl-1,4,7-triazacyclononane, piv = pivalate(1-)), **291**, 258
Hydroxo bridging complexes
synthesis, structure and magnetic properties of a linear trimanganese(II,III,II) complex with hydroxo and 1,2,4-triazole as bridging ligands, **295**, 229
Hydroxylamine derivatives
iminoacylation: 2. Addition of alkylated hydroxylamines via oxygen to platinum(IV)-bound nitriles, **292**, 272
Hydroxylated acid complexes
three mononuclear copper complexes of 9-hydroxy-9*H*-fluorene-9-carboxylic acid, **294**, 163
Hydroxymethyl-phosphine complexes
a molybdenum(0) complex of bis(hydroxymethyl)phenylphosphine, **287**, 209
Hydroxypyridone complexes
halfsandwich complexes of ruthenium(II), rhodium(III) and iridium(III) with *N*-substituted 3-hydroxy-2-methyl-4-pyridone ligands, **294**, 62
Hypervalent complexes
synthesis and structures of 2-$[Me_2NCH_2CH_2N(Me)CH_2]C_6H_4$ substituted tin compounds, **290**, 100

Imidazoles
crystal structures of two new heptamolybdates and of a pyrazole incorporating a γ-octamolybdate anion, **295**, 106
steric effect on the axial substitution of bis(dimethyl sulphoxide)phthalocyaninatoiron(II) by substituted imidazoles, **295**, 200
Imidazole basicity
5,10,15,20 - tetrakis(α,α,α,α - *o* - pivalamidophenyl)porphinatoiron(II) bearing a covalently linked axial imidazole via *m*-aminobenzoic acid. Synthesis and influence of imidazole basicity on O_2-binding affinity, **295**, 234
Imidazole complexes
copper(I) and copper(II) complexes of the bidentate imidazole/thioether ligand 1-methyl-2-(methylthiomethyl)-1*H*-benzimidazole, **287**, 204
cyanamidonitrate–copper(II) complexes of imidazole ligands: X-ray crystallography and physical investigation, **287**, 186
electronic spectra of 1,2-diiminetricarbonylrhenium(I)chloride complexes with imidazole derivatives as ligands, **288**, 150
Imidazolyl-borate complexes
new phosphino silver(I) derivatives of hydrotris(3-methyl-2-thioxo-1-imidazolyl)borate. X-ray crystal structure of tricyclohexylphosphinesilver(I) hydrotris(3-methyl-2-thioxo-1-imidazolyl)borate, **285**, 81
Imido-bridged complexes
spontaneous formation of trinickel clusters containing μ^3-imido bridges from 2-nitrophenols and $Ni(PMe_3)_4$, **287**, 199
Imidocarbyne bridged complexes
reaction patterns of 'tetraferrio-azaallenium', $[(OC)_3Cp_2Fe_2(\mu_4\text{-}C{=}N{=}C)Fe_2Cp_2(CO)_3]^+$, and its 'activated cyanide'-precursor (μ-phthalimidocarbyne)tricarbonyldicyclopentadienyldiiron(1+), **291**, 266
Imido complexes
synthesis and structure of a novel terminal imido complex of titanium, **292**, 105
Iminoacylation
iminoacylation: 2. Addition of alkylated hydroxylamines via oxygen to platinum(IV)-bound nitriles, **292**, 272
Iminodiacetic acid complexes
solution studies of complexes of iron(III) with iminodiacetic, alkyl-substituted iminodiacetic and nitrilotriacetic acids by potentiometry and cyclic voltammetry, **291**, 158
Indenyl complexes
monoindenyl titanium alkyl halides. The synthesis and molecular structures of $(\eta^5\text{-}C_9H_7)TiBr_3$, $(\eta^5\text{-}C_9H_7)Ti(CH_3)Br_2$, and $(\eta^5\text{-}C_9H_7)Ti(CH_3)Cl_2$, **292**, 220

synthesis of (ferrocenyl-indenyl)cyclopentadienyliron compounds with and without a bridging group via a CpFe transfer reaction, **286**, 215

Indium amine
synthesis of new Group III fluoride–ammonia adducts in supercritical ammonia: structures of $AlF_3(NH_3)_2$ and $InF_2(NH_2)(NH_3)$, **294**, 200

Indium complexes
indate(III) phthalocyanines: synthesis, spectroscopy and crystal structure of iodophthalocyaninato(2-) indate(III), **288**, 174

Indium compound
electrochemical tuning of indium phosphide|poly(acetylene) interfaces, **294**, 207

Indium(III) halide complexes
main group metal halide complexes with sterically hindered thioureas XVI: The synthesis, characterization, and crystal structures of two new complexes of 1,3-dimethyl-2(3*H*)-imidazolethione with indium trihalides, **285**, 217

Infrared spectra
lattice vibrations of the icosahedral solid boron arsenide, $B_{12}As_2$, **289**, 198

Inner-sphere adducts
high coordination numbers of metal ions in chelate complexes: molecular adducts of scandium tris-α-diketonates in solution, **286**, 175

Intervalence transfer
intervalence transfer properties of the binuclear μ-benzotriazolate- and μ-benzimidazolate-bis{ruthenium(II)/(III)-edta} complexes, **285**, 197

Intramolecular charge transfer
properties of Pt(II) complexes containing both a pyridyl N–N chelate ligand having a long alkyl chain and a sulfur-rich dithiolate ligand and their molecular interactions in the solid state, **284**, 55

Intramolecular coupling
intramolecular coupling of conjugated enynes with cyclopropylcarbene–chromium complexes; a complex reaction pathway, **296**, 273

Intramolecular phosphorous coordination
synthesis of the first chiral phosphorous chelate aminophosphino–carbene complex and its applications in stereoselective addition reactions, **296**, 236

Iodide complexes
structural and spectroscopic characterisation of the mixed valence platinum complex $PtdabI_3$ (dab = *o*-diaminobenzene), **287**, 218

Iodide oxidation
kinetics of the molybdate and tungstate catalyzed oxidation of iodide by hydrogen peroxide, **288**, 229

Iodine dimerization
a neutral molecule of I_2 as a bridge for dimerization of iodine iron(III) phthalocyanine, **287**, 101

Iodocarbon complexes
silver(I) tetraiodoethylene complexes with twisted olefin moiety, **290**, 251

Iridium
an approach to *megalo*-boranes. Mixed and multiple cluster fusions involving iridaborane and platinaborane cluster compounds. Crystal structure determinations by conventional and synchrotron methods, **289**, 95
application of the NOE experiment to the analysis of boron hydride derivatives: confirming the assignments of the *pseudocloso*-complex [1,2-Ph_2-3-{Cp*}-3,1,2-$IrC_2B_9H_9$] (Cp* = η^5-C_5Me_5) and the *closo*-compounds 1-Ph-1,2-$C_2B_{10}H_{11}$ and 1-Ph-2-Me-1,2-$C_2B_{10}H_{10}$, **289**, 125

Iridium complexes
B-frame-supported bimetallics. Isoelectronic *arachno*-structured $[(PMe_2Ph)_4Pd_2B_2H_{10}]$ and *closo*-structured $[(PMe_3)_4(CO)_2Ir_2B_8H_8]$, **285**, 290
halfsandwich complexes of ruthenium(II), rhodium(III) and iridium(III) with *N*-substituted 3-hydroxy-2-methyl-4-pyridone ligands, **294**, 62
preparation, structure and electrochemical behavior of dinuclear cyclooctadiene-chelated Ir(I) complexes with 2-aminopyridinato bridges, **292**, 244
reaction of $M(CO)Cl(L)_2$ with mercury(II) chloride (M = Ir, Rh; L = Ph_2Ppy, P(fur)$_3$) (Ph_2Ppy = 2-(diphenylphospino)pyridine, P(fur)$_3$ = tri-(2-furyl)phosphine), **284**, 119
synthesis, X-ray crystal structure, and reactivity of the monomeric dithio-*o*-carboranyl iridium complex $[Ir(\eta^5\text{-}C_5Me_5)(\eta^2\text{-}S_2C_2B_{10}H_{10})]$, **289**, 141
the ruthenium and iridium coordination chemistry of the tripodal ligand $CH_3C\{CH_2P(m\text{-}CF_3\text{-}C_6H_4)_2\}_3$, **293**, 70

Iron clusters
relativistic density functional study on the bimetallic cluster $[Pt_3Fe_3(CO)_{15}]^{n-}$ ($n = 0, 1, 2$), **286**, 30

Iron complexes
about the synthesis of the prismane $[NEt_4]_2[Fe_6S_6I_6]$, **284**, 296
aminophosphanes derived from chiral β-aminoalcohols and carbonyliron complexes thereof, **291**, 300
a neutral molecule of I_2 as a bridge for dimerization of iodine iron(III) phthalocyanine, **287**, 101
a new linear polymer chain iron complex; synthesis, crystal and molecular structural characterization of catena-μ-acetato-[*NN'*-ethylenebis(salicylaldiminato)]iron(III), **287**, 109
a polyoxoanion-based iron(II) catechol extradiol dioxygenase: oxygenation of 3,5-di-tert-butylcatechol by the polyoxoanion-supported iron(II) complex $(n\text{-}Bu_4N)_7[(CH_3CN)_3\text{-}Fe^{II}\text{-}P_2W_{15}Nb_3O_{62}]$, **291**, 426
a reversible NO complex of [Fe^{II}(edampda)] and the [Ni^{II}(edampda)] analogue (edampda^{2-} = *N,N'*-bis(pyridylmethyl)ethylenediamine-*N,N'*-diacetate), **286**, 197
bisphenolate iron(II) complexes with intramolecularly coordinating nitrogen Lewis bases, **291**, 326
carbosilane dendrimers functionalized with $AuFe_3$ clusters, **291**, 247
chemoselective synthesis of (allyloxy)carbene chelate complexes of iron, **291**, 449
comparing structures and reactivity in analogous Fe and Ru complexes. (^{i}Pr-DAB)Fe(CO)$_2$I$_2$ and (^{i}Pr-DAB)FeI$_2$: a perfectly reversible CO-carrier system. (R-DAB = *N,N'*-R_2-1,4-diaza-1,3-butadiene), **291**, 438
conformational diversity in (octaethylporphinato)(trichloroacetato)iron(III) derivatives, **291**, 49
crystal structure of (μ-oxo)bis{aquatris(2-pyridylmethyl)amine}iron(III) perchlorate, **284**, 124
C–S, C–H, and N–H bond cleavage of heterocycles by a zero-valent iron complex, Fe(N_2)(depe)$_2$ [depe = 1,2-bis(diethylphosphino)ethane], **291**, 341
cyano-bridged dinuclear iron complexes in different oxidation states, **287**, 95
formation of donor–acceptor Fe(0)–Hg(II) bond in separation and stabilization of optically active iron(0) phosphine complexes. Absolute configuration of (+)-(*R*)-(CO)$_4$Fe(μ-EtPhPpy)$HgCl_2$, **293**, 106
iron and ruthenium organometallic derivatives of spiro-substituted cyclotriphosphazenes bearing nitrile spacer: synthesis and electronic structure, **294**, 20
iron carbonyls with bulky thiolate ligands: crystal structures of $[Fe_2(CO)_6(\mu\text{-}SC_6H_2{}^iPr_3\text{-}2,4,6)_2]$ and $(C_6H_2{}^iPr_3\text{-}2,4,6)_2S_2$, **284**, 14
iron complexes of dioxoborolanylbutadienes, **290**, 159

iron(II/III) complexes of non-porphyrin macrocyclic $[N_4]^{2-}$ ligands: structure and axial reactivity, **291**, 190
kinetics of the photooxidation of tris(bipyridine)iron(II) in chloroform, **291**, 136
lattice effects in the Mössbauer spectra of salts of $[Fe_4S_4\{S(CH_2)_nOH\}_4]^{2-}$. Crystal structures of $[PPh_4]_2$-$[Fe_4S_4\{S(CH_2)_nOH\}_4]$ ($n = 2$, 3 and 4), **291**, 101
low-spin iron(III) complexes with N,S coordination: syntheses, structures, and properties of bis(*N*-2-mercaptophenyl-2′-pyridylmethyleniminato)iron(III) tetraphenylborate and bis(*N*-2-mercapto-2-methylpropyl-2′-pyridylmethylenimimato)iron(III) tetraphenylborate, **285**, 269
macrocyclic complexes derived from the cyclocondensation of sodium 2,6-diformyl-4-substituted phenolates with *N*,*N*-bis(2-aminoethyl)-2-hydroxybenzylamine and in situ transmetallation. An anticipated dinuclear Cd(II) pendant-arm macrocyclic complex and an unexpected mononuclear Fe(III) macrocyclic complex, **294**, 1
magnetic iron phosphates with an open framework, **291**, 317
mixed-valence states of iron long-chain carboxylate complexes, **291**, 127
molecular orbital-theoretical studies of a dinuclear Fe(II)–Fe(III) complex, **291**, 82
new capping agents for oximehydrazonate clathrochelates: sterically controlled synthesis, structural characterization and intramolecular reactions, **284**, 180
new ferracyclic π-allyl and η^3-pentadienyl complexes by carbonylative lactonization of unsaturated diols, **291**, 91
novel features associated with the electrochemically driven bis(η^5-pentaphenylcyclopentadienyl)iron(II)–iron(III) redox transformation at an electrode–microcrystal–solvent (electrolyte) interface, **291**, 21
novel iron(III) complexes with imidazole containing tripodal ligands as model systems for catechol dioxygenases, **291**, 289
nucleophilic substitutions of iodide with bromides in the $Fe(CO)_2(PMe_3)_2MeI$ complex, **288**, 200
on the contrasting reactivity of the phosphaalkene $[(\eta^5\text{-}C_5Me_5)(CO)_2FeP{=}C(NMe_2)_2]$ and the arsaalkene $[(\eta^5\text{-}C_5Me_5)(CO)_2FeAs{=}C(NMe_2)_2]$ towards $[(\eta^5\text{-}C_5H_5)Rh(CO)_2]$, **291**, 311
organometallic electron reservoir sandwich iron complexes as potential agents for redox and electron transfer chain catalysis, **291**, 1
oxidation state dependent *cis*/*trans* isomerization in ML_2 complexes of the heteroscorpionate ligand, (2-hydroxyphenyl)-bis(pyrazolyl)methane and its derivatives with iron and cobalt, **286**, 80
photo-assisted preparation of double isocyanide insertion complexes of η^5-cyclopentadienyliron(II) bearing only isocyanide: the molecular structure of $(\eta^5\text{-}C_5H_5)Fe(XylNC)_2$-$[C(=NXyl)C(=NXyl)CH_3]$ (XylNC = 2,6-$Me_2C_6H_3NC$), **286**, 233
potentiometric determination of the equilibrium constants for the formation of Fe(III)–sorbitol complexes, **293**, 206
protonation and substitution reactions of Fe–S 'basket' clusters including extracted FeMo-cofactor of nitrogenase, **291**, 212
reaction patterns of 'tetraferrio-azaallenium', $[(OC)_3Cp_2Fe_2(\mu_4\text{-}C{=}N{=}C)Fe_2Cp_2(CO)_3]^+$, and its 'activated cyanide'-precursor (μ-phthalimidocarbyne)tricarbonyldicyclopentadienyldiiron(1 +), **291**, 266
reactions of σ-π alkenyl complexes $[Fe_2(CO)_6(\mu\text{-alkenyl})(\mu\text{-}PPh_2)]$ with bis(diphenylphosphino)methane (dppm): carbonyl substitution, migratory carbonyl insertion, phosphorus–carbon bond formation and cleavage, proton shift and alkenyl isomerisation reactions, **291**, 178
receptors based on 2,2′:6′,2″-terpyridine fragments containing peripheral amino groups, **292**, 28
redox active organoiron alkynyl centers with a coordination site: an approach toward molecular plugs. Complexation reactions with $(THF)W(CO)_5$ and $(PhCN)_2MCl_2$ (M = Pd, Pt), **291**, 403
reduction of cationic P-coordinated (diphenylphosphino)alkyne iron complexes with sodium borohydride, **291**, 207
soluble complexes of $[FeH_6]^{4-}$, **291**, 142
solution studies of complexes of iron(III) with iminodiacetic, alkyl-substituted iminodiacetic and nitrilotriacetic acids by potentiometry and cyclic voltammetry, **291**, 158
solution thermochemical study of ligand substitution reactions of hybrid alkyl/fluoroalkoxy phosphorus ligands in the $L_2Fe(CO)_3$ system, **291**, 32
spin-crossover in the $[Fe(abpt)_2(NCX)_2]$ (X = S, Se) system: structural, magnetic, calorimetric and photomagnetic studies, **291**, 279
structure and characterization of a mononuclear Fe^{II} dialkyldithiocarbamate complex $Fe(OC_4H_9dtc)_2(DMF)$, **293**, 95
structure and magnetic properties of (μ-oxo)bis[*meso*-tetrakis(*p*-tolyl-porphyrinato)iron(III)] and (μ-oxo)bis[*N*,*N*′-ethylene-bis(2-acetylphenoliminato)iron(III)], **290**, 51
syntheses, properties and Mössbauer studies of cyanamide and cyanoguanidine complexes of iron(II). Crystal structures of *trans*-$[FeH(NCNH_2)(Ph_2PCH_2CH_2PPh_2)_2][BF_4]$ and *trans*-$[Fe(NCNEt_2)_2(Et_2PCH_2CH_2PEt_2)_2][BF_4]_2$, **291**, 39
synthesis and characterization of a ferric complex of the tripodal ligand tris(2-benzimidazolylmethyl)amine—a superoxide dismutase mimic, **290**, 21
synthesis and characterization of long chain acyl and alkyl complexes of the types $[(\eta^5\text{-}C_5R'_5)Fe(CO)_2\{C(O)R\}]$ (R′ = H, CH_3), $[CpW(CO)_3\{C(O)R\}]$ and $[CpW(CO)_3R]$ (R = *n*-alkyl), **286**, 221
synthesis and structural characterization of $[NEt_4][Fe_3(\mu_3\text{-}O)(\mu_3\text{-}AuPPh_3)(\mu\text{-}CO)_3(CO)_6]$, the new $[Au_6(\mu_3\text{-}S)_2(PPh_3)_6][Fe_3(\mu_3\text{-}S)(\mu\text{-}AuPPh_3)(CO)_9]_2$ and $[Au_6(\mu_3\text{-}S)_2(PPh_3)_6][Fe_4(\mu_3\text{-}S)_2\text{-}(CO)_{14}]$ ionic solids containing assemblages of clustercations and cluster-anions, **291**, 372
synthesis of bis-(acyl, carbamoyl or alkoxycarbonyl) iron monomer or dimer complexes by reaction of oxalyl chloride with various tetracarbonyl ferrates: $\{Fe[C(O)R](CO)_4\}^-$, **291**, 166
synthesis of (ferrocenyl-indenyl)cyclopentadienyliron compounds with and without a bridging group via a CpFe transfer reaction, **286**, 215
synthesis of hexametallic $CH_2\{(\eta^5\text{-}C_5H_4)Fe(CO)_2SnPh_2(\eta^5\text{-}C_5H_5)Fe(CO)_2\}_2$, **291**, 252
synthesis of transition metal isocyanide complexes containing hydrogen bonding sites in peripheral locations, **285**, 223
synthesis, spectroscopic, redox properties, and DNA cleavage activity of low-spin iron(III) complexes of bleomycin analogues, **294**, 56
synthesis, structure, and redox properties of the sulfido bridged tetranuclear cluster complex $[(Ph_3P)_2N][Fe_3(CO)_9\{\mu_3\text{-}SFeCp(CO)_2\}]$, **286**, 142
synthesis, studies and molecular structure of trinuclear cyanide-bridged copper–iron complexes, **292**, 157
synthetic approaches to Fe(II) complexes of the bulky phosphine P^tBu_2Me, **291**, 226
the molecular and electronic structure of the asymmetric, trinuclear complex $[LFe^{III}(\mu\text{-}O)(\mu\text{-piv})_2Fe^{III}(\mu\text{-}OH)(\mu\text{-piv})_2Fe^{III}(piv)_2]\cdot C_6H_5CH_3$ (L = 1,4,7-trimethyl-1,4,7-triazacyclononane, piv = pivalate-(1-)), **291**, 258
the relationship between surface derivatization and the physical properties of bisarylphosphinidenetriironnonacarbonyl clusters, **294**, 140

thermodynamic and kinetic studies of the sulfonated derivative of the iron chelator TRENCAM, an analog of enterobactin, **291**, 148
the μ-sulfonium–methylidene diiron complexes $[Fe_2\{\mu\text{-}C(X)SMe_2\}(\mu\text{-}CO)(CO)_2(Cp)_2]SO_3CF_3$ (X = CN, H) as precursors of μ-alkylidene complexes, **291**, 333
transformation of the carboxyl group adjacent to the $(\eta^4$-diene)$Fe(CO)_3$ moiety: a convenient route to $(\eta^4$-diene)$Fe(CO)_3$ complexes containing α,β- and β,γ-unsaturated carbonyl groups, and 1,3-dicarbonyl groups, **291**, 231
visible and resonance Raman spectra of low valent iron porphyrins, **287**, 21
widely differing stabilities of molecule-bridged cyanodiiron(III, II) species in non-aqueous solvents, **291**, 66

Iron(II) complexes
5,10,15,20-tetrakis(α,α,α,α-*o*-pivalamidophenyl)porphinatoiron(II) bearing a covalently linked axial imidazole via *m*-aminobenzoic acid. Synthesis and influence of imidazole basicity on O_2-binding affinity, **295**, 234

Iron(II) phthalocyanine
steric effect on the axial substitution of bis(dimethyl sulphoxide)phthalocyaninatoiron(II) by substituted imidazoles, **295**, 200

Iron overload
new chelating agents suitable for the treatment of iron overload, **291**, 238

Iron silicides
formation of FeSi and $FeSi_2$ films from *cis*-$Fe(SiCl_3)_2(CO)_4$ by MOCVD –precursor versus substrate control, **291**, 380

Isocyanide complexes
isocyanide derivatives of manganese cobalt nonacarbonyl: synthesis and characterization, **288**, 159
photo-assisted preparation of double isocyanide insertion complexes of η^5-cyclopentadienyliron(II) bearing only isocyanide: the molecular structure of $(\eta^5\text{-}C_5H_5)Fe(XylNC)_2[C(=NXyl)C(=NXyl)CH_3$ (XylNC = 2,6-$Me_2C_6H_3NC$), **286**, 233
synthesis and formation of metal complexes of 4-alkynyl and 4-cyano-2,6-diisopropylphenylisocyanides, **284**, 205

Isocyanide metal complexes
synthesis of transition metal isocyanide complexes containing hydrogen bonding sites in peripheral locations, **285**, 223

Isocyanides
synthesis of transition metal isocyanide complexes containing hydrogen bonding sites in peripheral locations, **285**, 223

Isoxazolines
synthesis and reactivity of acyclic (pentadienyl)iron(1 +) cations: model studies for the preparation of the 8*E*,10*Z*,16*E*,18*E*-tetraene segment of macrolactin A, **296**, 261

Jahn–Teller distortion
a new linear polymer chain iron complex; synthesis, crystal and molecular structural characterization of catena-μ-acetato-[*NN'*-ethylenebis(salicylaldiminato)]iron(III), **287**, 109
structure and physical properties of hexacyanomanganate(III), $[Mn^{III}(CN)_6]^{3-}$, **284**, 175

Kinetics
effects of β-cyclodextrin inclusion on the kinetics of the oxidation of bisferrocenyl cations by bis(pyridine-2,6-dicarboxylato)cobaltate(III) in aqueous solution: crystal structure of dimethyldi-(1-methylferrocene)ammonium bromide, **291**, 365

Kinetics and mechanism
anionopentaaminecobalt(III) complexes with polyamine ligands: Part 30. The base hydrolysis kinetics and structure determination of some $[CoCl(N)_5]^{2+}$ complexes containing pyridine ligands, **288**, 181
kinetics and mechanism of the reduction of ferrate by one-electron reductants, **293**, 229
kinetics of the photooxidation of tris(bipyridine)iron(II) in chloroform, **291**, 136
ligand substitution kinetics in $M(CO)_4$ ($\eta^{2\text{-}2}$-1,5-cyclooctadiene) complexes (M = Cr, Mo, W) — substitution of 1,5-cyclooctadiene by bis(diphenylphosphino)alkanes, **284**, 229
mechanistic study on the reaction of boronic acids with H-resorcinol, **290**, 222
$[(NH_3)_5CoOP(H)(O)OCo(NH_3)_5]^{4+}$ and its conjugate acid; evidence for the $P^+\text{–}O^-$ description of a terminal P–O bond, **294**, 14
nucleophilic substitutions of iodide with bromides in the $Fe(CO)_2(PMe_3)_2MeI$ complex, **288**, 200
palladium(II) allyl complexes with potentially terdentate ancillary ligands. Mechanism of allyl amination by piperidine, **293**, 44
protonation and substitution reactions of Fe–S 'basket' clusters including extracted FeMo-cofactor of nitrogenase, **291**, 212
the kinetics of the sequential hydrolysis reactions of the *xxm*-$[CoCl_2(dien)(py)]^+$ cation and the crystal structure of the $ZnCl_4^{2-}\cdot 2H_2O$ salt, **292**, 198
thermodynamic and kinetic studies of the sulfonated derivative of the iron chelator TRENCAM, an analog of enterobactin, **291**, 148

Lactam complexes
platina- and palladalactam complexes derived from 2-benzoylacetanilide; syntheses and X-ray structure of $[Pd\{NPhC(O)CHC(O)Ph\}(bipy)]\cdot CH_2Cl_2$, **292**, 260

Lanthanide complexes
asymmetric compartmental macrocyclic ligands and related mononuclear and hetero-dinuclear complexes with d- and/or f-metal ions, **293**, 178
crystal structures and stereochemical properties of lanthanide(III) complexes with ethylenediamine-*N,N,N',N'*-tetraacetate, **288**, 7
mononuclear lanthanide complexes of tripodal ligands: synthesis and spectroscopic studies, **285**, 49
spectroscopic study of lanthanide(III) complexes with carboxylic acids, **293**, 167
syntheses, crystal structures and magnetic properties of heterometallic copper–lanthanide clusters $[Cu_{12}Ln_6(\mu_3\text{-}OH)_{24}(\mu\text{-}O_2CR)_{12}(H_2O)_{18}(\mu_{12}\text{-}ClO_4)]^{5+}$ (Ln = La, Nd; R = CH_2Cl, CCl_3), **293**, 129
synthesis, characterization, and structural analyses of three high-coordination tetra-n-butylammonium lanthanide(III) complexes, **292**, 137
thermodynamic and structural study of inclusion complexes between trivalent lanthanide ions and native cyclodextrins, **293**, 53

Lanthanide dithionates
complexes of lanthanide dithionates 2. Electrospray mass spectrometry of $Ln_2(S_2O_6)_3$ and their Ph_3PO complexes; the X-ray crystal structure of $Pr_2(S_2O_6)_3(Ph_3PO)_6(H_2O)_6$, **295**, 146

L-Arginine complexes
aromatic carboxylate-controlled self-organization of copper(II)–L-arginine complexes, **286**, 37

Laser photolysis
photooxidation of water and isobutane in aqueous uranyl solutions, **285**, 322

Lead complexes
18-membered tetraimino macrocyclic complexes containing furan groups: stacking of two macrocycles and characterization of organic derivatives, **284**, 149

Lead(II) salts
Barbier-type allylation of carbonyl derivatives by use of aluminum as an electron pool. Double allylation of carboxylic esters, **296**, 204
Lewis acids
synthetic reactions using organometallics in water. Aldol and allylation reactions catalyzed by Lewis acid–surfactant-combined catalysts/Brønsted acids systems, **296**, 158
Ligand bites
high coordination numbers of metal ions in chelate complexes: molecular adducts of scandium tris-α-diketonates in solution, **286**, 175
Ligand substitution
iron(II/III) complexes of non-porphyrin macrocyclic $[N_4]^{2-}$ ligands: structure and axial reactivity, **291**, 190
ligand substitution kinetics in $M(CO)_4$ ($\eta^{2,2}$-1,5-cyclooctadiene) complexes (M = Cr, Mo, W) — substitution of 1,5-cyclooctadiene by bis(diphenylphosphino)alkanes, **284**, 229
Linear chain complexes
synthesis, crystal structure and magnetic properties of a copper(II)–nitroxide linear chain complex, **287**, 181
Linear polymer chain complexes
a new linear polymer chain iron complex; synthesis, crystal and molecular structural characterization of catena-μ-acetato-[*NN'*-ethylenebis(salicylaldiminato)]iron(III), **287**, 109
Linkage isomerism
synthesis, structures, and linkage isomerism of (allylbenzylmalonate)platinum(II) complexes, **295**, 214
Lipophilicity
synthesis and properties of hydrophobic $[Mn^{IV}_2(\mu\text{-}O)_3(L)_2]^{2+}$ complexes, derived from alkyl substituted 1,4,7-triazacyclononane ligands, **295**, 189
Lithium-bis(diphenylphosphanyl)amide
chemistry of polyfunctional molecules : CXXXI. Crystal structure and preparation of [bis(diphenyl–methyl–phosphino)iminium] tetraphenyloborate, $[(C_6H_5)_2(CH_3)P{=}N{=}P(CH_3)(C_6H_5)_2]^+[B(C_6H_5)_4]^-$, **284**, 288
Low-dimensional materials
hydrothermal assembly and structural characterization of two low-dimensional molybdenum oxide/2,2′-dipyridylamine hybrid materials, **295**, 222
Low valent complexes
visible and resonance Raman spectra of low valent iron porphyrins, **287**, 21
Luminescence
electronic spectra of 1,2-diiminetricarbonylrhenium(I)chloride complexes with imidazole derivatives as ligands, **288**, 150
heteroleptic poly(pyrazol-1-yl)borate derivatives of the lanthanides. The syntheses of picolinate-*N*-oxide complexes and the X-ray crystal structure of $[Tb\{HB(C_3H_3N_2)_3\}_2(ONC_5H_4CO_2\text{-}2)]$, **285**, 283
synthesis, characterization and optical properties of novel carbazole–ruthenium compounds: carbazole as a new building block?, **288**, 134
the crystal structure and doublet emission properties of *trans*-$[Cr(NH_3)_2(cyclam)](ClO_4)Cl_2$, **285**, 313

Macrocycle complexes
iron(II/III) complexes of non-porphyrin macrocyclic $[N_4]^{2-}$ ligands: structure and axial reactivity, **291**, 190
18-membered tetraimino macrocyclic complexes containing furan groups: stacking of two macrocycles and characterization of organic derivatives, **284**, 149
Macrocyclic complexes
^{1}H NMR study of new heterodinuclear macrocyclic complexes, **285**, 233
macrocyclic complexes derived from the cyclocondensation of sodium 2,6-diformyl-4-substituted phenolates with *N,N*-bis(2-aminoethyl)-2-hydroxybenzylamine and in situ transmetallation. An anticipated dinuclear Cd(II) pendant-arm macrocyclic complex and an unexpected mononuclear Fe(III) macrocyclic complex, **294**, 1
synthesis, structure and magnetic behavior of a di-μ-chloro-di-μ-imidazolato-tetracopper(II) complex with two dinucleating macrocycles, **287**, 52
template synthesis, crystal structure, and solution behavior of a hexaaza macrocyclic nickel(II) complex containing two *N*-aminoethyl pendant arms, **293**, 140
the interaction of BMXD and its copper(II) complexes with glycine, aspartic acid, and histidine, **286**, 55
Macrocyclic complexes equilibrium studies
equilibrium studies of reactions of 4-chloro-1,2-diaminobenzene, 4-methyl-1,2-diaminobenzene, 4-methoxy-1,2-diaminobenzene and 4,5-dimethyl-1,2-diaminobenzene with dinuclear cobalt(II) 1,4,7,13,16,19-hexaaza-10,22-dioxacyclotetracosane, **284**, 30
Macrocyclic ligand complexes
catalysis of hydrolysis of a phosphate diester by mono- and dinuclear macrocyclic zinc(II) complexes, **287**, 47
catalytic oxidation of 1,2-diaminobenzene by the dinuclear cobalt(II)–OBISDIEN dioxygen complex, **287**, 193
complexation of macrocyclic ligands with relatively non-solvated metal ions generated in dichloromethane by electrochemical oxidation of amalgam electrodes, **293**, 223
diphenoxo-bridged dinuclear nickel(II) complex of a dinucleating macrocycle with $N(amine)_6O_2$ metal-binding sites: synthesis, X-ray crystal structure and electrochemistry, **287**, 212
potentiometric and spectroscopic studies of reactions of histidine with 13-membered amide-based macrocyclic Cu^{2+} chelates, **295**, 115
the synthesis and structure of some, *rac–trans-*, *meso–trans-* and (*S,S*)-*trans-* Co(III) complexes with Me_6-[14]-diene-(5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene), **292**, 73
Macropolyhedral clusters
an approach to *megalo*-boranes. Mixed and multiple cluster fusions involving iridaborane and platinaborane cluster compounds. Crystal structure determinations by conventional and synchrotron methods, **289**, 95
Magnesium
Grignard reagent formation from aryl halides. There is no aryl radical intermediate along the dominant reaction channel, **296**, 52
Magnetic properties
a new bidimensional compound containing (μ-thiocyanate)(bpy) copper(II) molecules: synthesis, crystal structure and magnetic properties of $[Cu(bpy)(NCS)_2]_n$ (bpy = 2,2′-bipyridyl), **286**, 108
crystal structure and magnetic properties of the single-μ-chloro copper(II) chain $[Cu(bipy)Cl_2]$ (bipy = 2,2′-bipyridine), **284**, 258
crystal structures and magnetic properties of the squarate-O^1,O''-bridged dinuclear copper(II) complexes $[Cu_2(phen)_4(C_4O_4)](CF_3SO_3)_2\cdot 3H_2O$ ($n = 2$) and $[Cu_2(bipy)_4(C_4O_4)](CF_3SO_3)_2\cdot 6H_2O$ ($n = 3$), **287**, 173
infinite two-dimensional sheet-like array of a 2,2′-bipyrimidine-bis(oxalato)chromate(III) complex: synthesis, crystal structure and magnetic properties of $[K(H_2O)Cr(C_2O_4)_2(bpym)]$, **284**, 8
spin-crossover in the $[Fe(abpt)_2(NCX)_2]$ (X = S, Se) system: structural, magnetic, calorimetric and photomagnetic studies, **291**, 279

structure and magnetic properties of dinuclear [Mn(III)(salen)$(H_2O)]_2(ClO_4)_2$ and polynuclear [Mn(III)(salen)$(NO_3)]_n$, **290**, 8

structure and magnetic properties of (μ-oxo)bis[*meso*-tetrakis(*p*-tolyl-porphyrinato)iron(III)] and (μ-oxo)bis[*N,N'*-ethylenebis(2-acetylphenoliminato)iron(III)], **290**, 51

structure and physical properties of hexacyanomanganate(III), $[Mn^{III}(CN)_6]^{3-}$, **284**, 175

synthesis, crystal structure and magnetic properties of *catena*-poly[(2,2'-bipyridylcopper)-di-μ-thiocyanato] and bis(2,2'-bipyridyl)thiocyanatocopper(II) perchlorate, **286**, 169

synthesis, spectroscopic, magnetic properties and X-ray crystal structure of di-μ-hydroxo-bis(μ-perchlorato-*O,O'*)-bis-[bis(2-amino-4-methylpyrimidine)copper(II)]: bridging perchlorate results in significant deviation from the predicted magnetic exchange, **286**, 24

synthesis, structure and magnetic properties of a linear trimanganese(II,III,II) complex with hydroxo and 1,2,4-triazole as bridging ligands, **295**, 229

versatility of the cyanato ligand: structure and magnetic properties of four new copper(II)–cyanato compounds derived from tridentate amines, **286**, 189

Magnetic susceptibility

B and B–C as interstitials in reduced rare earth halides, **289**, 174

Magnetism

an alternating copper(II) chain with bridging oxamidato and nitrito ligands: crystal structure and magnetic properties of $[Cu(NO_2)_2CuL]_n$ (L = *N,N'*-bis(2-methyl-2-aminopropyl) oxamide), **294**, 8

magnetic iron phosphates with an open framework, **291**, 317

magneto–structural correlation of dimeric copper(II) carboxylates with pyridyl-substituted nitronyl nitroxides, **293**, 30

structure and magnetic properties of *catena*-(μ-CrO_4-*O,O'*)$[M^{II}(L)]\cdot 5H_2O$ (M = Ni^{II}, Cu^{II}; L = 3,14-dimethyl-2,6,13,17-tetraazatricyclo[14,4,$0^{1,18}$,$0^{7,12}$]docosane), **293**, 100

syntheses, crystal structures and magnetic properties of heterometallic copper–lanthanide clusters $[Cu_{12}Ln_6(\mu_3\text{-}OH)_{24}(\mu\text{-}O_2CR)_{12}(H_2O)_{18}(\mu_{12}\text{-}ClO_4)]^{5+}$ (Ln = La, Nd; R = CH_2Cl, CCl_3), **293**, 129

synthesis, characterization, spectroscopy, magnetism and X-ray structures of red and green Cu(II) chloride adducts with bis(2-benzimidazolyl)propane, **288**, 220

synthesis, crystal structure and magnetic properties of a copper(II)–nitroxide linear chain complex, **287**, 181

synthesis, crystal structures and magnetic properties of trinuclear oxo-centered manganese complexes of the general formula $[Mn_3O(O_2CR)_6L_3]^{z+}$ (R = Me, Et, Ph, CH_2Cl; L = 3-methylpyridine or water $z = 1$, 0), **287**, 82

Manganese complexes

a series of heteroleptic complexes of the type *fac*-$[Mn^{III}L_2]^-$ [H_2L = derivatives of *N*-(2-hydroxybenzyl)glycine or *N*-(5-nitro-2-hydroxybenzyl)sarcosine] possessing unusual Mn(III) co-ordination spheres, **290**, 228

a seven-coordinate manganese(II) complex formed with the single heptadentate ligand *N,N,N'*-tris(2-pyridylmethyl)-*N'*-(2-salicylideneethyl)ethane-1,2-diamine, **290**, 139

chelating and ligand-transfer studies of the complexes *cis*-$[Mn(CO)_4(SeR)_2]^-$: crystal structures of heterotrinuclear $[(CO)_4Mn(\mu\text{-}SeMe)_2Ni(\mu\text{-}SeMe)_2Mn(CO)_4]$ with a distorted square planar Ni^{II}–selenolate core and $[PPN]_2[Cl_2Pd(\mu\text{-}SePh)_2PdCl_2]$, **286**, 155

crystal structure and characterization of an unexpected dinuclear Mn(II)Mn(III) azacryptate, **287**, 14

formation pathway of a Mn(IV),Mn(IV) bis(μ-oxo) dimer that incorporates alkali and alkaline earth cations and electron transfer properties of the dimer, **294**, 133

interaction between heterobinuclear molecules and nature of the ground spin state in oximato-bridged $[Cu^{II}M^{II}]_2$ bis-binuclear complexes (M = Cu, Ni, Mn): crystal structure of [Cu(pdmg)Ni(Me_3[12]N_3)(EtOH)]$(ClO_4)_2$, **288**, 57

isocyanide derivatives of manganese cobalt nonacarbonyl: synthesis and characterization, **288**, 159

polyiodides of transition metal trischelate cations: syntheses, structures, and spectral and electrical conductivity studies of $[Mn(phen)_3](I_3)_2$ and $[Mn(bpy)_3](I_3)_{1.5}(I_8)_{0.25}$, **284**, 158

reactions of $Cp'(CO)_2MnPPh_2H$ with CH_3COCl and $CH_3S(O)_2Cl$ in THF/triethylamine: evidence of the first complex stabilization of a phosphorus homolog of the sulfonamides, **288**, 101

structure and magnetic properties of dinuclear [Mn(III)(salen)$(H_2O)]_2(ClO_4)_2$ and polynuclear [Mn(III)(salen)$(NO_3)]_n$, **290**, 8

structure and physical properties of hexacyanomanganate(III), $[Mn^{III}(CN)_6]^{3-}$, **284**, 175

studies of the binding modes of carboxylate donors with manganese(III) complexes containing tetradentate Schiff base ligands. Crystal structures of the complexes $[Mn(3\text{-}CH_3Osalpn)(HO_2CC_6H_4CO_2)]_n$ and $[\{Mn(5\text{-}NO_2salpn)(OH)(H_2O)\}\{Mn(5\text{-}NO_2salpn)(HO_2CC_6H_4CO_2)(H_2O)\}]\cdot C_2H_5OH$, **293**, 210

synthesis, crystal structures and magnetic properties of trinuclear oxo-centered manganese complexes of the general formula $[Mn_3O(O_2CR)_6L_3]^{z+}$ (R = Me, Et, Ph, CH_2Cl; L = 3-methylpyridine or water $z = 1$, 0), **287**, 82

synthesis, structure and magnetic properties of oxo-centered trinuclear manganese complex $[Mn_3O(O_2CCH_3)_6(C_5H_5N)_3]\cdot ClO_4$, **294**, 109

Manganese coordination complexes

synthesis and properties of hydrophobic $[Mn_2^{IV}(\mu\text{-}O)_3(L)_2]^{2+}$ complexes, derived from alkyl substituted 1,4,7-triazacyclononane ligands, **295**, 189

Manganese oxide

a new synthetic route to todorokite-type manganese oxides, **294**, 123

Matrix isolation

matrix isolation investigation of the reaction of B_2H_6 with carbon–carbon multiple bonds, **286**, 1

Mechanochemical reactions

mechanochemical reactions in the chemistry of boranes, **289**, 51

Mellitic acid compounds

syntheses and structures of Group 1 mellitate compounds, $K_6[C_6(COO)_6]\cdot 8H_2O$ and $Cs_5[C_6(COO)_6H]\cdot 7H_2O$, **286**, 207

72-membered macrocycle

a two-dimensional network constructed via 72-membered heart-shaped macrocycles, a copper(II) complex with 1,3,5-tris(imidazol-1-ylmethyl)-2,4,6-trimethylbenzene and diethylenetriamine ligands, **295**, 129

Mercury complexes

formation of donor–acceptor Fe(0)–Hg(II) bond in separation and stabilization of optically active iron(0) phosphine complexes. Absolute configuration of (+)-(*R*)-$(CO)_4Fe(\mu\text{-}EtPhPpy)HgCl_2$, **293**, 106

mercury(II) compounds with 1,3-imidazole-2-thione and its 1-methyl analogue. Preparative and NMR spectroscopic studies. The crystal structures of di-μ-iodo-bis[iodo(1,3-imidazolium-2-thiolato-*S*)mercury(II)], bis[bromo(1,3-imidazolium-2-thiolato-*S*)]mercury(II) and bis[μ-(1-*N*-methyl-1,3-imidazole-2-thiolato-*S*)]mercury(II), **294**, 35

preparation and spectroscopic properties of the complexes of mercuric thiocyanate with pyridine-2-thione and pyridine-2-carboxylic acid. Crystal and molecular structure of two polymorphs of $Hg(SCN)_2(C_5H_5NS)_2$, **285**, 208

reaction of $M(CO)Cl(L)_2$ with mercury(II) chloride (M = Ir, Rh; L = Ph_2Ppy, $P(fur)_3$) (Ph_2Ppy = 2-(diphenylphosphino)-pyridine, $P(fur)_3$ = tri-(2-furyl)phosphine), **284**, 119

Metal complexes
absence of 1,3-metallotropic shifts in $M(CO)_5$ complexes of pyrimidine and 1,3,5-triazine (M = W, Mo, Cr), **292**, 225
complexation of macrocyclic ligands with relatively non-solvated metal ions generated in dichloromethane by electrochemical oxidation of amalgam electrodes, **293**, 223
effects of electronic structure on chiral induction in outer-sphere electron transfer reactions of complexes with the C_3 symmetric ligand, 1,4,7-triazacyclononane-1,4,7-tris[2′(*R*)-2′-propionate](-3), **290**, 197
structure and magnetic properties of *catena*-(μ-CrO_4-*O*,*O*′)[M^{II}(L)]·5H_2O (M = Ni^{II}, Cu^{II}; L = 3,14-dimethyl-2,6,13,17-tetraazatricyclo[14,4.0^{1,18},0^{7,12}]docosane), **293**, 100
the bonding and structure of B–N sandwich complexes, **288**, 17

Metal coordination polymer
synthesis and characterization of a metal coordination polymer consisting of ruthenium(III) β-diketone units linked by butadiyne bridges, **292**, 64

Metallaboranes
an approach to *megalo*-boranes. Mixed and multiple cluster fusions involving iridaborane and platinaborane cluster compounds. Crystal structure determinations by conventional and synchrotron methods, **289**, 95
[4-{(*p*-cym)Ru}-5-{$(Ph_3P)_2(CO)Os$}B_4H_8]: the first *nido*-heterobimetallahexaborane cluster, **289**, 134

Metallacyceic complexes
platina- and palladalactam complexes derived from 2-benzoylacetanilide; syntheses and X-ray structure of [Pd{NPhC(O)CHC(O)Ph}(bipy)]·CH_2Cl_2, **292**, 260

Metallated complexes
synthesis and characterization of a series of hexadecachlorinated phthalocyanines, **293**, 80

Metallocylic complexes
synthesis, structure and EPR spectroscopy of a vanadium(IV)–amide metallacyclic complex, **285**, 134

Metallodendrimers
retention of optical activity during conversion of Λ-[Ru(1,10-phenanthroline)$_3$]$^{2+}$ to Λ-[Ru(1,10-phenanthroline-5,6-dione)$_3$]$^{2+}$ and Λ-[Ru(dipyrido[*a*:3,2-*h*:2′,3′-*c*-]-phenazine)$_3$]$^{2+}$, **293**, 37

Metallomesogens
synthesis and characterisation of main-chain oligomeric cyclopalladated azobenzene complexes, **292**, 163

Metal–metal bond complexes
metal–metal multiply-bonded complexes of technetium. Part 8. Synthesis and characterization of the α- and β-isomers of $Tc_2Cl_4(dppe)_2$, **288**, 69

Metal–metal bonds
relativistic density functional study on the bimetallic cluster $[Pt_3Fe_3(CO)_{15}]^{n-}$ (n = 0, 1, 2), **286**, 30

Methanide complexes
synthesis of gold ferrocenyl substituted ylides and methanides: crystal structure of $[FcCH(AuPPh_3)PPh_3]ClO_4$ [Fc = (η^5-C_5H_5)Fe(η^5-C_5H_4)], **291**, 60

Methyl ferrocenes
the quadrupole hyperfine interaction of iron organometallics: an interpretation of its temperature dependence, **291**, 74

Michael additions
copper-mediated 1,6-, 1,8-, 1,10- and 1,12-addition and 1,5-substitution reactions in organic synthesis, **296**, 1

Mixed ligand complexes
preparation spectroscopy and conductivity of [Pt(SS)(NN)]-type complexes and their iodine oxidized complexes, **284**, 112
structural studies of ReO(V) mixed ligand [SNS][Cl] and [SNS][S] complexes, **287**, 142
study on the formation of mixed ligand oxorhenium and oxotechnetium complexes (SNS/S combination), **295**, 1
synthesis and characterization of five-coordinate rhenium(V) and technetium(V) mixed ligand bifunctional complexes carrying the SNS/S or the SNN/S donor atom set. Crystal structure of ReO{[$(C_2H_5)_2NCH_2CH_2N(CH_2CH_2S)_2$](*p*-$H_2N$–PhS)} and ReO{[$(CH_2)_4NCH_2CH_2NCH_2CH_2S$](*p*-$H_2N$–PhS)}, **285**, 97
crystal structure and characterization of an unexpected dinuclear Mn(II)Mn(III) azacryptate, **287**, 14
mixed-valence states of iron long-chain carboxylate complexes, **291**, 127
molecular orbital-theoretical studies of a dinuclear Fe(II)–Fe(III) complex, **291**, 82
structural and spectroscopic characterisation of the mixed valence platinum complex $PtdabI_3$ (dab = *o*-diaminobenzene), **287**, 218
widely differing stabilities of molecule-bridged cyanodiiron(III, II) species in non-aqueous solvents, **291**, 66

Mixed valence state
spectroscopic and magnetic properties of Chevreul's salt, a mixed valence copper sulfite $Cu_3(SO_3)_2{\cdot}2H_2O$, **295**, 125

MLCT absorptions
synthesis, physical and photophysical properties of Ru(II) pyridyl-pyrimidine and pyridyl-pyrimidinone complexes: effect of substituents on the pyrimidine ring, **296**, 114

MO calculations
molecular orbital-theoretical studies of a dinuclear Fe(II)–Fe(III) complex, **291**, 82

MOCVD
formation of FeSi and $FeSi_2$ films from *cis*-$Fe(SiCl_3)_2(CO)_4$ by MOCVD–precursor versus substrate control, **291**, 380

Molecular dynamics
the quadrupole hyperfine interaction of iron organometallics: an interpretation of its temperature dependence, **291**, 74

Molecular magnetism
relativistic density functional study on the bimetallic cluster $[Pt_3Fe_3(CO)_{15}]^{n-}$ (n = 0, 1, 2), **286**, 30

Molecular mechanics
structural aspects of cesium ion selectivity by tetrabenzo-24-crown-8, **290**, 86

Molecular orbital theory
early versus late transition metals. Electronic structure of *nido*-2-$CpML_nB_4H_8$, $CpML_n$ = $CpTaCl_2$, $CpWH_3$ and CpCo, **289**, 85

Molecular recognition
molecular recognition of dipeptides. Catalysis of deuteration and hydrolysis of glycylglycine by dinuclear OBISDIEN Zn(II) complexes, **287**, 134

Molecular sieves
hydrothermal synthesis of microporous materials with the direct incorporation of porphyrin molecules, **294**, 179

Molybdate
kinetics of the molybdate and tungstate catalyzed oxidation of iodide by hydrogen peroxide, **288**, 229

Molybdate complexes
cluster synthesis by the reactions of [$Cp'_2Fe_2S_4$] with transition-metal complexes (Cp' = C_5Me_5, 1,3-$C_5H_3(SiMe_3)_2$), **291**, 395
dipyridylamine-ligated octamolybdate clusters aggregated into extended ribbons and layers via hydrogen bonding, **292**, 131

Molybdenum arsenate
hydrothermal synthesis and crystal structure of a new molybdenum(VI) arsenate(III), $Co^{III}(en)_3H_3O[(Co^{II}O_6)Mo^{VI}{}_6O_{18}(As^{III}{}_3O_3)_2]\cdot 2H_2O$, **295**, 244
Molybdenum complexes
a *cyclo*-tetraphosphoxane cage complex of molybdenum(0) and palladium(II), **287**, 223
a molybdenum(0) complex of bis(hydroxymethyl)phenylphosphine, **287**, 209
an EXAFS study of Mo(VI) and Mo(V) complexes with proline dithiocarbamate, **288**, 127
dimolybdenum complexes of bis(2-pyridyl)amine (Hdpa) ligand and its anion (dpa^-); transformation of chelating η^2-Hdpa to bridging η^2-dpa^-, **288**, 82
electrochemical and ^{95}Mo NMR studies of triply-bridged dinuclear oxomolybdenum(V) complexes with various dithiocarbamates, **292**, 7
ligand substitution kinetics in $M(CO)_4$ ($\eta^{2,2}$-1,5-cyclooctadiene) complexes (M = Cr, Mo, W) — substitution of 1,5-cyclooctadiene by bis(diphenylphosphino)alkanes, **284**, 229
preparation of cyclopentadienyl molybdenum clusters linked by a cyclotetradeca-1,8-diyne ligand, **294**, 232
ring-opening polymerization of norbornene by η-cycloheptatrienyl molybdenum and tungsten complexes, **288**, 226
syntheses, crystal structures and nonlinear optical properties of heteronuclear clusters $[MS_4Cu_4(SCN)_2(NC_5H_5)_6]$ (M = W, Mo), **294**, 73
synthesis and structural characterization of face-sharing bioctahedral complexes containing poly(pyrazolyl)borate ligands: $[HB(Me_2Pz)_3BH][X_3Mo(\mu\text{-}X)_2(\mu\text{-}H)MoTp^*]$ (X = Cl or Br; $Tp^* = HB(Me_2Pz)_3$; Pz = pyrazolyl), **292**, 182
synthesis, reactivity, and structures of (μ-acetamidato)-hexakis(acetonitrilo)dimolybdenum(II) tris(tetrafluoroborate) and derivatives, **287**, 159
the effect of replacing carbonyl with acetonitrile on the reactivity of the bimetallic system $[Mo_2Cp_2(\mu\text{-}SMe)_3(L)_2]^+$ (L = CO, CH_3CN). Crystal structure of $[Mo_2Cp_2(\mu\text{-}SMe)_2(CH_3CN)_4](BF_4)_2$, **284**, 107
the nitrido complex $(NEt_4)[MoCl_3(N)(OH)(MeCN)]$, **295**, 249
Molybdoarsenate complexes
formation and conversion of molybdophosphate and -arsenate complexes in aqueous solution, **284**, 237
Molybdophosphate complexes
formation and conversion of molybdophosphate and -arsenate complexes in aqueous solution, **284**, 237
^{95}Mo NMR
electrochemical and ^{95}Mo NMR studies of triply-bridged dinuclear oxomolybdenum(V) complexes with various dithiocarbamates, **292**, 7
Monocyclic ligand complexes
asymmetric compartmental macrocyclic ligands and related mononuclear and hetero-dinuclear complexes with d- and/or f-metal ions, **293**, 178
Mononuclear complexes
mononuclear lanthanide complexes of tripodal ligands: synthesis and spectroscopic studies, **285**, 49
Mo–S cluster complexes
the first binuclear Mo–S cluster compound containing citrate ligands, $K_5(NH_4)[Mo_2O_2(\mu\text{-}S)_2(C_6H_4O_7)_2]\cdot CH_3OH\cdot 5H_2O$, characterized by X-ray single crystal structure determination, **285**, 152
Mössbauer spectra
Mössbauer studies of different types of hematin, **286**, 229
Mössbauer spectroscopy
about the synthesis of the prismane $[NEt_4]_2[Fe_6S_6I_6]$, **284**, 296
lattice effects in the Mössbauer spectra of salts of $[Fe_4S_4\{S(CH_2)_nOH\}_4]^{2-}$. Crystal structures of $[PPh_4]_2[Fe_4S_4\{S(CH_2)_nOH\}_4]$ (n = 2, 3 and 4), **291**, 101
mixed-valence states of iron long-chain carboxylate complexes, **291**, 127
strained, ring tilted dicarbon-bridged [2]ferrocenophanes and ferrocene revisited: ^{57}Fe Mössbauer spectroscopic study of bonding, hyperfine interactions, and lattice dynamics, **287**, 152
the quadrupole hyperfine interaction of iron organometallics: an interpretation of its temperature dependence, **291**, 74
Mössbauer study
syntheses, properties and Mössbauer studies of cyanamide and cyanoguanidine complexes of iron(II). Crystal structures of *trans*-$[FeH(NCNH_2)(Ph_2PCH_2CH_2PPh_2)_2][BF_4]$ and *trans*-$[Fe(NCNEt_2)_2(Et_2PCH_2CH_2PEt_2)_2][BF_4]_2$, **291**, 39
Multidentate ligand complexes
the nickel(II) complex of a new multidentate ligand containing both imidazole and pyrazole moieties, **288**, 239
Multimetallic complexes
multimetallic, double-stranded helical complexes derived from hexa(*n*-propylthio)novipyridine: synthesis, structure and redox properties, **288**, 189

Nanoclusters
synthesis of organo(siloxo)platinum and -palladium complexes and preparation of supported nanoclusters by facile ligand reduction, **294**, 266
Naphthyridine complexes
di- and trinuclear Pd(II) and Pt(II) complexes containing bridging 2,7-disubstituted naphthyridines, **287**, 72
Natural bile acid complexes
organometallic radiopharmaceuticals: rhenium(I) carbonyl complexes of natural bile acids and derivatives, **286**, 46
Neodymium complexes
neodymium and europium picrate complexes with 3-picoline-*N*-oxide (3-picNO), **292**, 249
Neutron diffraction
hydrogen bonding in *cis*-bis(L-alaninato)copper(II): a single crystal neutron diffraction study, **288**, 29
Nickel complexes
ab initio calculations and X-ray structure determinations of dichloro(*O*-*O*′-*O*″-1,3,5-cyclohexanetriol)copper(II) and bis(*O*-*O*′-*O*″-1,3,5-hexanetriol)nickel(II), **284**, 284
a new structurally constrained dioxocyclam ligand: synthesis, characterization and molecular structure comparison of its nickel(II) complex with the nickel(II) complex of dioxocyclam, **292**, 254
an optically active nickel(II) Schiff base coordination compound *N*,*N*′-(1*R*,2*R*)-(−)-1,2-cyclohexylenebis(2-hydroxyacetophenonylideneiminato)nickel(II), **293**, 239
a novel oligomerization versus metal chelation: synthesis and crystal structure of pentakis[di-μ-(diisopropylaminoethanethiolato)nickel], **294**, 83
chelating and ligand-transfer studies of the complexes *cis*-$[Mn(CO)_4(SeR)_2]^-$: crystal structures of heterotrinuclear $[(CO)_4Mn(\mu\text{-}SeMe)_2Ni(\mu\text{-}SeMe)_2Mn(CO)_4]$ with a distorted square planar Ni^{II}–selenolate core and $[PPN]_2[Cl_2Pd(\mu\text{-}SePh)_2PdCl_2]$, **286**, 155
control of the coordination mode of NO_2^- by the nature of the amine ligand: the importance of H-bond. Crystal structures of the [*mer*-Ni(dpt)$(NO_2)_2$] (**I**) and [*trans*-Ni(333-tet)$(NO_2)_2$] (**II**), **292**, 149
coordination of *ortho* and *para* quinones and oxidation reduction processes in trimethylphosphine complexes of cobalt and nickel, **287**, 167
crystal structures and magnetic studies of six new nickel(II) amine μ-oxalato compounds, **295**, 91
diphenoxo-bridged dinuclear nickel(II) complex of a dinucleating macrocycle with $N(amine)_6O_2$ metal-binding sites: synthesis, X-ray crystal structure and electrochemistry, **287**, 212

fluctuations between square-planar and tetrahedral coordination geometry with bis(2-benzimidazolyl)alkane ligands. Synthesis, spectroscopic properties and X-ray crystal structure of four representative examples, **290**, 105
interaction between heterobinuclear molecules and nature of the ground spin state in oximato-bridged $[Cu^{II}M^{II}]_2$ bis-binuclear complexes (M = Cu, Ni, Mn): crystal structure of $[Cu(pdmg)Ni(Me_3[12]N_3)(EtOH)](ClO_4)_2$, **288**, 57
metal(II) complexes of the carbonylnitronate ligands $O_2NCHC(O)R^-$ (R = Ph, OMe or OEt) and X-ray structure of (DMANH)$[Ni(O_2NCHC(O)Ph)_3]$ (DMAN = 1,8-bis(dimethylamino)naphthalene), **285**, 18
nickel complexes of tetradentate asymmetric ligands, **290**, 14
nickel(II) and palladium(II) complexes with acylhydrazone ligands of α-diketones: the electronic and steric factors affecting the formation of the dimeric palladium(II) complexes, **295**, 171
$Ni(COD)_2$: an efficient promoter in the construction of 5-cyclopent-2-enone derivatives from allyl halides, alkynes and carbon monoxide, **296**, 94
non-linear optical absorption of $[\{Ni(mnt)_2\}_2]^{2-}$ and its unique counter ion, a disodium triple-decker dication $[Na_2(benzo-15-crown-5)_3(H_2O)_2]^{2+}$, **293**, 89
preliminary studies of $Ni(dmen)_2(CF_3CO_2)_2$, $Ni(deen)_2(CF_3CO_2)_2$ and $Ni(en)(CF_3C(O)CHC(O)CF_3)_2$ as precursors for metalorganic chemical vapour deposition: structures, thermal behaviours and vapour pressures, **288**, 233
spontaneous formation of trinickel clusters containing μ^3-imido bridges from 2-nitrophenols and $Ni(PMe_3)_4$, **287**, 199
structural characterization of a new ligand mode of 2,6-diacetylpyridine bis(semicarbazone), H_2daps, **292**, 121
syntheses and structures of nickel(II) and copper(II) complexes with a 14-membered tetraaza macrocyclic ligand, **286**, 67
synthesis and coordination chemistry of 1-phenyl-1-phospha-4,7-dithiacyclononane, **294**, 170
synthesis and X-ray crystallographic structures of dichloronickel(II) and -palladium(II) complexes containing the biphenyl diamine ligand 6-(2-dimethylaminoethyl)-6,7-dihydro-5-H-dibenz[c,e]azepin, **294**, 28
synthesis of the ligand *N*,*N*′-bis(2-tert-butylthiobenzenylidene)-diethylenetriamine; its reactivity with nickel(II) salts, **287**, 105
synthesis, vibrational study, crystal structure and density functional calculations of $[Ni(dien)_2]^{2+}$ complexes. Configurational and conformational study of $[M(dien)_2]^{n+}$ complexes, **290**, 57
template synthesis, crystal structure, and solution behavior of a hexaaza macrocyclic nickel(II) complex containing two *N*-aminoethyl pendant arms, **293**, 140
terminal S-coordinated thiocyanate in a nickel(II) complex. X-ray structures of $Ni(poph)(H_2O)(SCN)(NCS)$ and $Ni(poqh)(H_2O)(NCS)_2$ (poph = 2-pyridinecarboxaldehyde 1-oxide 2′-pyridinylhydrazone, poqh = 2-pyridinecarboxaldehyde 1-oxide 2′-quinolynylhydrazone), **285**, 297
the hydrothermal synthesis and structural characterization of a new class of compounds: nickel organoamine-halocadmates, **292**, 41
the nickel(II) complex of a new multidentate ligand containing both imidazole and pyrazole moieties, **288**, 239
the optically active and racemic products of monocondensation of 1,2-diaminopropane and 2,4-pentadione as ligands and precursors for preparation of unsymmetrical Schiff bases and their nickel(II) complexes, **293**, 115
tridentate N_3 capping coordination behaviour of potentially multi-site coordinating cyclotriphosphazenes: synthesis and spectroscopic studies of 2,2-spiro(1,3-propanediamino)-4,4,6,6-tetrakis(3,5-dimethylpyrazolyl)cyclotriphosphazene, 2,2-diamino-4,4,6,6-tetrakis(3,5-dimethylpyrazolyl)-cyclotriphosphazene and their mononuclear complexes; X-ray structure of ATPCTP·$CoCl_2$, **286**, 127
Nickel on charcoal
kumada couplings catalyzed by nickel on charcoal (Ni/C), **296**, 164
nido and *arachno* structures
mechanistic aspects of boron hydride reactions, **289**, 1
Niobium complexes
synthesis and structure of niobium and tantalum derivatives of bis(dicyclohexylphosphino)methane (dcpm), **288**, 35
Nitric oxide complexes
a reversible NO complex of $[Fe^{II}$(edampda)] and the $[Ni^{II}$(edampda)] analogue (edampda^{2-} = *N*,*N*′-bis(pyridylmethyl)ethylenediamine-*N*,*N*′-diacetate), **286**, 197
Nitride complexes
vanadium phenoxide complexes with oxide, nitride or hydrazide co-ligands: preparation and crystal structures of $[V(OC_6H_3Pr^i_2\text{-}2,6)_3NLi(C_4H_8O)_3]$, $[\{VO_2(OC_6H_3Pr^i_2\text{-}2,6)_2\}_2\text{-}\{\mu\text{-}Li(C_4H_8O)_2\}_2]$ and $[V(NNMe_2)(OC_6H_3Pr^i_2\text{-}2,6)_3]$, **285**, 178
Nitrides
synthesis of new Group III fluoride–ammonia adducts in supercritical ammonia: structures of $AlF_3(NH_3)_2$ and $InF_2(NH_2)(NH_3)$, **294**, 200
Nitrido-bridged complexes
heterobimetallic nitrido-bridged Ru(II)NOs(VIII) and Ru(II)-NOs(VI) complexes containing ruthenium porphyrins, **290**, 28
Nitrido complexes
the nitrido complex $(NEt_4)[MoCl_3(N)(OH)(MeCN)]$, **295**, 249
Nitrile
nitrite ion-promoted hydration of acetonitrile by the bis(μ-hydroxo)bis{tris(2-pyridylmethyl)amine)chromium(III)} ion, **288**, 211
Nitrile complexes
iminoacylation: 2. Addition of alkylated hydroxylamines via oxygen to platinum(IV)-bound nitriles, **292**, 272
synthesis and formation of metal complexes of 4-alkynyl and 4-cyano-2,6-diisopropylphenylisocyanides, **284**, 205
synthesis, reactivity, and structures of (μ-acetamidato)-hexakis(acetonitrilo)dimolybdenum(II) tris(tetrafluoroborate) and derivatives, **287**, 159
Nitrile oxide–olefin cycloadditions
synthesis and reactivity of acyclic (pentadienyl)iron(1 +) cations: model studies for the preparation of the 8*E*,10*Z*,16*E*,18*E*-tetraene segment of macrolactin A, **296**, 261
Nitrilotriacetic acid complexes
solution studies of complexes of iron(III) with iminodiacetic, alkyl-substituted iminodiacetic and nitrilotriacetic acids by potentiometry and cyclic voltammetry, **291**, 158
Nitrite
crystal structures and magnetic studies of six new nickel(II) amine μ-oxalato compounds, **295**, 91
nitrite ion-promoted hydration of acetonitrile by the bis(μ-hydroxo)bis{tris(2-pyridylmethyl)amine)chromium(III)} ion, **288**, 211
Nitrite complexes
control of the coordination mode of NO_2^- by the nature of the amine ligand: the importance of H-bond. Crystal structures of the [*mer*-Ni(dpt)$(NO_2)_2$] (**I**) and [*trans*-Ni(333-tet)$(NO_2)_2$] (**II**), **292**, 149
Nitrito complexes
an alternating copper(II) chain with bridging oxamidato and nitrito ligands: crystal structure and magnetic properties of $[Cu(NO_2)_2CuL]_n$ (L = *N*,*N*′-bis(2-methyl-2-aminopropyl) oxamide), **294**, 8

Nitrogenase
the first binuclear Mo–S cluster compound containing citrate ligands, $K_5(NH_4)[Mo_2O_2(\mu\text{-}S)_2(C_6H_4O_7)_2]\cdot CH_3OH\cdot 5H_2O$, characterized by X-ray single crystal structure determination, **285**, 152
Nitrosyl complexes
evidence for a new nitrosyl-to-nitro reaction and an acetamide formation in the $\{RuNO\}^6$-type complex, *cis*-$[Ru(NO)(CH_3CN)(bpy)_2]^{3+}$, **290**, 113
reaction of hydrous nitrosylruthenium trichloride with 2-pyridinemethanol, **290**, 36
reduction of the NO^+ ligand in the pentacyanonitrosylosmate(II) ion, **285**, 129
Nitroxide complexes
synthesis, crystal structure and magnetic properties of a copper(II)–nitroxide linear chain complex, **287**, 181
NMR
new phosphino silver(I) derivatives of hydrotris(3-methyl-2-thioxo-1-imidazolyl)borate. X-ray crystal structure of tricyclohexylphosphinesilver(I)-hydrotris(3-methyl-2-thioxo-1-imidazolyl)borate, **285**, 81
NMR spectroscopy
a dinuclear platinum(II) complex with hydrazine and chloro bridges: $(NPr^n_4)[PtCl_2(\mu\text{-}Cl)(\mu\text{-}N_2H_4)PtCl_2]$, **284**, 103
structural studies of ReO(V) mixed ligand [SNS][Cl] and [SNS][S] complexes, **287**, 142
synthesis, structures, and linkage isomerism of (allylbenzylmalonate)platinum(II) complexes, **295**, 214
N,N,N-trimethylglycine
characterization and crystal structure of copper(II) complex with betaine having a water-wheel-like structure, **295**, 121
N,O-bidentate ligands
X-ray crystal structures of copper(II) nitrate complexes of two isomeric N,O-chelating ligands, **284**, 273
NOE
alkyl derivatives of $[B_6H_6]^{2-}$: NMR and vibrational spectra and crystal structure of $(Ph_4P)[B_6H_6CH_2Ph]$, **289**, 70
Non-linear optics
$[Cd_2(4,4'\text{-bipy})_2(H_2O)_3(SO_4)_2]\cdot 3H_2O$: a novel three-dimensional coordination polymer with nonlinear optical properties, **294**, 95
non-linear optical absorption of $[\{Ni(mnt)_2\}_2]^{2-}$ and its unique counter ion, a disodium triple-decker dication $[Na_2(benzo\text{-}15\text{-}crown\text{-}5)_3(H_2O)_2]^{2+}$, **293**, 89
syntheses, crystal structures and nonlinear optical properties of heteronuclear clusters $[MS_4Cu_4(SCN)_2(NC_5H_5)_6]$ (M = W, Mo), **294**, 73
Norbornene
ring-opening polymerization of norbornene by η-cycloheptatrienyl molybdenum and tungsten complexes, **288**, 226
Novipyridine complexes
multimetallic, double-stranded helical complexes derived from hexa(*n*-propylthio)novipyridine: synthesis, structure and redox properties, **288**, 189
Nuclear magnetic resonance
application of the NOE experiment to the analysis of boron hydride derivatives: confirming the assignments of the *pseudocloso*-complex [1,2-Ph_2-3-{Cp*}-3,1,2-$IrC_2B_9H_9$] (Cp* = η^5-C_5Me_5) and the *closo*-compounds 1-Ph-1,2-$C_2B_{10}H_{11}$ and 1-Ph-2-Me-1,2-$C_2B_{10}H_{10}$, **289**, 125
conversion of site-specific deuterated *closo*-1,2- and 1,7-$C_2B_{10}H_{12}$ to deuterated *nido*-7,8- and 7,9-$[C_2B_9H_{12}]^-$ ions, respectively; mechanistic inferences. Comparisons of GIAO-NMR generated $[C_2B_9H_{12}]^-$ chemical shifts with experimental values, and consideration of two principal isomers of the 7,8-$[C_2B_9H_{12}]^-$ ion by comparison of geometry optimization data, NMR chemical shift data, and NMR coupling data to respective experimental data, **289**, 11
NMR and NQR studies of boron in vitreous and crystalline borates, **289**, 158
NMR as a tool for elucidation of structures and estimation of electron distribution in boranes and their derivatives, **289**, 20
NMR investigation of Pd(II)–Pd(0) reduction in the presence of mono- and ditertiary phosphines, **286**, 93
Nuclear magnetic resonance imaging
interaction of ATP with a Gd^{3+} complex employed as paramagnetic contrast agent in NMR imaging, **288**, 244
Nuclear quadrupole resonance
NMR and NQR studies of boron in vitreous and crystalline borates, **289**, 158
Nucleophilic substitution
nucleophilic substitutions of iodide with bromides in the $Fe(CO)_2(PMe_3)_2MeI$ complex, **288**, 200
Nucleotide complexes
interaction of ATP with a Gd^{3+} complex employed as paramagnetic contrast agent in NMR imaging, **288**, 244

O_2-Binding affinity
5,10,15,20 - tetrakis(α,α,α,α - *o* - pivalamidophenyl)porphinatoiron(II) bearing a covalently linked axial imidazole via *m*-aminobenzoic acid. Synthesis and influence of imidazole basicity on O_2-binding affinity, **295**, 234
Octamolybdate
crystal structures of two new heptamolybdates and of a pyrazole incorporating a γ-octamolybdate anion, **295**, 106
Olefin complexes
reactions of H_2, silanes, and olefins with superelectrophilic cationic rhenium complexes: heterolytic cleavage of H_2 and relation to the structure and function of hydrogenases, **294**, 240
Open framework
magnetic iron phosphates with an open framework, **291**, 317
Optical activity
retention of optical activity during conversion of Λ-$[Ru(1,10\text{-phenanthroline})_3]^{2+}$ to Λ-$[Ru(1,10\text{-phenanthroline-5,6-dione})_3]^{2+}$ and Λ-$[Ru(dipyrido[a:3,2\text{-}h:2',3'\text{-}c]\text{-phenazine})_3]^{2+}$, **293**, 37
Optical properties
synthesis and photophysical properties of *trans*-dithiocyanato bis(4,4′-dicarboxylic acid-2,2′-bipyridine) ruthenium(II) charge transfer sensitizer, **296**, 250
Organoboranes
spirocyclic stannole derivatives by 1,1-organoboration of 2,2-bis(1-alkynyl)-1,3-di-tert-butyl-1,3,2-diazastannacyclohexanes, **296**, 26
Organocopper reagents
copper-mediated 1,6-, 1,8-, 1,10- and 1,12-addition and 1,5-substitution reactions in organic synthesis, **296**, 1
Organogermanium complexes
synthesis of five-coordinate Pt(II) complexes containing organogermanium fragments in axial position. Structure of $[PtCl(GePh_2Cl)(2,9\text{-}Me_2\text{-}1,10\text{-phen})(CH_2{=}CH_2)]$, **285**, 70
Organoiron complexes
novel organoiron compounds resulting from the attempted syntheses of dibenzofulvalene complexes, **291**, 109
Organometallic polymers
synthesis and characterisation of main-chain oligomeric cyclopalladated azobenzene complexes, **292**, 163
Organometallics
synthetic reactions using organometallics in water. Aldol and allylation reactions catalyzed by Lewis acid–surfactant-combined catalysts/Brønsted acids systems, **296**, 158
Organotin complexes
synthesis and structures of 2-$[Me_2NCH_2CH_2N(Me)CH_2]C_6H_4$ substituted tin compounds, **290**, 100

Organotin compounds
spirocyclic stannole derivatives by 1,1-organoboration of 2,2-bis(1-alkynyl)-1,3-di-tert-butyl-1,3,2-diazastannacyclohexanes, **296**, 26
synthesis and characterization of compounds of di- and tributyltin chloride with adenine and guanine mononucleotides, **285**, 60

Ortho lithiation
diastereoselective *ortho*-lithiation of tricarbonylchromium complexes of chiral benzaldehyde acetals, **296**, 80

Osmium chemistry
[4-{(*p*-cym)Ru}-5-{$(Ph_3P)_2(CO)Os$}B_4H_8]: the first *nido*-heterobimetallahexaborane cluster, **289**, 134

Osmium complexes
heterobimetallic nitrido-bridged Ru(II)NOs(VIII) and Ru(II)NOs(VI) complexes containing ruthenium porphyrins, **290**, 28
oxidative addition reactions of benzo[*b*]thiophene and dibenzothiophene with $[Os_3(CO)_{10}(MeCN)_2]$: room-temperature ring opening of benzo[*b*]thiophene at triosmium clusters, **285**, 277
reduction of the NO^+ ligand in the pentacyanonitrosylosmate(II) ion, **285**, 129

Overhauser effect
application of the NOE experiment to the analysis of boron hydride derivatives: confirming the assignments of the *pseudocloso*-complex [1,2-Ph_2-3-{Cp*}-3,1,2-$IrC_2B_9H_9$] (Cp* = η^5-C_5Me_5) and the *closo*-compounds 1-Ph-1,2-$C_2B_{10}H_{11}$ and 1-Ph-2-Me-1,2-$C_2B_{10}H_{10}$, **289**, 125

Oxadiazole complexes
coordination complexes of bifunctional compounds: I. Synthesis and properties of bis[5-(2-oxyphenyl)-3-phenyl-1,2,4-oxadiazolyl]copper(II). A fluorescent coordination compound of Cu(II), **292**, 1

Oxalate complexes
infinite two-dimensional sheet-like array of a 2,2′-bipyrimidine-bis(oxalato)chromate(III) complex: synthesis, crystal structure and magnetic properties of $[K(H_2O)Cr(C_2O_4)_2(bpym)]$, **284**, 8

Oxalato complexes
crystal structures and magnetic studies of six new nickel(II) amine μ-oxalato compounds, **295**, 91

Oxamato complexes
synthesis and crystal structure of $Na[Pd(Hpba)]\cdot 2H_2O$ [H_4pba = 1,3-propylenebis(oxamic acid)]. The first bis(oxamato)-palladium(II) complex and an unexpected trideprotonated form of the ligand, **284**, 292

Oxamido complexes
an alternating copper(II) chain with bridging oxamidato and nitrito ligands: crystal structure and magnetic properties of $[Cu(NO_2)_2CuL]_n$ (L = *N*,*N*′-bis(2-methyl-2-aminopropyl) oxamide), **294**, 8

Oxidation
formation pathway of a Mn(IV),Mn(IV) bis(μ-oxo) dimer that incorporates alkali and alkaline earth cations and electron transfer properties of the dimer, **294**, 133
molecular structure of $[PPh_4]_2[Co(C_3S_5)_2]$ [$C_3S_5^{2-}$ = 4,5-disulfanyl-1,3-dithiole-2-thionate(2-)] and spectroscopic and electrical properties of its oxidized species, **295**, 239
oxidation of adamantane by palladium acetate systems, **294**, 99

Oxidation potentials
synthesis, physical and photophysical properties of Ru(II) pyridyl-pyrimidine and pyridyl-pyrimidinone complexes: effect of substituents on the pyrimidine ring, **296**, 114

Oxidation/reduction
coordination of *ortho* and *para* quinones and oxidation reduction processes in trimethylphosphine complexes of cobalt and nickel, **287**, 167

Oximato complexes
interaction between heterobinuclear molecules and nature of the ground spin state in oximato-bridged $[Cu^{II}M^{II}]_2$ bisbinuclear complexes (M = Cu, Ni, Mn): crystal structure of $[Cu(pdmg)Ni(Me_3[12]N_3)(EtOH)](ClO_4)_2$, **288**, 57

Oxime complexes
different chlorination modes of oximes: chlorination of salicylaldoxime coordinated to platinum, **285**, 116
unusual coordination mode of a 2-pyridyl ketone oxime ligand in bis(4-butylphenyl 2-pyridyl ketone oximate)palladium(II), **285**, 138

Oximes
synthesis of transition metal isocyanide complexes containing hydrogen bonding sites in peripheral locations, **285**, 223

Oxo-bridged complexes
crystal structure of (μ-oxo)bis{aquatris(2-pyridylmethyl)amine}-iron(III) perchlorate, **284**, 124
structure and magnetic properties of (μ-oxo)bis[*meso*-tetrakis(*p*-tolyl-porphyrinato)iron(III)] and (μ-oxo)bis[*N*,*N*′-ethylenebis(2-acetylphenoliminato)iron(III)], **290**, 51
the molecular and electronic structure of the asymmetric, trinuclear complex $[LFe^{III}(\mu\text{-}O)(\mu\text{-piv})_2Fe^{III}(\mu\text{-}OH)(\mu\text{-piv})_2Fe^{III}(piv)_2]\cdot C_6H_5CH_3$ (L = 1,4,7-trimethyl-1,4,7-triazacyclononane, piv = pivalate(1-)), **291**, 258

Oxo complexes
deboronation of $HB(3,5\text{-}Me_2C_3HN_2)_3^-$ in a reaction with $[Re(O)Cl_4]^-$, the X-ray crystal structures of [{Re(O)Cl(3,5-$Me_2C_3HN_2H)_3$}(μ-O){$Re(O)Cl_3(3,5\text{-}Me_2C_3HN_2H)$}] and [{$Re(O)Cl_2(3,5\text{-}Me_2C_3HN_2H)_2$}(μ-O){$Re(O)Cl_2(3,5\text{-}Me_2C_3HN_2H)_2$}], **285**, 55
molecular assembly of BINOL–Ti complexes into an active μ_3-oxo titanium catalyst, **296**, 267
synthesis and characterization of oxorhenium-'3 + 1' mixed-thiolate complexes. Crystal and molecular structures of $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}(C_6H_4X\text{-}4\text{-}CH_2S)]$ (X = F, Cl, Br, OMe) and of the pendant thiolate compounds $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}(\eta^1\text{-}SCH_2CH_2SCH_2CH_2SH)]$ and $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}\{\eta^1\text{-}SCH_2CH(OH)CH(OH)CH_2SH\}]$, **284**, 252
variations in capping the $Zr_6O_4(OH)_4$ cluster core: X-ray structure analyses of $[Zr_6(OH)_4O_4(OOC\text{–}CH\text{=}CH_2)_{10}]_2(\mu\text{-}OOC\text{–}CH\text{=}CH_2)_4$ and $Zr_6(OH)_4O_4(OOCR)_{12}(PrOH)$ (R = Ph, CMe = CH_2), **284**, 1
1:1 adducts of triphenyltin chloride with oxovanadium(IV) tetradentate Schiff-base complexes, **293**, 147
coordination mode of dihydroxybenzoic acids in oxorhenium(V) complexes, **294**, 114
electrochemical and ^{95}Mo NMR studies of triply-bridged dinuclear oxomolybdenum(V) complexes with various dithiocarbamates, **292**, 7
mixed-valence states of iron long-chain carboxylate complexes, **291**, 127
synthesis and structural characterisation of rhenium(V) oxo-thioether thiolate complexes with pendant ester or carboxylate groups, **285**, 76
synthesis, characterization and crystal structures of rhenium(V) complexes with diphosphines, **294**, 47
synthesis, crystal structures and magnetic properties of trinuclear oxo-centered manganese complexes of the general formula $[Mn_3O(O_2CR)_6L_3]^{z+}$ (R = Me, Et, Ph, CH_2Cl; L = 3-methylpyridine or water z = 1, 0), **287**, 82
synthesis, structure and magnetic properties of oxo-centered trinuclear manganese complex $[Mn_3O(O_2CC_3H_7)_6(C_5H_5N)_3]\cdot ClO_4$, **294**, 109

Oxomolybdenum complexes
oxygen atom transfer reaction involving oxomolybdenum complexes with sterically bulky thiocarboxylate ligands and biochemical interesting substrates in methanol at neutral pH, **285**, 25
synthesis and characterization of a series of new thiocarboxylate complexes of molybdenum(V), **284**, 223
Oxomolybdenum compounds
hydrothermal assembly and structural characterization of two low-dimensional molybdenum oxide/2,2′-dipyridylamine hybrid materials, **295**, 222
Oxorhenium complexes
structural studies of ReO(V) mixed ligand [SNS][Cl] and [SNS][S] complexes, **287**, 142
Oxovanadium ions
high pressure NMR studies on ligand exchange reactions in the equatorial sites of $[VO(L)_5]^{2+}$ (L = acetonitrile, *N*,*N*-dimethylformamide, and *N*,*N*-dimethylacetamide), **294**, 119
Oxydiacetate complexes
adduct formation between zinc and cadmium oxydiacetates and imidazol. Crystal structures of $[Zn(Him)_2(oda)]_n$ and $[Cd_2(Him)_4(oda)_2(H_2O)]_2 \cdot 6H_2O$ (Him = imidazol, oda = oxydiacetate), **284**, 49
Oxygenation
a polyoxoanion-based iron(II) catechol extradiol dioxygenase: oxygenation of 3,5-di-tert-butylcatechol by the polyoxoanion-supported iron(II) complex $(n\text{-}Bu_4N)_7[(CH_3CN)_3Fe^{II}P_2W_{15}Nb_3O_{62}]$, **291**, 426
Oxygen ligands
the preparation and coordination chemistry of $^iPr_2P(E)NHP(E')^iPr_2$ (E, E′ = Se; E = Se, E′ = S; E = S, E′ = O; E,E′ = O, **290**, 1
Oxygen transfer
oxygen atom transfer reaction involving oxomolybdenum complexes with sterically bulky thiocarboxylate ligands and biochemical interesting substrates in methanol at neutral pH, **285**, 25
Oxythiamine
novel structural aspects of oxythiamine, an antagonist of thiamine. Crystal structures of three salts, (oxythiamineH)(picrolonate)$_2 \cdot 2H_2O$, (oxythiamineH)$(PtCl_6)$, (oxythiamineH)$_2(V_{10}O_{28}H_2) \cdot 8H_2O$, and a metal complex Pt(oxythiamine)$Cl_3 \cdot H_2O$, **295**, 71

Palladium acetate
oxidation of adamantane by palladium acetate systems, **294**, 99
Palladium catalysis
preparation of poly(aryleneethynylene) type π-conjugated polymers constituted of 2-alkylbenzimidazole-4,7-diyl units by using palladium catalyzed cross coupling reaction, **296**, 254
Palladium complexes
a *cyclo*-tetraphosphoxane cage complex of molybdenum(0) and palladium(II), **287**, 223
B-frame-supported bimetallics. Isoelectronic *arachno*-structured $[(PMe_2Ph)_4Pd_2B_8H_{10}]$ and *closo*-structured $[(PMe_3)_4(CO)_2Ir_2B_8H_8]$, **285**, 290
^{13}C *ipso* chemical shifts in the aryl phosphino-thioether complexes $PdX(p\text{-}RC_6H_4)(P,S)$. On resonance effects into the aryl ring, **290**, 95
coordination geometries for palladium and platinum: theoretical studies and the synthesis and structure of tris(indazol-1-yl)borate complexes $MMe_3\{(ind)_2BH\}$, **287**, 27
cross-coupling reactions proceeding through η^1- and η^3-propargyl/allenyl–palladium(II) intermediates, **296**, 37
crystallographic evidence for the coordination performance of $WO_2S_2^{2-}$. Synthesis, characterization and crystal structure of palladium complexes with mixed-ligand thiotungstates and 1,2-dithiolates: $[NBu_4]_2[PdWO_2S_2(C_3S_5)]$ and $[NBu_4]_2[PdWS_4(C_4N_2S_2)]$, **285**, 241
di- and trinuclear Pd(II) and Pt(II) complexes containing bridging 2,7-disubstituted naphthyridines, **287**, 72
metal(II) complexes of the carbonylnitronate ligands $O_2NCHC(O)R^-$ (R = Ph, OMe or OEt) and X-ray structure of $(DMANH)[Ni(O_2NCHC(O)Ph)_3]$ (DMAN = 1,8-bis(dimethylamino)naphthalene), **285**, 18
new palladium(II) complexes with bis(pyrazol-1-yl)alkanes, **295**, 136
nickel(II) and palladium(II) complexes with acylhydrazone ligands of α-diketones: the electronic and steric factors affecting the formation of the dimeric palladium(II) complexes, **295**, 171
NMR investigation of Pd(II)–Pd(0) reduction in the presence of mono- and ditertiary phosphines, **286**, 93
palladium(II) allyl complexes with potentially terdentate ancillary ligands. Mechanism of allyl amination by piperidine, **293**, 44
palladium(II) complexes with hemilabile etherdiphos ligands in the alternating copolymerization of carbon monoxide with olefins, **296**, 103
platina- and palladalactam complexes derived from 2-benzoylacetanilide; syntheses and X-ray structure of $[Pd\{NPhC(O)CHC(O)Ph\}(bipy)] \cdot CH_2Cl_2$, **292**, 260
platinum(II) and palladium(II) complexes with 3,3,8,8-tetramethyl-1,6-dioxa-4,9-diaza-5λ^5-phosphaspiro[4,4]nonane, **295**, 164
platinum(II) and palladium(II) saccharinate complexes, **285**, 145
P ^ N versus N ^ N ligands for the palladium-catalyzed alternating copolymerization of styrene and carbon monoxide, **296**, 45
reactions of bis(silyl)palladium(II) complexes with allyl halides. Synthesis of mono(silyl)palladium(II) halides and X-ray structure of *trans*-$PdCl(SiF_2Ph)(PMe_2Ph)_2$, **296**, 19
reactivity towards some weak protic electrophiles of the di-μ-hydroxo complex $[\{Pd(\mu\text{-}OH)(tmeda)\}_2][BPh_4]_2$: crystal structure of $[\{Pd(tmeda)\}_2(\mu\text{-}OH)(\mu\text{-}HNC_6H_4Cl\text{-}p)][BPh_4]_2$, **287**, 37
stereochemical investigation of a kinetically labile chiral organopalladium complex in solution, **284**, 99
synthesis and characterisation of mono- and binuclear palladium and platinum complexes with organochalcogenides, **285**, 190
synthesis and crystal structure of $Na[Pd(Hpba)] \cdot 2H_2O$ [H_4pba = 1,3-propylenebis(oxamic acid)]. The first bis(oxamato)-palladium(II) complex and an unexpected tride-protonated form of the ligand, **284**, 292
synthesis and X-ray crystallographic structures of dichloronickel(II) and -palladium(II) complexes containing the biphenyl diamine ligand 6-(2-dimethylaminoethyl)-6,7-dihydro-5-H-dibenz[c,e]azepin, **294**, 28
synthesis of diphenylphosphine palladium complexes, **284**, 116
synthesis of organo(siloxo)platinum and -palladium complexes and preparation of supported nanoclusters by facile ligand reduction, **294**, 266
synthesis of transition metal isocyanide complexes containing hydrogen bonding sites in peripheral locations, **285**, 223
synthesis, platinum(II) complexes and structural aspects of the new tetradentate phosphine *cis*,*trans*,*cis*-1,2,3,4-tetrakis(diphenylphosphino)cyclobutane, **290**, 167
synthesis, structure, and reactivity of new chiral ferrocenyl pyrazole ligands, **296**, 183
unusual coordination mode of a 2-pyridyl ketone oxime ligand in bis(4-butylphenyl 2-pyridyl ketone oximate)palladium(II), **285**, 138
water soluble platinum(II) and palladium(II) complexes of alkyl sulfonated phosphines, **290**, 189

Palladium compounds
bonding of the β-pinene allyl, η^3-$C_{10}H_{15}$ to Pd(II): X-ray structure of [Pd(η^3-$C_{10}H_{15}$)(Josiphos)]CF_3SO_3, **296**, 246
Palladium(I) complexes
palladium(I) phosphido-bridged dinuclear derivatives with both phosphine and amine ligands, **284**, 246
Paramagnetic complexes
redox active organoiron alkynyl centers with a coordination site: an approach toward molecular plugs. Complexation reactions with (THF)W(CO)$_5$ and $(PhCN)_2MCl_2$ (M = Pd, Pt), **291**, 403
Pentadienyl complexes
synthesis and characterization of siloxy-substituted diene and dienyl complexes of the Ru(C_5Me_5) fragment, **296**, 170
synthesis and characterization of the triethylphosphine adduct of bis(2,4-dimethylpentadienyl)tungsten, [W(2,4-C_7H_{11})$_2$-(PEt_3)], a complex with one η^5-**U** and one η^5-**S** pentadienyl ligand, **288**, 154
Pentanuclear complexes
a novel oligomerization versus metal chelation: synthesis and crystal structure of pentakis[di-μ-(diisopropylaminoethanethiolato)nickel], **294**, 83
Perchlorato-bridged complexes
synthesis, spectroscopic, magnetic properties and X-ray crystal structure of di-μ-hydroxo-bis(μ-perchlorato-*O,O'*)-bis-[bis(2-amino-4-methylpyrimidine)copper(II)]: bridging perchlorate results in significant deviation from the predicted magnetic exchange, **286**, 24
Phenanthroline complexes
Cu–O stretching frequency correlation with phenanthroline pK_a values in mixed copper complexes, **288**, 106
retention of optical activity during conversion of Λ-[Ru(1,10-phenanthroline)$_3$]$^{2+}$ to Λ-[Ru(1,10-phenanthroline-5,6-dione)$_3$]$^{2+}$ and Λ-[Ru(dipyrido[*a*:3,2-*h*:2',3'-*c*-]-phenazine)$_3$]$^{2+}$, **293**, 37
synthesis and electrochemical properties of a new C_{60}-phenanthroline derivative and its dimeric complex with silver, **292**, 34
synthesis and structure of a heptacoordinated cadmium(II) complex, **288**, 118
Phenol
β-silylethenylation of phenol with silylethyne using gallium trichloride, **296**, 67
Phenolate complexes
bisphenolate iron(II) complexes with intramolecularly coordinating nitrogen Lewis bases, **291**, 326
Phenoxide complexes
vanadium phenoxide complexes with oxide, nitride or hydrazide co-ligands: preparation and crystal structures of [V($OC_6H_3Pr^i_2$-2,6)$_3$NLi(C_4H_8O)$_3$], [{VO_2($OC_6H_3Pr^i_2$-2,6)$_2$}$_2$-{μ-Li(C_4H_8O)$_2$}$_2$] and [V(NNMe$_2$)($OC_6H_3Pr^i_2$-2,6)$_3$], **285**, 178
Phenyl bromide
Grignard reagent formation from aryl halides. There is no aryl radical intermediate along the dominant reaction channel, **296**, 52
Phenyl chloride
Grignard reagent formation from aryl halides. There is no aryl radical intermediate along the dominant reaction channel, **296**, 52
Phenyl iodide
Grignard reagent formation from aryl halides. There is no aryl radical intermediate along the dominant reaction channel, **296**, 52
Phosphaadamantane complexes
catalytic hydrogenation and deuteration of phospholipid model membranes with a water-soluble chlorotris(1,3,5-triaza-7-phosphaadamantane)rhodium(I) complex catalyst, **293**, 218
Phosphaalkene complexes
on the contrasting reactivity of the phosphaalkene [(η^5-C_5Me_5)(CO)$_2$FeP=C(NMe$_2$)$_2$] and the arsaalkene [(η^5-C_5Me_5)(CO)$_2$FeAs=C(NMe$_2$)$_2$] towards [(η^5-C_5H_5)Rh(CO)$_2$], **291**, 311
Phosphate complexes
magnetic iron phosphates with an open framework, **291**, 317
synthesis of the {Co[(6-amino-6-(4-aminobenzyl)-1,4,8,11-tetraazacyclotetradecane)][PO_4]}, a potential phosphate-binding site for molecular recognition, **293**, 245
Phosphate diester complexes
catalysis of hydrolysis of a phosphate diester by mono- and dinuclear macrocyclic zinc(II) complexes, **287**, 47
Phosphathia chelate ligand
platinum(II) complexes with the tridentate phosphino–thioether ligand bis(2-diphenylphosphinoethyl)sulfide, (PPh_2CH_2-$CH_2)_2S$. X-ray crystal structures of [Pt{(PPh_2CH_2-$CH_2)_2S$}Cl]ClO_4 and [Pt{($PPh_2CH_2CH_2)_2S$}I]I, **295**, 153
Phosphathiamacrocycle complexes
synthesis and coordination chemistry of 1-phenyl-1-phospha-4,7-dithiacyclononane, **294**, 170
Phosphide complexes
palladium(I) phosphido-bridged dinuclear derivatives with both phosphine and amine ligands, **284**, 246
Phosphide compound
electrochemical tuning of indium phosphide|poly(acetylene) interfaces, **294**, 207
Phosphine complexes
coordination chemistry and X-ray crystal structure of (ferrocenylmethyl)diphenylphosphine, $FcCH_2PPh_2$ [Fc = (η^5-C_5H_5)Fe(η^5-C_5H_4)], **295**, 18
electrospray mass spectrometric investigation of the relative ligand properties of EPh_3 ligands (E = P, As, Sb or Bi) towards Ag^+ and Cu^+, **290**, 213
NMR investigation of Pd(II)–Pd(0) reduction in the presence of mono- and ditertiary phosphines, **286**, 93
reaction of M(CO)Cl(L)$_2$ with mercury(II) chloride (M = Ir, Rh; L = Ph_2Ppy, P(fur)$_3$) (Ph_2Ppy = 2-(diphenylphosphino)pyridine, P(fur)$_3$ = tri-(2-furyl)phosphine), **284**, 119
syntheses and X-ray structures of mixed-metal gold phosphine clusters including an example having a highly asymmetric Re_2Au_2 skeleton, **294**, 255
synthesis and characterization of the triethylphosphine adduct of bis(2,4-dimethylpentadienyl)tungsten, [W(2,4-C_7H_{11})$_2$-(PEt_3)], a complex with one η^5-**U** and one η^5-**S** pentadienyl ligand, **288**, 154
synthetic approaches to Fe(II) complexes of the bulky phosphine P^tBu_2Me, **291**, 226
the reaction of triorganophosphorus diiodides, R_3PI_2, with zinc metal powder; dependency of product on R; the X-ray crystal structures of dimeric {ZnI_2[P(NMe$_2$)$_3$]}$_2$ and monomeric $ZnI_2(PPh_2Me)_2$, **292**, 213
Phosphine-substituted complexes
chiral phosphine ligands derived from sugars. 15. Synthesis, structure and spectroscopic studies of platinum(II) complexes with chiral phosphine substituted carbohydrate ligands, **290**, 121
Phosphinoalkyne complexes
reduction of cationic P-coordinated (diphenylphosphino)alkyne iron complexes with sodium borohydride, **291**, 207
Phosphino-imines
chemistry of polyfunctional molecules : CXXXI. Crystal structure and preparation of [bis(diphenyl–methyl–phosphino)iminium] tetraphenyloborate, [(C_6H_5)$_2$(CH_3)P=N=P-(CH_3)(C_6H_5)$_2$]$^+$[B(C_6H_5)$_4$]$^-$, **284**, 288

Phosphino-thioether complexes
^{13}C *ipso* chemical shifts in the aryl phosphino-thioether complexes PdX(*p*-RC_6H_4)(P,S). On resonance effects into the aryl ring, **290**, 95
Phosphite complexes
$[(NH_3)_5CoOP(H)(O)OCo(NH_3)_5]^{4+}$ and its conjugate acid; evidence for the P^+–O^- description of a terminal P–O bond, **294**, 14
Phospholipids
catalytic hydrogenation and deuteration of phospholipid model membranes with a water-soluble chlorotris(1,3,5-triaza-7-phosphaadamantane)rhodium(I) complex catalyst, **293**, 218
Phosphonate ligands
interaction of beryllium(II) in aqueous solution with bidentate ligands containing phosphonate groups, **285**, 39
Phosphorous dendrimers
carbosilane dendrimers functionalized with $AuFe_3$ clusters, **291**, 247
Phosphorous ligands
the preparation and coordination chemistry of $^iPr_2P(E)NHP(E')^iPr_2$ (E, E' = Se; E = Se, E' = S; E = S, E' = O; E,E' = O, **290**, 1
Phosphorus ligands
new versatile phosphorus-containing ligands—asymmetric $(XPR_2)(YPR'_2)NH$ (X, Y = O, S; R, R' = Ph, OEt) and their potassium salts: the crystal and molecular structures of $(SPPh_2)[OP(OEt)_2]NH$, $K[(SPPh_2)\{OP(OEt)_2\}N]\cdot H_2O$ and $(OPPh_2)[OP(OEt)_2]NH\cdot 1/2HCl\cdot 1/4H_2O$, **287**, 61
Photoacoustic calorimetry
volume and thermodynamic profiles of CO-binding to Fe(II) protoporphyrin IX in detergent micelles, **288**, 74
Photochemistry
photo-assisted preparation of double isocyanide insertion complexes of η^5-cyclopentadienyliron(II) bearing only isocyanide: the molecular structure of $(\eta^5\text{-}C_5H_5)$-$Fe(XylNC)_2[C(=NXyl)C(=NXyl)CH_3$ (XylNC = 2,6-$Me_2C_6H_3NC$), **286**, 233
syntheses, spectroscopic properties and photochemistry of [Re(R)$(CO)_3$(4,4'-Me_2-bpy)] (R = alkyl) complexes, **284**, 61
Photochromism
theoretical study of photochromic dinuclear molybdenum complexes, **295**, 180
Photooxidation
kinetics of the photooxidation of tris(bipyridine)iron(II) in chloroform, **291**, 136
photooxidation of water and isobutane in aqueous uranyl solutions, **285**, 322
Photophysical properties
synthesis and photophysical properties of *trans*-dithiocyanato bis(4,4'-dicarboxylic acid-2,2'-bipyridine) ruthenium(II) charge transfer sensitizer, **296**, 250
Photoreactions
photochemical reactions of transition metal organyl complexes with olefins: XXII. Photoreactions of pentacarbonyl(μ-$\eta^{5:3}$-4*H*-indenyl)diiron(Fe–Fe) with alkynes—substituent controlled C–H transfer or C–C bond formation, **296**, 127
Phthalocyanine complexes
a neutral molecule of I_2 as a bridge for dimerization of iodine iron(III) phthalocyanine, **287**, 101
indate(III) phthalocyanines: synthesis, spectroscopy and crystal structure of iodophthalocyaninato(2-) indate(III), **288**, 174
synthesis and characterization of a series of hexadecachlorinated phthalocyanines, **293**, 80
synthesis and X-ray characterization of phthalocyaninato(2-)-antimony(III) triiodide, **293**, 155
Picrate complexes
neodymium and europium picrate complexes with 3-picoline-*N*-oxide (3-picNO), **292**, 249
Picrolonate
novel structural aspects of oxythiamine, an antagonist of thiamine. Crystal structures of three salts, (oxythiamineH)-(picrolonate)$_2\cdot 2H_2O$, (oxythiamineH)$(PtCl_6)$, (oxythiamineH)$_2(V_{10}O_{28}H_2)\cdot 8H_2O$, and a metal complex Pt(oxythiamine)$Cl_3\cdot H_2O$, **295**, 71
Pincer complexes
synthesis and characterization of ruthenium 'pincer' complexes containing sulfur atom donor ligands. The X-ray crystal structure of $[RuCl_2(\eta^3\text{-}NN'N)(SMe_2)]$ (NN'N = {2,6-$(Me_2NCH_2)_2C_5H_3N$}), **287**, 113
β-Pinene allyl complexes
bonding of the β-pinene allyl, $\eta^3\text{-}C_{10}H_{15}$ to Pd(II): X-ray structure of $[Pd(\eta^3\text{-}C_{10}H_{15})(Josiphos)]CF_3SO_3$, **296**, 246
Planar chirality
diastereoselective *ortho*-lithiation of tricarbonylchromium complexes of chiral benzaldehyde acetals, **296**, 80
Platinum
an approach to *megalo*-boranes. Mixed and multiple cluster fusions involving iridaborane and platinaborane cluster compounds. Crystal structure determinations by conventional and synchrotron methods, **289**, 95
synthesis of chiral diphosphite ligands and comparison of their rhodium and platinum complexes in the asymmetric hydroformylation of styrene, **296**, 222
Platinum clusters
relativistic density functional study on the bimetallic cluster $[Pt_3Fe_3(CO)_{15}]^{n-}$ ($n = 0, 1, 2$), **286**, 30
Platinum complexes
a dinuclear platinum(II) complex with hydrazine and chloro bridges: $(NPr^n_4)[PtCl_2(\mu\text{-}Cl)(\mu\text{-}N_2H_4)PtCl_2]$, **284**, 103
chiral phosphine ligands derived from sugars. 15. Synthesis, structure and spectroscopic studies of platinum(II) complexes with chiral phosphine substituted carbohydrate ligands, **290**, 121
coordination geometries for palladium and platinum: theoretical studies and the synthesis and structure of tris(indazol-1-yl)borate complexes $MMe_3\{(ind)_3BH\}$, **287**, 27
di- and trinuclear Pd(II) and Pt(II) complexes containing bridging 2,7-disubstituted naphthyridines, **287**, 72
different chlorination modes of oximes: chlorination of salicylaldoxime coordinated to platinum, **285**, 116
facile synthesis and structural properties of (diamine)tetracarboxylatoplatinum(IV) complexes, **292**, 52
formation of two stable seven-membered chelate rings bound to one metal center due to addition of bifunctional amine to coordinated nitriles, **285**, 203
iminoacylation: 2. Addition of alkylated hydroxylamines via oxygen to platinum(IV)-bound nitriles, **292**, 272
interactions between thiamine and large anions. Crystal structures of (H-thiamine)$[Pt^{II}(SCN)_4]\cdot 3H_2O$ and (H-thiamine)$[Pt^{IV}(SCN)_6]\cdot H_2O$, **290**, 145
[*meso*- and *rac*-1,2-bis(4-fluorophenyl)ethylenediamine]chloro-[sulfinylbis(methane)-S]platinum(II) chloride new water soluble platinum complexes with high anti-breast cancer activities, **285**, 184
multinuclear NMR study and crystal structures of complexes of the types *cis*- and *trans*-$Pt(Ypy)_2X_2$, where Ypy = pyridine derivative and X = Cl and I, **295**, 25
phase dependent η^2-alkene versus η^1-alkyl transformation in platinum(II) complexes, **285**, 301
platina- and palladalactam complexes derived from 2-benzoylacetanilide; syntheses and X-ray structure of [Pd{NPhC-(O)CHC(O)Ph}(bipy)]$\cdot CH_2Cl_2$, **292**, 260
platinum(II) and palladium(II) complexes with 3,3,8,8-tetramethyl-1,6-dioxa-4,9-diaza-5λ^5-phosphaspiro[4,4]nonane, **295**, 164

platinum(II) and palladium(II) saccharinate complexes, **285**, 145
preparation spectroscopy and conductivity of [Pt(SS)(NN)]-type complexes and their iodine oxidized complexes, **284**, 112
properties of Pt(II) complexes containing both a pyridyl N–N chelate ligand having a long alkyl chain and a sulfur-rich dithiolate ligand and their molecular interactions in the solid state, **284**, 55
'ship-in-bottle' synthesis of platinum carbonyl clusters in titania- or zirconia-modified mesoporous channels of FSM-16. Structural characterization and catalysis in CO hydrogenation, **294**, 281
structural and spectroscopic characterisation of the mixed valence platinum complex $PtdabI_3$ (dab = *o*-diaminobenzene), **287**, 218
structures of 1,3-bis(diphenylphosphino)propane platinum(II) alditolate complexes, **285**, 10
studies on Pt–S bonds. Complexes of methylsulfanyl- and methylsulfynil-acetate, benzoate and phenolate with the diethylenetriamineplatinum(II) moiety, **285**, 249
synthesis and biological activity of *cis*-dichloro mono- and bis-(platinum) complexes with *N*-alkyl-ethylenediamine ligands, **290**, 237
synthesis and characterisation of mono- and binuclear palladium and platinum complexes with organochalcogenides, **285**, 190
synthesis and properties of (diamine)platinum(II) complexes of pyridine carboxylate isomers and their antitumor activity, **292**, 238
synthesis and structures of (*SP*-4-2)-diiodobis(2-hydroxyethylamine)platinum(II), (*SP*-4-2)-dichlorobis(2-hydroxyethylamine)platinum(II) and (*OC*-6-22)-bis(2-hydroxyethylamine)tetrachloroplatinum(IV) in the crystal, **292**, 127
synthesis of five-coordinate Pt(II) complexes containing organogermanium fragments in axial position. Structure of $[PtCl(GePh_2Cl)(2,9\text{-}Me_2\text{-}1,10\text{-}phen)(CH_2{=}CH_2)]$, **285**, 70
synthesis of organo(siloxo)platinum and -palladium complexes and preparation of supported nanoclusters by facile ligand reduction, **294**, 266
synthesis of transition metal isocyanide complexes containing hydrogen bonding sites in peripheral locations, **285**, 223
synthesis, platinum(II) complexes and structural aspects of the new tetradentate phosphine *cis,trans,cis*-1,2,3,4-tetrakis-(diphenylphosphino)cyclobutane, **290**, 167
synthetic control of intrinsic ground-state defects in a mixed valence quasi-one-dimensional Pt halide chain, **294**, 275
water soluble platinum(II) and palladium(II) complexes of alkyl sulfonated phosphines, **290**, 189

Platinum(II) complexes
synthesis, structures, and linkage isomerism of (allylbenzylmalonate)platinum(II) complexes, **295**, 214

Platinum(II) phosphine complexes
platinum(II) complexes with the tridentate phosphino–thioether ligand bis(2-diphenylphosphinoethyl)sulfide, $(PPh_2CH_2CH_2)_2S$. X-ray crystal structures of $[Pt\{(PPh_2CH_2CH_2)_2S\}Cl]ClO_4$ and $[Pt\{(PPh_2CH_2CH_2)_2S\}I]I$, **295**, 153

PM3 calculations
a semi-empirical quantum chemical study (PM3) of structure–reactivity relationships for mono- and binuclear bismuthines, **284**, 167

P,N-ligand complexes
platinum(II) and palladium(II) complexes with 3,3,8,8-tetramethyl-1,6-dioxa-4,9-diaza-5λ^5-phosphaspiro[4,4]nonane, **295**, 164

Polyamine complexes
anionopentaaminecobalt(III) complexes with polyamine ligands: Part 30. The base hydrolysis kinetics and structure determination of some $[CoCl(N)_5]^{2+}$ complexes containing pyridine ligands, **288**, 181
complexation of thorium(IV) by tris((2,3-dihydroxybenzylamino)ethyl)amine—a new strong chelating agent, **286**, 103
synthesis, structure and magnetic behavior of a di-μ-chloro-di-μ-imidazolato-tetracopper(II) complex with two dinucleating macrocycles, **287**, 52

Polydentate amine complexes
the methylcobalt(III) complex of a tetrapodal pentadentate amine ligand, 2,6-bis(1′,3′-diamino-2′-methyl-prop-2′-yl)-pyridine, **286**, 98

Polydentate ligand complexes
a reversible NO complex of $[Fe^{II}(edampda)]$ and the $[Ni^{II}(edampda)]$ analogue (edampda^{2-} = *N,N′*-bis(pyridylmethyl)ethylenediamine-*N,N′*-diacetate), **286**, 197

Polydentate phosphine complexes
synthesis and structural characterization of binuclear silver(I) complexes bridged by three polydentate phosphine ligands, **284**, 278

Polyiodide chains
polyiodides of transition metal trischelate cations: syntheses, structures, and spectral and electrical conductivity studies of $[Mn(phen)_3](I_3)_2$ and $[Mn(bpy)_3](I_3)_{1.5}(I_8)_{0.25}$, **284**, 158

Polymer compound
electrochemical tuning of indium phosphide|poly(acetylene) interfaces, **294**, 207

Polymeric complexes
a new self-assembled copper(I) complex of 2-thioxo-5H-1,3-dithiolo[4,5-*b*][1,4]dithiepin-6(7*H*)-one showing three-dimensional supramolecular networks, **292**, 117
a novel polymeric disilver(I) complex containing both macrocyclic- and linear-coordinate silver(I) centres: $[Ag_2(mn\text{-}15S_2O_3)]_n(ClO_4)_{2n}$ (mn-15S_2O_3 = maleonitrile-dithia-[15]crown-5), **285**, 305
coordination compounds with 4-aminopyridine. An unusual in situ formation of a bridging formato compound. The structure of *catena*-poly[bis(4-amino-pyridine)(formato-*O*)-copper(II)-μ-formato(*O,O′*)], **287**, 226
synthesis, characterization, spectroscopy, magnetism and X-ray structures of red and green Cu(II) chloride adducts with bis(2-benzimidazolyl)propane, **288**, 220

Polymerization
cleavage of substituents upon oxidative polymerization of substituted thiophenes, **287**, 232

Polymers
manufacturing of specialty materials in supercritical fluid carbon dioxide, **294**, 214
Rh-complex promoted synthesis of poly(arylacetylene) with pendant TTF units and formation of CT complexes of the polymer with π-acceptors, **296**, 278

Polynuclear complexes
di- and trinuclear Pd(II) and Pt(II) complexes containing bridging 2,7-disubstituted naphthyridines, **287**, 72
reaction patterns of 'tetraferrio-azaallenium', $[(OC)_3Cp_2Fe_2(\mu_4\text{-}C{=}N{=}C)Fe_2Cp_2(CO)_3]^+$, and its 'activated cyanide'-precursor (μ-phthalimidocarbyne)tricarbonyldicyclopentadienyldiiron(1+), **291**, 266
redox active organoiron alkynyl centers with a coordination site: an approach toward molecular plugs. Complexation reactions with $(THF)W(CO)_5$ and $(PhCN)_2MCl_2$ (M = Pd, Pt), **291**, 403
structure and magnetic properties of dinuclear [Mn(III)-(salen)$(H_2O)]_2(ClO_4)_2$ and polynuclear [Mn(III)(salen)-$(NO_3)]_n$, **290**, 8
synthesis and characterization of a series of new thiocarboxylate complexes of molybdenum(V), **284**, 223
synthesis, crystal structure and magnetic properties of *catena*-poly[(2,2′-bipyridylcopper)-di-μ-thiocyanato] and bis(2,2′-bipyridyl)thiocyanatocopper(II) perchlorate, **286**, 169

synthesis, crystal structures and magnetic properties of trinuclear oxo-centered manganese complexes of the general formula $[Mn_3O(O_2CR)_6L_3]^{z+}$ (R = Me, Et, Ph, CH_2Cl; L = 3-methylpyridine or water $z = 1, 0$), **287**, 82

synthesis, structure and magnetic behavior of a di-μ-chloro-di-μ-imidazolato-tetracopper(II) complex with two dinucleating macrocycles, **287**, 52

Polynuclear compounds

crystal structures and magnetic properties of the squarate-O^1,O^n-bridged dinuclear copper(II) complexes $[Cu_2(phen)_4(C_4O_4)](CF_3SO_3)_2 \cdot 3H_2O$ ($n = 2$) and $[Cu_2(bipy)_4(C_4O_4)](CF_3SO_3)_2 \cdot 6H_2O$ ($n = 3$), **287**, 173

Polyoxometalate complexes

a polyoxoanion-based iron(II) catechol extradiol dioxygenase: oxygenation of 3,5-di-tert-butylcatechol by the polyoxoanion-supported iron(II) complex $(n\text{-}Bu_4N)_7[(CH_3CN)_3\text{-}Fe^{II}\cdot P_2W_{15}Nb_3O_{62}]$, **291**, 426

Poly(pyrazolyl)borate complexes

synthesis and structural characterization of face-sharing bioctahedral complexes containing poly(pyrazolyl)borate ligands: $[HB(Me_2Pz)_3BH][X_3Mo(\mu\text{-}X)_2(\mu\text{-}H)MoTp^*]$ (X = Cl or Br; $Tp^* = HB(Me_2Pz)_3$; Pz = pyrazolyl), **292**, 182

Polysilacycloalkanes

a facile synthesis of 1,3-disilacyclohexanes, **296**, 231

Porphinatoiron(II) derivative

5,10,15,20 - tetrakis(α,α,α,α - *o* - pivalamidophenyl)porphinatoiron(II) bearing a covalently linked axial imidazole via *m*-aminobenzoic acid. Synthesis and influence of imidazole basicity on O_2-binding affinity, **295**, 234

Porphyrin aggregates

spectroscopic characterization of porphyrin supramolecular aggregates on poly-lysine and their application to quantitative DNA determination, **286**, 121

Porphyrin complexes

conformational diversity in (octaethylporphinato)(trichloroacetato)iron(III) derivatives, **291**, 49

heterobimetallic nitrido-bridged Ru(II)NOs(VIII) and Ru(II)-NOs(VI) complexes containing ruthenium porphyrins, **290**, 28

structure and magnetic properties of (μ-oxo)bis[*meso*-tetrakis(*p*-tolyl-porphyrinato)iron(III)] and (μ-oxo)bis[*N,N'*-ethylene-bis(2-acetylphenoliminato)iron(III)], **290**, 51

visible and resonance Raman spectra of low valent iron porphyrins, **287**, 21

Porphyrins

hydrothermal synthesis of microporous materials with the direct incorporation of porphyrin molecules, **294**, 179

Potassium compounds

syntheses and structures of Group 1 mellitate compounds, $K_6[C_6(COO)_6] \cdot 8H_2O$ and $Cs_5[C_6(COO)_6H] \cdot 7H_2O$, **286**, 207

Proline complexes

an EXAFS study of Mo(VI) and Mo(V) complexes with proline dithiocarbamate, **288**, 127

Propargyl complexes

cross-coupling reactions proceeding through η^1- and η^3-propargyl/allenyl–palladium(II) intermediates, **296**, 37

Propene

palladium(II) complexes with hemilabile etherdiphos ligands in the alternating copolymerization of carbon monoxide with olefins, **296**, 103

2-Propynylamine

synthesis of 2-silylmethyl-2-alkenals by rhodium-catalyzed interaction among 2-propynylamine, hydrosilane, and CO, **296**, 72

Protonation

alkyl derivatives of $[B_6H_6]^{2-}$: NMR and vibrational spectra and crystal structure of $(Ph_4P)[B_6H_6CH_2Ph]$, **289**, 70

protonation and substitution reactions of Fe–S 'basket' clusters including extracted FeMo-cofactor of nitrogenase, **291**, 212

Pseudohalogens

the preparation and characterisation of a new interpseudohalogen, $ClCS_2N_3$, and $(CS_2N_3)SS(CS_2N_3)$, a novel disulfur-dipseudohalide, **294**, 68

$PtCl_3^-$

novel structural aspects of oxythiamine, an antagonist of thiamine. Crystal structures of three salts, (oxythiamineH)-(picrolonate)$_2 \cdot 2H_2O$, (oxythiamineH)$(PtCl_6)$, (oxythiamineH)$_2(V_{10}O_{28}H_2) \cdot 8H_2O$, and a metal complex Pt(oxythiamine)$Cl_3 \cdot H_2O$, **295**, 71

Pyrazine complexes

a large C_6D_6 solvent effect on the ^1H NMR shifts of pyrazines coordinated to the $M(CO)_5$ series (M = W, Mo, Cr), **285**, 157

Pyrazole

crystal structures of two new heptamolybdates and of a pyrazole incorporating a γ-octamolybdate anion, **295**, 106

Pyrazolylalkane complexes

new palladium(II) complexes with bis(pyrazol-1-yl)alkanes, **295**, 136

Pyrazolyl ligands

synthesis, structure, and reactivity of new chiral ferrocenyl pyrazole ligands, **296**, 183

Pyridine–carbaldimine complexes

properties of Pt(II) complexes containing both a pyridyl N–N chelate ligand having a long alkyl chain and a sulfur-rich dithiolate ligand and their molecular interactions in the solid state, **284**, 55

Pyridine complexes

coordination compounds with 4-aminopyridine. An unusual in situ formation of a bridging formato compound. The structure of *catena*-poly[bis(4-amino-pyridine)(formato-*O*)-copper(II)-μ-formato(*O,O'*)], **287**, 226

multinuclear NMR study and crystal structures of complexes of the types *cis*- and *trans*-$Pt(Ypy)_2X_2$, where Ypy = pyridine derivative and X = Cl and I, **295**, 25

new synthesis of $[RuH(CO)(PPh_3)_2L_2][BF_4]$ [L_2 = 2py, 2(4-Mepy) or bpy] and X-ray crystal structure of $[RuH(CO)(PPh_3)_2(4\text{-}Mepy)_2][BF_4]$, **285**, 318

synthesis and dynamic NMR studies of stereochemical non-rigidity in rhenium(I) complexes of 2,6-bis[1-(phenylimino)-ethyl]pyridine derivatives, **295**, 56

Pyridine ligands

X-ray crystal structures of copper(II) nitrate complexes of two isomeric N,O-chelating ligands, **284**, 273

Pyridines

pyridines from allyl cobaloximes: a new decomposition pathway, **288**, 112

Pyridine-2-selenolato complexes

electrochemical synthesis of copper and silver pyridine-2-selenolato complexes: crystal structure of (6-trimethylsilylpyridine-2-selenolato)silver(I), $[Ag_6(6\text{-}Me_3SipySe)_6]$, **284**, 133

Pyridyl-amine complexes

dimolybdenum complexes of bis(2-pyridyl)amine (Hdpa) ligand and its anion (dpa^-); transformation of chelating η^2-Hdpa to bridging η^2-dpa^-, **288**, 82

Pyridylamine complexes

complexes of copper(II) dihalide with 2,2'-dipyridylamine. X-ray diffraction structures of the [dibromo-bis(dipyam) copper(II)]–water (1:2) and di[chloro-bis(dipyam)-copper(II)]diiodide–acetonitrile (1:2) complexes, **290**, 180

Pyridyl complexes

solvatochromism in ligand-field absorption bands and solvent-dependent deuteron nuclear magnetic resonance spectra of $[CrF_3(tri(2\text{-}pyridyl)amine)]$: implication of trigonal splitting for the first spin-allowed d–d band and evaluation of π bonding peculiarities, **295**, 84

Pyridyl-pyrimidine complexes
synthesis, physical and photophysical properties of Ru(II) pyridyl-pyrimidine and pyridyl-pyrimidinone complexes: effect of substituents on the pyrimidine ring, **296**, 114
Pyridyl-pyrimidinone complexes
synthesis, physical and photophysical properties of Ru(II) pyridyl-pyrimidine and pyridyl-pyrimidinone complexes: effect of substituents on the pyrimidine ring, **296**, 114
Pyrimidine complexes
absence of 1,3-metallotropic shifts in $M(CO)_5$ complexes of pyrimidine and 1,3,5-triazine (M = W, Mo, Cr), **292**, 225
Pyrimidine ring
synthesis, physical and photophysical properties of Ru(II) pyridyl-pyrimidine and pyridyl-pyrimidinone complexes: effect of substituents on the pyrimidine ring, **296**, 114

Quadruply-bonded complexes
two diphenylformamidinate compounds of the quadruply-bonded Re_2^{6+} core, **285**, 149
Quadrupole splitting
Mössbauer studies of different types of hematin, **286**, 229
strained, ring tilted dicarbon-bridged [2]ferrocenophanes and ferrocene revisited: ^{57}Fe Mössbauer spectroscopic study of bonding, hyperfine interactions, and lattice dynamics, **287**, 152
Quasi-one-dimensional chain
synthetic control of intrinsic ground-state defects in a mixed valence quasi-one-dimensional Pt halide chain, **294**, 275
Quinone complexes
coordination of *ortho* and *para* quinones and oxidation reduction processes in trimethylphosphine complexes of cobalt and nickel, **287**, 167

Raman spectra
lattice vibrations of the icosahedral solid boron arsenide, $B_{12}As_2$, **289**, 198
Raman spectroscopy
vibrational spectra of the series CdX_2(18-crown-6) and CdXY(18-crown-6) (X or Y = Cl, Br or I), **292**, 108
Rare earth complexes
synthesis and crystal structure of two new rare earth trimesate complexes: $ErTMA(H_2O)_5{\cdot}3.5H_2O$ and $YTMA(H_2O)_6$, **284**, 139
Rare earth halides
B and B–C as interstitials in reduced rare earth halides, **289**, 174
Reduction
kinetics and mechanism of the reduction of ferrate by one-electron reductants, **293**, 229
NMR investigation of Pd(II)–Pd(0) reduction in the presence of mono- and ditertiary phosphines, **286**, 93
Regioselectivity
copper-mediated 1,6-, 1,8-, 1,10- and 1,12-addition and 1,5-substitution reactions in organic synthesis, **296**, 1
$[Rh(\mu\text{-}Cl)(\eta^4\text{-norbornadiene})]_2$
Rh-complex promoted synthesis of poly(arylacetylene) with pendant TTF units and formation of CT complexes of the polymer with π-acceptors, **296**, 278
Rhenium carboxylates
synthesis and crystal structure of $[Re(CO)_3(\mu\text{-}CH_3CO_2)(THF)]_4$, **284**, 304
Rhenium complexes
coordination mode of dihydroxybenzoic acids in oxorhenium(V) complexes, **294**, 114
deboronation of $HB(3,5\text{-}Me_2C_3HN_2)_3^-$ in a reaction with $[Re(O)Cl_4]^-$, the X-ray crystal structures of $[\{Re(O)Cl(3,5\text{-}Me_2C_3HN_2H)_3\}(\mu\text{-}O)\{Re(O)Cl_3(3,5\text{-}Me_2C_3HN_2H)\}]$ and $[\{Re(O)Cl_2(3,5\text{-}Me_2C_3HN_2H)_2\}(\mu\text{-}O)\{Re(O)Cl_2(3,5\text{-}Me_2C_3HN_2H)_2\}]$, **285**, 55
electronic spectra of 1,2-diiminetricarbonylrhenium(I)chloride complexes with imidazole derivatives as ligands, **288**, 150
five-coordinate rhenium(III)-thiolato complexes: the structure of $[Re(SCH_2PhOCH_3)_3(PPh_3)_2]$, **290**, 247
organometallic radiopharmaceuticals: rhenium(I) carbonyl complexes of natural bile acids and derivatives, **286**, 46
reactions of H_2, silanes, and olefins with superelectrophilic cationic rhenium complexes: heterolytic cleavage of H_2 and relation to the structure and function of hydrogenases, **294**, 240
study on the formation of mixed ligand oxorhenium and oxotechnetium complexes (SNS/S combination), **295**, 1
syntheses, spectroscopic properties and photochemistry of $[Re(R)(CO)_3(4,4'\text{-}Me_2\text{-bpy})]$ (R = alkyl) complexes, **284**, 61
synthesis and characterization of five-coordinate rhenium(V) and technetium(V) mixed ligand bifunctional complexes carrying the SNS/S or the SNN/S donor atom set. Crystal structure of $ReO\{[(C_2H_5)_2NCH_2CH_2N(CH_2CH_2S)_2](p\text{-}H_2N\text{–}PhS)\}$ and $ReO\{[(CH_2)_4NCH_2CH_2NCH_2CH_2S](p\text{-}H_2N\text{–}PhS)\}$, **285**, 97
synthesis and characterization of oxorhenium-'3 + 1' mixed-thiolate complexes. Crystal and molecular structures of $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}(C_6H_4X\text{-}4\text{-}CH_2S)]$ (X = F, Cl, Br, OMe) and of the pendant thiolate compounds $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}(\eta^1\text{-}SCH_2CH_2SCH_2CH_2SH)]$ and $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}\{\eta^1\text{-}SCH_2CH(OH)CH(OH)CH_2SH\}]$, **284**, 252
synthesis and crystal structure of $[Re(CO)_3(\mu\text{-}CH_3CO_2)(THF)]_4$, **284**, 304
synthesis and dynamic NMR studies of stereochemical non-rigidity in rhenium(I) complexes of 2,6-bis[1-(phenylimino)ethyl]pyridine derivatives, **295**, 56
synthesis and structural characterisation of rhenium(V) oxo-thioether thiolate complexes with pendant ester or carboxylate groups, **285**, 76
synthesis, characterization and crystal structures of rhenium(V) complexes with diphosphines, **294**, 47
two diphenylformamidinate compounds of the quadruply-bonded Re_2^{6+} core, **285**, 149
Rhenium–oxo system
synthesis, molecular and crystal structure of the zwitterionic Re(V)–oxo complex $ReOCl_3(L)(PPh_3)$ with L = 2-(diethylaminomethyl)-4-methylphenol, **295**, 209
Rhodium
synthesis of chiral diphosphite ligands and comparison of their rhodium and platinum complexes in the asymmetric hydroformylation of styrene, **296**, 222
Rhodium catalyst
rhodium-catalyzed cyclic carbonylation of 2-phenylethynylbenzamide under water–gas shift reaction conditions, **296**, 195
Rhodium-catalyzed silylformylation
synthesis of 2-silylmethyl-2-alkenals by rhodium-catalyzed interaction among 2-propynylamine, hydrosilane, and CO, **296**, 72
Rhodium complexes
catalytic hydrogenation and deuteration of phospholipid model membranes with a water-soluble chlorotris(1,3,5-triaza-7-phosphaadamantane)rhodium(I) complex catalyst, **293**, 218
diphosphinito-bridged ruthenium(II) and rhodium(III) complexes with stereogenic metal centers, **292**, 84
halfsandwich complexes of ruthenium(II), rhodium(III) and iridium(III) with *N*-substituted 3-hydroxy-2-methyl-4-pyridone ligands, **294**, 62
reaction of $M(CO)Cl(L)_2$ with mercury(II) chloride (M = Ir, Rh; L = Ph_2Ppy, $P(fur)_3$) (Ph_2Ppy = 2-(diphenylphosphino)pyridine, $P(fur)_3$ = tri-(2-furyl)phosphine), **284**, 119

structural, spectroscopic and catalytic properties of water-soluble hydride rhodium complexes $[RhH(Rtpa^{+}I^{-})_4]H_2O$ (R = Me, Et), **293**, 110
synthesis, reactivity and catalytic properties of rhodium complexes of (*R,R*)-1-benzyl-3,4-dithioetherpyrrolidines, **295**, 64
synthetic routes for hydrotris(pyrazolyl)borate complexes of rhodium(III), **284**, 191
Ring closure
a facile synthesis of 1,3-disilacyclohexanes, **296**, 231
Ring-opening polymerization
ring-opening polymerization of norbornene by η-cycloheptatrienyl molybdenum and tungsten complexes, **288**, 226
Ring strain
strained, ring tilted dicarbon-bridged [2]ferrocenophanes and ferrocene revisited: ^{57}Fe Mössbauer spectroscopic study of bonding, hyperfine interactions, and lattice dynamics, **287**, 152
Rotation barriers
early versus late transition metals. Electronic structure of *nido*-2-$CpML_nB_4H_8$, $CpML_n$ = $CpTaCl_2$, $CpWH_3$ and CpCo, **289**, 85
Rubredoxin models
effect of N–H···S amide–thiolate hydrogen bonding on bond lengths in rubredoxin models $[(CH_3)_3NCH_2CONH_2]_2$-$[M(S_2$-*o*-xyl$)_2]$, **291**, 388
Ruthenium
preparation and characterization of various *N*-substituted-2-aza-[3]-ferrocenophanes and their chemical and electrochemical properties, **296**, 176
Ruthenium chemistry
[4-{(*p*-cym)Ru}-5-{$(Ph_3P)_2(CO)Os\}B_8H_8$]: the first *nido*-heterobimetallahexaborane cluster, **289**, 134
Ruthenium clusters
the versatile nature of [ClAu(L–L)AuCl] (L–L = bis(diphenylphosphino)methane (dppm) or -butane (dppb)) in reactions with $[Rh_2Ru_4(CO)_{16}B]^-$: crystal structures of $[\{Rh_2Ru_4(CO)_{16}B\}_2\{\mu$-Au(dppb)Au$\}]$ and $[RhRu_4(CO)_{14}B\{$Au(dppm)Au$\}]$, **289**, 149
Ruthenium complexes
a simple route to bis(terpyridyl)transition metal oligomers, **284**, 76
bdmpp complexes of Ru(II) as structural analogs for Ru(II)-2,2′:6′,2″-terpyridine species (where bdmpp is 2,6-bis(3,5-dimethyl-*N*-pyrazoyl)pyridine), **286**, 181
cationic 16-electron half-sandwich ruthenium complexes containing asymmetric diamines: understanding the stability and reactivity of coordinatively unsaturated two-legged piano stool complexes, **286**, 114
^{13}C NMR confirmation that $[Ru^{II}$(hedta)(6-azauridine)$]^-$ has η^2(5,6) and not σ(C-5) coordination, **293**, 123
comparing structures and reactivity in analogous Fe and Ru complexes. (^{i}Pr-DAB)$Fe(CO)_2I_2$ and (^{i}Pr-DAB)FeI_2: a perfectly reversible CO-carrier system. (R-DAB = *N,N′*-R_2-1,4-diaza-1,3-butadiene), **291**, 438
coordination chemistry and X-ray crystal structure of (ferrocenylmethyl)diphenylphosphine, $FcCH_2PPh_2$ [Fc = (η^5-C_5H_5)Fe(η^5-C_5H_4)], **295**, 18
detection of N-3 and N-7-coordinated $[Ru^{II}$(edta)(5′-GMP)$]^{4-}$ complexes and the N-1 protonation equilibrium of the Ru^{III} derivative, **285**, 170
dinuclear complexes of a new bridging ligand containing 2,2′-bipyridyl and dioxolene binding sites: syntheses, electrochemical and electronic spectroscopic properties, **285**, 89
diphosphinito-bridged ruthenium(II) and rhodium(III) complexes with stereogenic metal centers, **292**, 84
evidence for a charge-transfer intermediate in the catalytic epoxidation of alkenes by ruthenium complexes: a comparison between electrochemical and kinetic data, **288**, 122
evidence for a new nitrosyl-to-nitro reaction and an acetamide formation in the $\{RuNO\}^6$-type complex, *cis*-$[Ru(NO)(CH_3CN)(bpy)_2]^{3+}$, **290**, 113
halfsandwich complexes of ruthenium(II), rhodium(III) and iridium(III) with *N*-substituted 3-hydroxy-2-methyl-4-pyridone ligands, **294**, 62
heterobimetallic nitrido-bridged Ru(II)NOs(VIII) and Ru(II)-NOs(VI) complexes containing ruthenium porphyrins, **290**, 28
intervalence transfer properties of the binuclear μ-benzotriazolate- and μ-benzimidazolate-bis{ruthenium(II)/(III)-edta} complexes, **285**, 197
iron and ruthenium organometallic derivatives of spiro-substituted cyclotriphosphazenes bearing nitrile spacer: synthesis and electronic structure, **294**, 20
new synthesis of $[RuH(CO)(PPh_3)_2L_2][BF_4]$ [L_2 = 2py, 2(4-Mepy) or bpy] and X-ray crystal structure of $[RuH(CO)(PPh_3)_2$(4-Mepy$)_2][BF_4]$, **285**, 318
reaction of hydrous nitrosylruthenium trichloride with 2-pyridinemethanol, **290**, 36
retention of optical activity during conversion of Λ-[Ru(1,10-phenanthroline$)_3]^{2+}$ to Λ-[Ru(1,10-phenanthroline-5,6-dione$)_3]^{2+}$ and Λ-[Ru(dipyrido[*a*:3,2-*h*:2′,3′-*c*]-phenazine$)_3]^{2+}$, **293**, 37
some new chain-like terdentate phosphines, their ruthenium(II) coordination chemistry and the activity of the cations $[Ru(MeCN)_3(PhP\{CH_2CH_2P(p$-X-$C_6H_4)_2\}_2)]^{2+}$ (X = H, F, Me and OMe) as acetalization catalysts, **290**, 64
synthesis and characterization of a metal coordination polymer consisting of ruthenium(III) β-diketone units linked by butadiyne bridges, **292**, 64
synthesis and characterization of ruthenium 'pincer' complexes containing sulfur atom donor ligands. The X-ray crystal structure of $[RuCl_2(\eta^3$-NN′N)$(SMe_2)]$ (NN′N = {2,6-$(Me_2NCH_2)_2C_5H_3N\}$), **287**, 113
synthesis and characterization of siloxy-substituted diene and dienyl complexes of the Ru(C_5Me_5) fragment, **296**, 170
synthesis, characterization and optical properties of novel carbazole–ruthenium compounds: carbazole as a new building block?, **288**, 134
the ruthenium and iridium coordination chemistry of the tripodal ligand $CH_3C\{CH_2P(m$-CF_3-$C_6H_4)_2\}_3$, **293**, 70
Ruthenium(II) complexes
synthesis and photophysical properties of *trans*-dithiocyanato bis(4,4′-dicarboxylic acid-2,2′-bipyridine) ruthenium(II) charge transfer sensitizer, **296**, 250
Ru(II) complexes
synthesis, physical and photophysical properties of Ru(II) pyridyl-pyrimidine and pyridyl-pyrimidinone complexes: effect of substituents on the pyrimidine ring, **296**, 114

Saccharin complexes
platinum(II) and palladium(II) saccharinate complexes, **285**, 145
Salicylalodiminate complexes
structure and DNA binding properties of the gold(III) complex $[AuCl_2$(esal)], **285**, 309
Salicylate complexes
some complexes derived from zinc salicylate or 3,5-di-tert-butylsalicylate. The crystal structure of (2,2′-bipyridyl)-(methanol)(*O*-salicylato)(*O,O′*-salicylato)zinc, **287**, 89
Sandwich complexes
organometallic electron reservoir sandwich iron complexes as potential agents for redox and electron transfer chain catalysis, **291**, 1
the bonding and structure of B–N sandwich complexes, **288**, 17

Sarcosine derivative complexes
a series of heteroleptic complexes of the type *fac*-$[Mn^{III}L_2]^-$ [H_2L = derivatives of *N*-(2-hydroxybenzyl)glycine or *N*-(5-nitro-2-hydroxybenzyl)sarcosine] possessing unusual Mn(III) co-ordination spheres, **290**, 228
Scandium complexes
high coordination numbers of metal ions in chelate complexes: molecular adducts of scandium tris-α-diketonates in solution, **286**, 175
solution state chemistry of scandium(III): a study of scandium-crown ether complexes by ^{45}Sc-NMR spectroscopy, **284**, 71
Schiff base complexes
a new linear polymer chain iron complex; synthesis, crystal and molecular structural characterization of catena-μ-acetato-[*NN*′-ethylenebis(salicylaldiminato)]iron(III), **287**, 109
1:1 adducts of triphenyltin chloride with oxovanadium(IV) tetradentate Schiff-base complexes, **293**, 147
an optically active nickel(II) Schiff base coordination compound *N*,*N*′-(1*R*,2*R*)-(−)-1,2-cyclohexylenebis(2-hydroxyacetophenonylideneiminato)nickel(II), **293**, 239
facile and stereoselective condensation of acetone with ammonia ligands on cobalt(III): structure of a N-bonded cyanate complex containing the 2-methyl-2-amino-4-imino-pentane ligand, **290**, 133
formation pathway of a Mn(IV),Mn(IV) bis(μ-oxo) dimer that incorporates alkali and alkaline earth cations and electron transfer properties of the dimer, **294**, 133
low-spin iron(III) complexes with N,S coordination: syntheses, structures, and properties of bis(*N*-2-mercaptophenyl-2′-pyridylmethyleniminato)iron(III) tetraphenylborate and bis(*N*-2-mercapto-2-methylpropyl-2′-pyridylmethyleniminato)iron(III) tetraphenylborate, **285**, 269
macrocyclic complexes derived from the cyclocondensation of sodium 2,6-diformyl-4-substituted phenolates with *N*,*N*-bis(2-aminoethyl)-2-hydroxybenzylamine and in situ transmetallation. An anticipated dinuclear Cd(II) pendant-arm macrocyclic complex and an unexpected mononuclear Fe(III) macrocyclic complex, **294**, 1
nickel complexes of tetradentate asymmetric ligands, **290**, 14
preparation and characterisation of new oxovanadium(IV) Schiff base complexes derived from amino acids and aromatic *o*-hydroxyaldehydes, **293**, 1
structure and magnetic properties of dinuclear [Mn(III)-(salen)$(H_2O)]_2(ClO_4)_2$ and polynuclear [Mn(III)(salen)-$(NO_3)]_n$, **290**, 8
studies of the binding modes of carboxylate donors with manganese(III) complexes containing tetradentate Schiff base ligands. Crystal structures of the complexes [Mn(3-CH_3Osalpn)$(HO_2CC_6H_4CO_2)]_n$ and [{Mn(5-NO_2salpn)-(OH) (H_2O)} {Mn(5-NO_2salpn) $(HO_2CC_6H_4CO_2)$ (H_2O)}]·C_2H_5OH, **293**, 210
synthesis of the ligand *N*,*N*′-bis(2-tert-butylthiobenzenylidene)-diethylenetriamine; its reactivity with nickel(II) salts, **287**, 105
synthesis, X-ray structural determination and Mössbauer characterization of Schiff bases bearing ferrocene groups, their reduced analogues and related complexes, **287**, 117
synthetic, spectroscopic and X-ray crystallographic structural studies on some copper(II) complexes of the 6-methylpyridine-2-carboxaldehyde Schiff base of *S*-methyldithiocarbazate, **287**, 8
the optically active and racemic products of monocondensation of 1,2-diaminopropane and 2,4-pentadione as ligands and precursors for preparation of unsymmetrical Schiff bases and their nickel(II) complexes, **293**, 115
^{45}Sc-NMR spectroscopy
solution state chemistry of scandium(III): a study of scandium-crown ether complexes by ^{45}Sc-NMR spectroscopy, **284**, 71
S-coordinated complexes
terminal S-coordinated thiocyanate in a nickel(II) complex. X-ray structures of Ni(poph)(H_2O)(SCN)(NCS) and Ni(poqh)-$(H_2O)(NCS)_2$ (poph = 2-pyridinecarboxaldehyde 1-oxide 2′-pyridinylhydrazone, poqh = 2-pyridinecarboxaldehyde 1-oxide 2′-quinolynylhydrazone), **285**, 297
Selenocyanate complexes
spin-crossover in the $[Fe(abpt)_2(NCX)_2]$ (X = S, Se) system: structural, magnetic, calorimetric and photomagnetic studies, **291**, 279
Selenolate complexes
chelating and ligand-transfer studies of the complexes *cis*-$[Mn(CO)_4(SeR)_2]^-$: crystal structures of heterotrinuclear $[(CO)_4Mn(\mu\text{-}SeMe)_2Ni(\mu\text{-}SeMe)_2Mn(CO)_4]$ with a distorted square planar Ni^{II}–selenolate core and $[PPN]_2[Cl_2Pd(\mu\text{-}SePh)_2PdCl_2]$, **286**, 155
Seleno ligands
the preparation and coordination chemistry of $^iPr_2P$-(E)NHP(E′)$^iPr_2$ (E, E′ = Se; E = Se, E′ = S; E = S, E′ = O; E,E′ = O, **290**, 1
Semicarbazone complexes
structural characterization of a new ligand mode of 2,6-diacetylpyridine bis(semicarbazone), H_2daps, **292**, 121
Siderophore complexes
copper(II) and zinc(II) complexes of a macrocyclic bis-(amine–amide–hydroxamate) siderophore analogue. Formation constants and coordination chemistry, **284**, 20
Silacyclopropane
a facile synthesis of 1,3-disilacyclohexanes, **296**, 231
Silane complexes
reactions of H_2, silanes, and olefins with superelectrophilic cationic rhenium complexes: heterolytic cleavage of H_2 and relation to the structure and function of hydrogenases, **294**, 240
Silanolato complexes
triphenylsilanolato complexes of Group 4 metals: syntheses, X-ray crystal structures and catalytic cyclization reaction of 1,5-hexadiene, **286**, 14
Siloxo complexes
synthesis of organo(siloxo)platinum and -palladium complexes and preparation of supported nanoclusters by facile ligand reduction, **294**, 266
Silver complexes
a novel polymeric disilver(I) complex containing both macrocyclic- and linear-coordinate silver(I) centres: $[Ag_2(mn\text{-}15S_2O_3)]_x(ClO_4)_{2x}$ (mn-$15S_2O_3$ = maleonitrile-dithia[15]-crown-5), **285**, 305
electrospray mass spectrometric investigation of the relative ligand properties of EPh_3 ligands (E = P, As, Sb or Bi) towards Ag^+ and Cu^+, **290**, 213
new phosphino silver(I) derivatives of hydrotris(3-methyl-2-thioxo-1-imidazolyl)borate. X-ray crystal structure of tricyclohexylphosphinesilver(I)-hydrotris(3-methyl-2-thioxo-1-imidazolyl)borate, **285**, 81
silver(I) tetraiodoethylene complexes with twisted olefin moiety, **290**, 251
studies of the construction of coordination polymers using linear pyridyl-donor ligands, **292**, 231
synthesis and crystal structure of tetranuclear copper(I) and silver(I) complexes bridged by 2-amino-1,3,4-thiadiazole (atdz): $[Cu_4(atdz)_6](ClO_4)_4{\cdot}2CH_3OH$ and $[Ag_4(atdz)_6]$-$(ClO_4)_4$, **290**, 153

synthesis and electrochemical properties of a new C_{60}-phenanthroline derivative and its dimeric complex with silver, **292**, 34

synthesis and silver(I) coordination of N-functionalized aza-crown ethers with pendant aromatic carbocyclic or heterocyclic side-arms, **285**, 1

synthesis and structural characterization of binuclear silver(I) complexes bridged by three polydentate phosphine ligands, **284**, 278

Silver(I) complexes

stability constants of copper(I) and silver(I) complexes with open-chain, macrocyclic and -bicyclic aza-ligands in acetonitrile and comparison with results in dimethylsulfoxide, **286**, 160

Silyl complexes

reactions of bis(silyl)palladium(II) complexes with allyl halides. Synthesis of mono(silyl)palladium(II) halides and X-ray structure of *trans*-$PdCl(SiF_2Ph)(PMe_2Ph)_2$, **296**, 19

β-Silylethenylation

β-silylethenylation of phenol with silylethyne using gallium trichloride, **296**, 67

2-Silylmethyl-2-alkenals

synthesis of 2-silylmethyl-2-alkenals by rhodium-catalyzed interaction among 2-propynylamine, hydrosilane, and CO, **296**, 72

Single-crystal X-ray diffraction

characterization and crystal structure of copper(II) complex with betaine having a water-wheel-like structure, **295**, 121

Sodium borohydride

optically active β-ketoiminato cobalt(II) complexes as highly efficient catalysts in enantioselective borohydride reduction of ketones, imines, and α,β-unsaturated carboxylates, **296**, 86

Sodium complexes

18-membered tetraimino macrocyclic complexes containing furan groups: stacking of two macrocycles and characterization of organic derivatives, **284**, 149

Solid-state/solution ^{119}Sn NMR

studies in aryltin chemistry: Part XIV. Crystal and molecular structures of tris(*o*-methoxyphenyl)tin(IV) chloride, fluoride and oxide, **294**, 224

Solvatochromism

solvatochromism in ligand-field absorption bands and solvent-dependent deuteron nuclear magnetic resonance spectra of $[CrF_3$(tri(2-pyridyl)amine)]: implication of trigonal splitting for the first spin-allowed d–d band and evaluation of π bonding peculiarities, **295**, 84

Sorbitol complexes

potentiometric determination of the equilibrium constants for the formation of Fe(III)–sorbitol complexes, **293**, 206

Spectroelectrochemistry

reduction of the NO^+ ligand in the pentacyanonitrosylosmate(II) ion, **285**, 129

widely differing stabilities of molecule-bridged cyanodiiron(III, II) species in non-aqueous solvents, **291**, 66

Spiro compound

rhodium-catalyzed cyclic carbonylation of 2-phenylethynylbenzamide under water–gas shift reaction conditions, **296**, 195

Squarate complexes

crystal structures and magnetic properties of the squarate-O^1,O^n-bridged dinuclear copper(II) complexes $[Cu_2(phen)_4(C_4O_4)](CF_3SO_3)_2\cdot 3H_2O$ ($n=2$) and $[Cu_2(bipy)_4(C_4O_4)](CF_3SO_3)_2\cdot 6H_2O$ ($n=3$), **287**, 173

Stability constants

catalysis of hydrolysis of a phosphate diester by mono- and dinuclear macrocyclic zinc(II) complexes, **287**, 47

complexation of thorium(IV) by tris((2,3-dihydroxybenzylamino)ethyl)amine—a new strong chelating agent, **286**, 103

high coordination numbers of metal ions in chelate complexes: molecular adducts of scandium tris-α-diketonates in solution, **286**, 175

interaction of beryllium(II) in aqueous solution with bidentate ligands containing phosphonate groups, **285**, 39

solid and solution studies of ternary copper(II) complexes formed by heteroaromatic N bases and the anion of *N*,*N*-bis(2-hydroxyethyl)glycine (bicine), **285**, 326

stability constants of copper(I) and silver(I) complexes with open-chain, macrocyclic and -bicyclic aza-ligands in acetonitrile and comparison with results in dimethylsulfoxide, **286**, 160

synthesis and silver(I) coordination of N-functionalized aza-crown ethers with pendant aromatic carbocyclic or heterocyclic side-arms, **285**, 1

Stereochemistry

crystal structures and stereochemical properties of lanthanide(III) complexes with ethylenediamine-*N*,*N*,*N*′,*N*′-tetraacetate, **288**, 7

retention of optical activity during conversion of Λ-$[Ru(1,10\text{-phenanthroline})_3]^{2+}$ to Λ-$[Ru(1,10\text{-phenanthroline-5,6-dione})_3]^{2+}$ and Λ-$[Ru(dipyrido[a:3,2\text{-}h:2',3'\text{-}c\text{-}]\text{-phenazine})_3]^{2+}$, **293**, 37

Stereoselectivity

copper-mediated 1,6-, 1,8-, 1,10- and 1,12-addition and 1,5-substitution reactions in organic synthesis, **296**, 1

effects of electronic structure on chiral induction in outer-sphere electron transfer reactions of complexes with the C_3 symmetric ligand, 1,4,7-triazacyclononane-1,4,7-tris[2′(*R*)-2′-propionate](-3), **290**, 197

Steric effect

steric effect on the axial substitution of bis(dimethyl sulphoxide)phthalocyaninatoiron(II) by substituted imidazoles, **295**, 200

Stibine complexes

electrospray mass spectrometric investigation of the relative ligand properties of EPh_3 ligands (E = P, As, Sb or Bi) towards Ag^+ and Cu^+, **290**, 213

Structure-directing templates

hydrothermal assembly and structural characterization of two low-dimensional molybdenum oxide/2,2′-dipyridylamine hybrid materials, **295**, 222

Structure–reactivity

a semi-empirical quantum chemical study (PM3) of structure–reactivity relationships for mono- and binuclear bismuthines, **284**, 167

Substitution reactions

protonation and substitution reactions of Fe–S 'basket' clusters including extracted FeMo-cofactor of nitrogenase, **291**, 212

Substitutions

copper-mediated 1,6-, 1,8-, 1,10- and 1,12-addition and 1,5-substitution reactions in organic synthesis, **296**, 1

Sulfido complexes

cluster synthesis by the reactions of $[Cp'_2Fe_2S_4]$ with transition-metal complexes ($Cp' = C_5Me_5$, $1,3\text{-}C_5H_3(SiMe_3)_2$), **291**, 395

synthesis, structure, and redox properties of the sulfido bridged tetranuclear cluster complex $[(Ph_3P)_2N][Fe_3(CO)_9\{\mu_3\text{-}SFeCp(CO)_2\}]$, **286**, 142

Sulfinyl complexes

[*meso*- and *rac*-1,2-bis(4-fluorophenyl)ethylenediamine]chloro-[sulfinylbis(methane)-S]platinum(II) chloride new water soluble platinum complexes with high anti-breast cancer activities, **285**, 184

Sulfonylphosphane complexes
reactions of $Cp'(CO)_2MnPPh_2H$ with CH_3COCl and $CH_3S(O)_2Cl$ in THF/triethylamine: evidence of the first complex stabilization of a phosphorus homolog of the sulfonamides, **288**, 101
Sulfur and nitrogen ligand complexes
metal(II) complexes with monodentate S and N ligands as structural models for zinc–sulfur DNA-binding proteins, **285**, 262
Sulfur complexes
about the synthesis of the prismane $[NEt_4]_2[Fe_6S_6I_6]$, **284**, 296
lattice effects in the Mössbauer spectra of salts of $[Fe_4S_4\{S(CH_2)_nOH\}_4]^{2-}$. Crystal structures of $[PPh_4]_2[Fe_4S_4\{S(CH_2)_nOH\}_4]$ ($n = 2$, 3 and 4), **291**, 101
protonation and substitution reactions of Fe–S 'basket' clusters including extracted FeMo-cofactor of nitrogenase, **291**, 212
Sulfur ligand complexes
C–S, C–H, and N–H bond cleavage of heterocycles by a zero-valent iron complex, $Fe(N_2)(depe)_2$ [depe = 1,2-bis(diethylphosphino)ethane], **291**, 341
studies on Pt–S bonds. Complexes of methylsulfanyl- and methylsulfynil-acetate, benzoate and phenolate with the diethylenetriamineplatinum(II) moiety, **285**, 249
synthesis and characterization of ruthenium 'pincer' complexes containing sulfur atom donor ligands. The X-ray crystal structure of $[RuCl_2(\eta^3\text{-}NN'N)(SMe_2)]$ (NN'N = {2,6-$(Me_2NCH_2)_2C_5H_3N$}), **287**, 113
Superconductivity
B and B–C as interstitials in reduced rare earth halides, **289**, 174
Supercritical
synthesis of new Group III fluoride–ammonia adducts in supercritical ammonia: structures of $AlF_3(NH_3)_2$ and $InF_2(NH_2)(NH_3)$, **294**, 200
Supercritical fluids
manufacturing of specialty materials in supercritical fluid carbon dioxide, **294**, 214
Superoxide dismutase
synthesis and characterization of a ferric complex of the tripodal ligand tris(2-benzimidazolylmethyl)amine—a superoxide dismutase mimic, **290**, 21
Superoxide radical
synthesis and characterization of a ferric complex of the tripodal ligand tris(2-benzimidazolylmethyl)amine—a superoxide dismutase mimic, **290**, 21
Supramolecular chemistry
spectroscopic characterization of porphyrin supramolecular aggregates on poly-lysine and their application to quantitative DNA determination, **286**, 121
Supramolecular complexes
$NH_4TlI_4 \cdot 2(18c6) \cdot NH_3$, a new type of a potential double-shell lattice clathrate, **293**, 234
Surface chemistry
comparison of the surface chemical reactivity of hafnium diboride and hafnium, **289**, 191
Surfactant
synthetic reactions using organometallics in water. Aldol and allylation reactions catalyzed by Lewis acid–surfactant-combined catalysts/Brønsted acids systems, **296**, 158
Synchrotron X-ray
an approach to *megalo*-boranes. Mixed and multiple cluster fusions involving iridaborane and platinaborane cluster compounds. Crystal structure determinations by conventional and synchrotron methods, **289**, 95

Tantalum complexes
synthesis and structure of niobium and tantalum derivatives of bis(dicyclohexylphosphino)methane (dcpm), **288**, 35
Technetium complexes
metal–metal multiply-bonded complexes of technetium. Part 8. Synthesis and characterization of the α- and β-isomers of $Tc_2Cl_4(dppe)_2$, **288**, 69
study on the formation of mixed ligand oxorhenium and oxotechnetium complexes (SNS/S combination), **295**, 1
synthesis and characterization of five-coordinate rhenium(V) and technetium(V) mixed ligand bifunctional complexes carrying the SNS/S or the SNN/S donor atom set. Crystal structure of $ReO\{[(C_2H_5)_2NCH_2CH_2N(CH_2CH_2S)_2](p\text{-}H_2N\text{–}PhS)\}$ and $ReO\{[(CH_2)_4NCH_2CH_2NCH_2CH_2S](p\text{-}H_2N\text{–}PhS)\}$, **295**, 97
Template synthesis
template synthesis, crystal structure, and solution behavior of a hexaaza macrocyclic nickel(II) complex containing two *N*-aminoethyl pendant arms, **293**, 140
Terbium complexes
heteroleptic poly(pyrazol-1-yl)borate derivatives of the lanthanides. The syntheses of picolinate-*N*-oxide complexes and the X-ray crystal structure of $[Tb\{HB(C_3H_3N_2)_3\}_2(ONC_5H_4CO_2\text{-}2)]$, **285**, 283
Terdentate ligand complexes
palladium(II) allyl complexes with potentially terdentate ancillary ligands. Mechanism of allyl amination by piperidine, **293**, 44
Terdentate phosphine complexes
some new chain-like terdentate phosphines, their ruthenium(II) coordination chemistry and the activity of the cations $[Ru(MeCN)_3(PhP\{CH_2CH_2P(p\text{-}X\text{-}C_6H_4)_2\}_2)]^{2+}$ (X = H, F, Me and OMe) as acetalization catalysts, **290**, 64
Terimine ligands
the syntheses, spectra and structures of five-coordinate cobalt(II) complexes of pyrazolyl-containing ligands, **284**, 266
Ternary complexes
solid and solution studies of ternary copper(II) complexes formed by heteroaromatic N bases and the anion of *N*,*N*-bis(2-hydroxyethyl)glycine (bicine), **285**, 326
Terpyridine analog complexes
bdmpp complexes of Ru(II) as structural analogs for Ru(II)-2,2′:6′,2″-terpyridine species (where bdmpp is 2,6-bis(3,5-dimethyl-*N*-pyrazoyl)pyridine), **286**, 181
Terpyridine-derivative complexes
receptors based on 2,2′:6′,2″-terpyridine fragments containing peripheral amino groups, **292**, 28
Terpyridyl complexes
a simple route to bis(terpyridyl)transition metal oligomers, **284**, 76
Tetraborabicyclobutanes
novel tetraalkyltetraboranes of the type B_4R_4, $B_4H_2R_4$ and $B_4H_4R_4$, **289**, 58
Tetraboranes
novel tetraalkyltetraboranes of the type B_4R_4, $B_4H_2R_4$ and $B_4H_4R_4$, **289**, 58
Tetraboratetrahedranes
novel tetraalkyltetraboranes of the type B_4R_4, $B_4H_2R_4$ and $B_4H_4R_4$, **289**, 58
Tetrameric complexes
synthesis and crystal structure of $[Re(CO)_3(\mu\text{-}CH_3CO_2)(THF)]_4$, **284**, 304
Tetrameric compound
triethanolamine complexes of copper, **294**, 153
Tetranuclear complexes
synthesis and crystal structure of tetranuclear copper(I) and silver(I) complexes bridged by 2-amino-1,3,4-thiadiazole (atdz): $[Cu_4(atdz)_6](ClO_4)_4 \cdot 2CH_3OH$ and $[Ag_4(atdz)_6](ClO_4)_4$, **290**, 153

synthesis, crystal structure and magnetic behaviour of the new linear copper(II) tetranuclear compound $[Cu_4(\mu_3\text{-}AcO)_6(AcO)_2(DPQ)_2]$ (DPQ = 2,3-bis(2-pyridyl)quinoxaline), **286**, 62

Tetranuclear molybdenum complexes
synthesis and characterization of a series of new thiocarboxylate complexes of molybdenum(V), **284**, 223

Tetraphosphine complexes
synthesis, platinum(II) complexes and structural aspects of the new tetradentate phosphine *cis,trans,cis*-1,2,3,4-tetrakis-(diphenylphosphino)cyclobutane, **290**, 167

Tetraphosphoxane complexes
a *cyclo*-tetraphosphoxane cage complex of molybdenum(0) and palladium(II), **287**, 223

Tetrathiafulvalene
Rh-complex promoted synthesis of poly(arylacetylene) with pendant TTF units and formation of CT complexes of the polymer with π-acceptors, **296**, 278

Thallium complexes
structural diversity in thallium chemistry: III. The first structurally characterized examples of the pentabromothallate(III) anion in the 1,1,4,4-tetramethylpiperazinium and *N,N'*-diethyltriethylenediammonium salts contrasted with the mixed anions in the 1,1,3,3-tetramethylimidazolidinium salt, **285**, 122
structural diversity in thallium chemistry: II. Potential pentabromothallate(III) derivatives: examples of incipient bromide ion coordination?, **284**, 215

Theoretical calculations
cationic 16-electron half-sandwich ruthenium complexes containing asymmetric diamines: understanding the stability and reactivity of coordinatively unsaturated two-legged piano stool complexes, **286**, 114

Thermochemistry
solution thermochemical study of ligand substitution reactions of hybrid alkyl/fluoroalkoxy phosphorus ligands in the $L_2Fe(CO)_3$ system, **291**, 32

Thermodynamics
thermodynamic and structural study of inclusion complexes between trivalent lanthanide ions and native cyclodextrins, **293**, 53

Thermolysis
pyridines from allyl cobaloximes: a new decomposition pathway, **288**, 112

Thiadiazole complexes
synthesis and crystal structure of tetranuclear copper(I) and silver(I) complexes bridged by 2-amino-1,3,4-thiadiazole (atdz): $[Cu_4(atdz)_6](ClO_4)_4 \cdot 2CH_3OH$ and $[Ag_4(atdz)_6](ClO_4)_4$, **290**, 153

Thiamine complexes
interactions between thiamine and large anions. Crystal structures of (H-thiamine)$[Pt^{II}(SCN)_4] \cdot 3H_2O$ and (H-thiamine)$[Pt^{IV}(SCN)_6] \cdot H_2O$, **290**, 145

Thiocarboxylate complexes
oxygen atom transfer reaction involving oxomolybdenum complexes with sterically bulky thiocarboxylate ligands and biochemical interesting substrates in methanol at neutral pH, **285**, 25
synthesis and characterization of a series of new thiocarboxylate complexes of molybdenum(V), **284**, 223

Thiocyanate complexes
a new bidimensional compound containing (μ-thiocyanate)(bpy) copper(II) molecules: synthesis, crystal structure and magnetic properties of $[Cu(bpy)(NCS)_2]_n$ (bpy = 2,2'-bipyridyl), **286**, 108
preparation and spectroscopic properties of the complexes of mercuric thiocyanate with pyridine-2-thione and pyridine-2-carboxylic acid. Crystal and molecular structure of two polymorphs of $Hg(SCN)_2(C_5H_5NS)_2$, **285**, 208
spin-crossover in the $[Fe(abpt)_2(NCX)_2]$ (X = S, Se) system: structural, magnetic, calorimetric and photomagnetic studies, **291**, 279
terminal S-coordinated thiocyanate in a nickel(II) complex. X-ray structures of Ni(popb)(H_2O)(SCN)(NCS) and Ni(poqb)$(H_2O)(NCS)_2$ (popb = 2-pyridinecarboxaldehyde 1-oxide 2'-pyridinylhydrazone, poqh = 2-pyridinecarboxaldehyde 1-oxide 2'-quinolynylhydrazone), **285**, 297

Thiocyanato complexes
synthesis, crystal structure and magnetic properties of *catena*-poly[(2,2'-bipyridylcopper)-di-μ-thiocyanato] and bis(2,2'-bipyridyl)thiocyanatocopper(II) perchlorate, **286**, 169

Thioether complexes
a dinuclear copper(II) complex involving monoatomic *O*-carboxylate bridging and Cu–S(thioether) bonds: $[Cu(tda)(phen)]_2 \cdot H_2tda$ (tda = thiodiacetate, phen = phenanthroline), **286**, 74
copper(I) and copper(II) complexes of the bidentate imidazole/thioether ligand 1-methyl-2-(methylthiomethyl)-1*H*-benzimidazole, **287**, 204
synthesis and coordination chemistry of 1-phenyl-1-phospha-4,7-dithiacyclononane, **294**, 170
synthesis and structural characterisation of rhenium(V) oxo-thioether thiolate complexes with pendant ester or carboxylate groups, **285**, 76

Thiolate
effect of N–H···S amide–thiolate hydrogen bonding on bond lengths in rubredoxin models $[(CH_3)_3NCH_2CONH_2]_2[M(S_2\text{-}o\text{-}xyl)_2]$, **291**, 388

Thiolate complexes
chemistry of functionalized thiolates: syntheses and structures of neutral, monomeric $[Zn(SC_6H_4X\text{-}4)_2(N\text{–}N)]$, where X = H, NH_2 and OH, and (N–N) = nitrogen bases 2,2'-bpy, phen and tmen, **295**, 9
iron carbonyls with bulky thiolate ligands: crystal structures of $[Fe_2(CO)_6(\mu\text{-}SC_6H_2{}^iPr_3\text{-}2,4,6)_2]$ and $(C_6H_2{}^iPr_3\text{-}2,4,6)_2S_2$, **284**, 14
synthesis and characterization of oxorhenium-'3 + 1' mixed-thiolate complexes. Crystal and molecular structures of $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}(C_6H_4X\text{-}4\text{-}CH_2S)]$ (X = F, Cl, Br, OMe) and of the pendant thiolate compounds $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}\{\eta^1\text{-}SCH_2CH_2SCH_2CH_2SH\}]$ and $[ReO\{\eta^3\text{-}(SCH_2CH_2)_2S\}\{\eta^1\text{-}SCH_2CH(OH)CH(OH)CH_2SH\}]$, **284**, 252
synthesis and structural characterisation of rhenium(V) oxo-thioether thiolate complexes with pendant ester or carboxylate groups, **285**, 76
the effect of replacing carbonyl with acetonitrile on the reactivity of the bimetallic system $[Mo_2Cp_2(\mu\text{-}SMe)_2(L)_2]^+$ (L = CO, CH_3CN). Crystal structure of $[Mo_2Cp_2(\mu\text{-}SMe)_2(CH_3CN)_4](BF_4)_2$, **284**, 107

Thiolate ligands
a novel oligomerization versus metal chelation: synthesis and crystal structure of pentakis[di-μ-(diisopropylaminoethanethiolato)nickel], **294**, 83

Thiolato complexes
five-coordinate rhenium(III)-thiolato complexes: the structure of $[Re(SCH_2PhOCH_3)_3(PPh_3)_2]$, **290**, 247

Thiomorpholine complexes
synthesis and comparative reactivity of thiomorpholine–borane: aqueous hydrolysis and oxidation by hypochlorite, **290**, 80
transition metal complexes of 1,3-bis(thiomorpholino)propane: crystal structure and dynamic ^1H NMR study, **285**, 31

THE PAPER AND INK US
PUBLICATION MAY AFF
THE MICROFORM EDIT

Thione complexes
a new self-assembled copper(I) complex of 2-thioxo-5H-1,3-dithiolo[4,5-*b*][1,4]dithiepin-6(7*H*)-one showing three-dimensional supramolecular networks, **292**, 117
mercury(II) compounds with 1,3-imidazole-2-thione and its 1-methyl analogue. Preparative and NMR spectroscopic studies. The crystal structures of di-μ-iodo-bis[iodo(1,3-imidazolium-2-thiolato-*S*)mercury(II)], bis[bromo(1,3-imidazolium-2-thiolato-*S*)]mercury(II) and bis[μ-(1-*N*-methyl-1,3-imidazole-2-thiolato-*S*)]mercury(II), **294**, 35
Thiophenes
cleavage of substituents upon oxidative polymerization of substituted thiophenes, **287**, 232
Thiosemicarbazones complexes
structural study of the electrochemically synthesized binuclear complex bis{1-phenylglyoxal bis(3-piperidylthiosemicarbazone)zinc(II)}, **294**, 79
synthesis, structural characterization and biological activity of helicin thiosemicarbazone monohydrate and a copper(II) complex of salicylaldehyde thiosemicarbazone, **286**, 134
Thiotungstate complexes
crystallographic evidence for the coordination performance of $WO_2S_2^{2-}$. Synthesis, characterization and crystal structure of palladium complexes with mixed-ligand thiotungstates and 1,2-dithiolates: $[NBu_4]_2[PdWO_2S_2(C_2S_2)]$ and $[NBu_4]_2[PdWS_4(C_4N_2S_2)]$, **285**, 241
Thiourea complexes
main group metal halide complexes with sterically hindered thioureas XVI: The synthesis, characterization, and crystal structures of two new complexes of 1,3-dimethyl-2(3*H*)-imidazolethione with indium trihalides, **285**, 217
Thiourea oxides
structure and stability of aminoiminomethanesulfonic acid, **286**, 149
Thorium complexes
complexation of thorium(IV) by tris((2,3-dihydroxybenzylamino)ethyl)amine—a new strong chelating agent, **286**, 103
Three-dimensional complexes
magnetic iron phosphates with an open framework, **291**, 317
Tin complexes
1:1 adducts of triphenyltin chloride with oxovanadium(IV) tetradentate Schiff-base complexes, **293**, 147
multinuclear NMR studies of diphosphine, diphosphine–dioxide and diarsine complexes of tin(IV) halides. Structures of $[SnI_4\{o\text{-}C_6H_4(AsMe_2)_2\}]$ and $[SnI_4\{o\text{-}C_6H_4(P(O)Ph_2)_2\}]$, **288**, 142
synthesis of hexametallic $CH_2\{(\eta^5\text{-}C_5H_4)Fe(CO)_2SnPh_2(\eta^5\text{-}C_5H_5)Fe(CO)_2\}_2$, **291**, 252
Tin-119 NMR
multinuclear NMR studies of diphosphine, diphosphine–dioxide and diarsine complexes of tin(IV) halides. Structures of $[SnI_4\{o\text{-}C_6H_4(AsMe_2)_2\}]$ and $[SnI_4\{o\text{-}C_6H_4(P(O)Ph_2)_2\}]$, **288**, 142
Titania
'ship-in-bottle' synthesis of platinum carbonyl clusters in titania- or zirconia-modified mesoporous channels of FSM-16. Structural characterization and catalysis in CO hydrogenation, **294**, 281
Titanium complexes
molecular assembly of BINOL–Ti complexes into an active μ_3-oxo titanium catalyst, **296**, 267
monoindenyl titanium alkyl halides. The synthesis and molecular structures of $(\eta^5\text{-}C_9H_7)TiBr_3$, $(\eta^5\text{-}C_9H_7)Ti(CH_3)Br_2$, and $(\eta^5\text{-}C_9H_7)Ti(CH_3)Cl_2$, **292**, 220
reactions of $Ti(O\text{-}i\text{-}Pr)Cl_3$ with bidentate ligands containing hydroxy and methoxy groups. The crystal structure of $Ti(O\text{-}i\text{-}Pr)Cl_3(HOCH_2CH_2OCH_3)$ and the ^1H variable-temperature NMR studies of solution structures of $Ti(O\text{-}i\text{-}Pr)Cl_3(HOCH_2CH_2OCH_3)$ and $Ti(O\text{-}i\text{-}Pr)Cl_3(CH_3OCH_2\text{-}CH_2OCH_3)$, **285**, 254
synthesis and structure of a novel terminal imido complex of titanium, **292**, 105
synthesis and structures of $TiCl_3(2\text{-}OC_6H_4OCH_3)_2$ and $[TiCl_2(2\text{-}OC_6H_4OCH_3)(\mu\text{-}Cl)]_2$ and stepwise reactions of $[TiCl_2(2\text{-}OC_6H_4OCH_3)(\mu\text{-}Cl)]_2$ with 2-propanol, **293**, 12
Todorokite
a new synthetic route to todorokite-type manganese oxides, **294**, 123
Trans influence
platinum(II) complexes with the tridentate phosphino–thioether ligand bis(2-diphenylphosphinoethyl)sulfide, $(PPh_2CH_2\text{-}CH_2)_2S$. X-ray crystal structures of $[Pt\{(PPh_2CH_2\text{-}CH_2)_2S\}Cl]ClO_4$ and $[Pt\{(PPh_2CH_2CH_2)_2S\}I]I$, **295**, 153
Transition metal
synthesis and formation of metal complexes of 4-alkynyl and 4-cyano-2,6-diisopropylphenylisocyanides, **284**, 205
Transition metal complexes
a large C_6D_6 solvent effect on the ^1H NMR shifts of pyrazines coordinated to the $M(CO)_5$ series (M = W, Mo, Cr), **285**, 157
electrospray mass spectrometric analysis of transition-metal halide complexes, **294**, 183
^1H NMR study of new heterodinuclear macrocyclic complexes, **285**, 233
syntheses, crystal structures and coordination compounds of some 2-hetarylcyanoximes, **284**, 85
structural trends in a series of divalent transition metal triazole complexes, **293**, 61
transition metal complexes of 1,3-bis(thiomorpholino)propane: crystal structure and dynamic ^1H NMR study, **285**, 31
Triamine complexes
base hydrolysis of $[Co(triamine)(diamine)Cl]^{2+}$ and epimerization of $[Co(triamine)(diamine)OH]^{2+}$: studies on resolved cobalt(III) complexes of 3-aza-hexane-1,6-diamine (2,3-tri), **284**, 37
synthesis, vibrational study, crystal structure and density functional calculations of $[Ni(dien)_2]^{2+}$ complexes. Configurational and conformational study of $[M(dien)_2]^{n+}$ complexes, **290**, 57
Triazine complexes
absence of 1,3-metallotropic shifts in $M(CO)_5$ complexes of pyrimidine and 1,3,5-triazine (M = W, Mo, Cr), **292**, 225
Triazole complexes
structural trends in a series of divalent transition metal triazole complexes, **293**, 61
Tricarbonylchromium complexes
tricarbonylchromium complexes with methoxy substituted benzocyclobutenone and benzocyclobutenedione ligands, **296**, 150
Tridentate ligand complexes
copper(II) complexes with (2-hydroxybenzyl-2-pyridylmethyl)-amine–Hbpa: syntheses, characterization and crystal structures of the ligand and $[Cu(II)(Hbpa)_2](ClO_4)_2\cdot 2H_2O$, **290**, 207
Triethanolamine complexes
triethanolamine complexes of copper, **294**, 153
Trigonal splitting
solvatochromism in ligand-field absorption bands and solvent-dependent deuteron nuclear magnetic resonance spectra of $[CrF_3(tri(2\text{-}pyridyl)amine)]$: implication of trigonal splitting for the first spin-allowed d–d band and evaluation of π bonding peculiarities, **295**, 84

IK USED IN THE ORIGINAL
AFFECT THE QUALITY OF
EDITION.

Triiodide anions
polyiodides of transition metal trischelate cations: syntheses, structures, and spectral and electrical conductivity studies of $[Mn(phen)_3](I_3)_2$ and $[Mn(bpy)_3](I_3)_{1.5}(I_8)_{0.25}$, **284**, 158
Trimesate complexes
synthesis and crystal structure of two new rare earth trimesate complexes: $ErTMA(H_2O)_5{\cdot}3.5H_2O$ and $YTMA(H_2O)_6$, **284**, 139
Trinuclear complexes
synthesis, structure and magnetic properties of oxo-centered trinuclear manganese complex $[Mn_3O(O_2CC_3H_7)_6(C_5H_5N)_3]ClO_4$, **294**, 109
synthesis, studies and molecular structure of trinuclear cyanide-bridged copper–iron complexes, **292**, 157
trinuclear vanadium dialkyldithiocarbamate complexes. Synthesis, structure and spectroscopic characterization of $[Et_4N][V_3(\mu_3\text{-}S)(\eta^2\text{-}\mu\text{-}S_2)_3(Et_2dtc)_3]\cdot 2CH_3CN$, **286**, 7
Trinuclear manganese(II,III,II)
synthesis, structure and magnetic properties of a linear trimanganese(II,III,II) complex with hydroxo and 1,2,4-triazole as bridging ligands, **295**, 229
Triol complexes
ab initio calculations and X-ray structure determinations of dichloro(*O*-*O*′-*O*″-1,3,5-cyclohexanetriol)copper(II) and bis(*O*-*O*′-*O*″-1,3,5-hexanetriol)nickel(II), **284**, 284
Triphenylphosphine oxide complexes
complexes of lanthanide dithionates 2. Electrospray mass spectrometry of $Ln_2(S_2O_6)_3$ and their Ph_3PO complexes; the X-ray crystal structure of $Pr_2(S_2O_6)_3(Ph_3PO)_6(H_2O)_6$, **295**, 146
Triple-decker cation complexes
non-linear optical absorption of $[\{Ni(mnt)_2\}_2]^{2-}$ and its unique counter ion, a disodium triple-decker dication $[Na_2(benzo\text{-}15\text{-}crown\text{-}5)_3(H_2O)_2]^{2+}$, **293**, 89
Tripodal amine complexes
a new carbonate bridged dinuclear zinc complex with tripodal amine ligands, **288**, 206
synthesis and characterization of a ferric complex of the tripodal ligand tris(2-benzimidazolylmethyl)amine—a superoxide dismutase mimic, **290**, 21
Tripodal ligand complexes
mononuclear lanthanide complexes of tripodal ligands: synthesis and spectroscopic studies, **285**, 49
novel iron(III) complexes with imidazole containing tripodal ligands as model systems for catechol dioxygenases, **291**, 289
Tripodal ligands
a two-dimensional network constructed via 72-membered heart-shaped macrocycles, a copper(II) complex with 1,3,5-tris(imidazol-1-ylmethyl)-2,4,6-trimethylbenzene and diethylenetriamine ligands, **295**, 129
the ruthenium and iridium coordination chemistry of the tripodal ligand $CH_3C\{CH_2P(m\text{-}CF_3\text{-}C_6H_4)_2\}_3$, **293**, 70
Tris–catechol complexes
thermodynamic and kinetic studies of the sulfonated derivative of the iron chelator TRENCAM, an analog of enterobactin, **291**, 148
Tris(indazol-1-yl)borate complexes
coordination geometries for palladium and platinum: theoretical studies and the synthesis and structure of tris(indazol-1-yl)borate complexes $MMe_3\{(ind)_3BH\}$, **287**, 27
Tris(*o*-methoxyphenyl)tin compounds
studies in aryltin chemistry: Part XIV. Crystal and molecular structures of tris(*o*-methoxyphenyl)tin(IV) chloride, fluoride and oxide, **294**, 224
T-state effect
steric effect on the axial substitution of bis(dimethyl sulphoxide)phthalocyaninatoiron(II) by substituted imidazoles, **295**, 200
Tungstate
kinetics of the molybdate and tungstate catalyzed oxidation of iodide by hydrogen peroxide, **288**, 229
Tungstate complexes
cluster synthesis by the reactions of $[Cp'_2Fe_2S_4]$ with transition-metal complexes ($Cp' = C_5Me_5$, $1,3\text{-}C_5H_3(SiMe_3)_2$), **291**, 395
Tungsten complexes
ligand substitution kinetics in $M(CO)_4$ ($\eta^{2\text{-}2}$-1,5-cyclooctadiene) complexes (M = Cr, Mo, W) — substitution of 1,5-cyclooctadiene by bis(diphenylphosphino)alkanes, **284**, 229
redox active organoiron alkynyl centers with a coordination site: an approach toward molecular plugs. Complexation reactions with $(THF)W(CO)_5$ and $(PhCN)_2MCl_2$ (M = Pd, Pt), **291**, 403
ring-opening polymerization of norbornene by η-cycloheptatrienyl molybdenum and tungsten complexes, **288**, 226
syntheses, crystal structures and nonlinear optical properties of heteronuclear clusters $[MS_4Cu_4(SCN)_2(NC_5H_5)_6]$ (M = W, Mo), **294**, 73
synthesis and characterization of long chain acyl and alkyl complexes of the types $[(\eta^5\text{-}C_5R'_5)Fe(CO)_2\{C(O)R\}]$ (R′ = H, CH_3), $[CpW(CO)_3\{C(O)R\}]$ and $[CpW(CO)_3R]$ (R = *n*-alkyl), **286**, 221
synthesis and characterization of the triethylphosphine adduct of bis(2,4-dimethylpentadienyl)tungsten, $[W(2,4\text{-}C_7H_{11})_2(PEt_3)]$, a complex with one η^5-**U** and one η^5-**S** pentadienyl ligand, **288**, 154
synthesis and structures of tungsten ferrocenyldiazoalkane complexes derived from a dinitrogen complex. Part 58. Preparation and properties of molybdenum and tungsten dinitrogen complexes, **285**, 336
synthesis of a dinuclear tungsten complex, $W_2(\mu\text{-}H)(\mu\text{-}OC_2H_5)(\mu\text{-}O_2CC_6H_5)(\eta^1\text{-}OC_2H_5)_2Cl_2(\eta^2\text{-}P,P'\text{-dppp})$ where dppp is 1,3 bis(diphenylphosphino) propane, **284**, 303
synthesis of *N*-allyl-2,3-dihydrobenzoxazol-2-ylidene complexes of chromium(0) and tungsten(0), **288**, 47
Two-dimensional network
a two-dimensional network constructed via 72-membered heart-shaped macrocycles, a copper(II) complex with 1,3,5-tris(imidazol-1-ylmethyl)-2,4,6-trimethylbenzene and diethylenetriamine ligands, **295**, 129

Unsaturated carbonyl group complexes
transformation of the carboxyl group adjacent to the $(\eta^4$-diene)$Fe(CO)_3$ moiety: a convenient route to $(\eta^4$-diene$)Fe(CO)_3$ complexes containing α,β- and β,γ-unsaturated carbonyl groups, and 1,3-dicarbonyl groups, **291**, 231
UPS
formation of FeSi and $FeSi_2$ films from *cis*-$Fe(SiCl_3)_2(CO)_4$ by MOCVD – precursor versus substrate control, **291**, 380
Uranium complexes
complexation of uranyl ion by tetrahexylmalonamides: an equilibrium modeling and infrared spectroscopic study, **293**, 195
Uranyl ion
photooxidation of water and isobutane in aqueous uranyl solutions, **285**, 322
Urea
crystal structures of two new heptamolybdates and of a pyrazole incorporating a γ-octamolybdate anion, **295**, 106

Valence-force-fields
lattice vibrations of the icosahedral solid boron arsenide, $B_{12}As_2$, **289**, 198
Vanadium complexes
1:1 adducts of triphenyltin chloride with oxovanadium(IV) tetradentate Schiff-base complexes, **293**, 147

preparation and characterisation of new oxovanadium(IV) Schiff base complexes derived from amino acids and aromatic *o*-hydroxyaldehydes, **293**, 1

synthesis, structure and EPR spectroscopy of a vanadium(IV)–amide metallacyclic complex, **285**, 134

trinuclear vanadium dialkyldithiocarbamate complexes. Synthesis, structure and spectroscopic characterization of $[Et_4N][V_3(\mu_3\text{-}S)(\eta^2\text{-}\mu\text{-}S_2)_3(Et_2dtc)_3]\cdot 2CH_3CN$, **286**, 7

vanadium phenoxide complexes with oxide, nitride or hydrazide co-ligands: preparation and crystal structures of $[V(OC_6H_3Pr^i_2\text{-}2,6)_3NLi(C_4H_8O)_3]$, $[\{VO_2(OC_6H_3Pr^i_2\text{-}2,6)_2\}_2\text{-}\{\mu\text{-}Li(C_4H_8O)_2\}_2]$ and $[V(NNMe_2)(OC_6H_3Pr^i_2\text{-}2,6)_3]$, **285**, 178

Vanadyl sulfate

hydrothermal synthesis and characterization of a new vanadyl sulfate compound: crystal structure of $[V(IV)OSO_4\text{-}(H_2O)_4]\cdot SO_4\cdot[H_2N(C_2H_4)_2NH_2]$, **292**, 112

Vibrational spectra

alkyl derivatives of $[B_6H_6]^{2-}$: NMR and vibrational spectra and crystal structure of $(Ph_4P)[B_6H_6CH_2Ph]$, **289**, 70

Vitamin B_{12} derivatives

vitamin B_{12}–transport protein interaction: electrochemistry of aquo- and glutathionyl-cobalamins adsorbed on carbon electrodes; role of the nucleotide chain, **292**, 204

Water soluble complexes

structural, spectroscopic and catalytic properties of water-soluble hydride rhodium complexes $[RhH(Rtpa^+I^-)_4]H_2O$ (R = Me, Et), **293**, 110

water soluble platinum(II) and palladium(II) complexes of alkyl sulfonated phosphines, **290**, 189

Water-wheel-like structures

characterization and crystal structure of copper(II) complex with betaine having a water-wheel-like structure, **295**, 121

X-ray absorption spectroscopy

novel iron(III) complexes with imidazole containing tripodal ligands as model systems for catechol dioxygenases, **291**, 289

X-ray crystal structures

bonding of the β-pinene allyl, $\eta^3\text{-}C_{10}H_{15}$ to Pd(II): X-ray structure of $[Pd(\eta^3\text{-}C_{10}H_{15})(Josiphos)]CF_3SO_3$, **296**, 246

X-ray crystal structures of copper(II) nitrate complexes of two isomeric N,O-chelating ligands, **284**, 273

X-ray diffraction

structures of 1,3-bis(diphenylphosphino)propane platinum(II) alditolate complexes, **285**, 10

X-ray structures

synthesis, structures, and linkage isomerism of (allylbenzylmalonate)platinum(II) complexes, **295**, 214

Ylide complexes

synthesis of gold ferrocenyl substituted ylides and methanides: crystal structure of $[FcCH(AuPPh_3)PPh_3]ClO_4$ $[Fc = (\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4)]$, **291**, 60

Zeolites

hydrothermal synthesis of microporous materials with the direct incorporation of porphyrin molecules, **294**, 179

Zinc complexes

adduct formation between zinc and cadmium oxydiacetates and imidazol. Crystal structures of $[Zn(Him)_2(oda)]_n$ and $[Cd_2(Him)_4(oda)_3(H_2O)]_2\cdot 6H_2O$ (Him = imidazol, oda = oxydiacetate), **284**, 49

a new carbonate bridged dinuclear zinc complex with tripodal amine ligands, **288**, 206

catalysis of hydrolysis of a phosphate diester by mono- and dinuclear macrocyclic zinc(II) complexes, **287**, 47

chemistry of functionalized thiolates: syntheses and structures of neutral, monomeric $[Zn(SC_6H_4X\text{-}4)_2(N\text{–}N)]$, where X = H, NH_2 and OH, and (N–N) = nitrogen bases 2,2′-bpy, phen and tmen, **295**, 9

highly enantioselective catalytic isopropenylation of aldehydes using diisopropenylzinc, **296**, 33

metal(II) complexes with monodentate S and N ligands as structural models for zinc–sulfur DNA-binding proteins, **285**, 262

molecular recognition of dipeptides. Catalysis of deuteration and hydrolysis of glycylglycine by dinuclear OBISDIEN Zn(II) complexes, **287**, 134

some complexes derived from zinc salicylate or 3,5-di-tert-butylsalicylate. The crystal structure of (2,2′-bipyridyl)-(methanol)(*O*-salicylato)(*O,O′*-salicylato)zinc, **287**, 89

structural study of the electrochemically synthesized binuclear complex bis{1-phenylglyoxal bis(3-piperidylthiosemicarbazone)zinc(II)}, **294**, 79

structure and properties of a novel OH^- bridged dimetal helical complex with 6,6‴-dimethyl-2,2′:6′,2″:6″,2‴:6‴,2⁗-quinquepyridine ligand, **290**, 127

the kinetics of the sequential hydrolysis reactions of the *xxm*-$[CoCl_2(dien)(py)]^+$ cation and the crystal structure of the $ZnCl_4^{2-}\cdot 2H_2O$ salt, **292**, 198

the reaction of triorganophosphorus diiodides, R_3PI_2, with zinc metal powder; dependency of product on R; the X-ray crystal structures of dimeric $\{ZnI_2[P(NMe_2)_3]\}_2$ and monomeric $ZnI_2(PPh_2Me)_2$, **292**, 213

Zintl complexes

synthesis and structural characterization of two new Zintl $[M_7As_3Q_{13}]^{4-}$ anions (M = Cu, Ag; Q = Se, Te) containing distorted cubane $[M_3Q]$ clusters, **288**, 1

Zirconium complexes

variations in capping the $Zr_6O_4(OH)_4$ cluster core: X-ray structure analyses of $[Zr_6(OH)_4O_4(OOC\text{–}CH\text{=}CH_2)_{10}]_2(\mu\text{-}OOC\text{–}CH\text{=}CH_2)_4$ and $Zr_6(OH)_4O_4(OOCR)_{12}(PrOH)$ (R = Ph, CMe = CH_2), **284**, 1